

NIRS-RSD-52

**RADIOACTIVITY
SURVEY DATA
in Japan**

NUMBER 52

Mar. 1980

National Institute of Radiological Sciences

Chiba, Japan

Radioactivity Survey Data in Japan Number 52

Mar. 1980

Contents

	Page
Environmental and Dietary Materials	1
<i>(Japan Chemical Analysis Center)</i>	
1. Collection and pretreatment of samples	1
2. Preparation of samples for analysis	3
3. Separation of strontium-90 and cesium-137	3
4. Determination of stable strontium, calcium and potassium	4
5. Counting	4
6. Results	5
(1)-1 Strontium-90 and Cesium-137 in Rain and dry fallout (for domestic program)	5
-2 Strontium-90 and Cesium-137 in Rain and dry fallout (for WHO program)	13
(2) Strontium-90 and Cesium-137 in Airborne dust	17
(3) Strontium-90 and Cesium-137 in Service water	19
(4) Strontium-90 and Cesium-137 in Soil	23
(5) Strontium-90 and Cesium-137 in Sea water	26
(6) Strontium-90 and Cesium-137 in Sea sediments	28
(7) Strontium-90 and Cesium-137 in Total diet	30
(8) Strontium-90 and Cesium-137 in Rice	33
(9)-1 Strontium-90 and Cesium-137 in Milk (producing districts for WHO program)	35
-2 Strontium-90 and Cesium-137 in Milk (producing districts for domestic program)	38
-3 Strontium-90 and Cesium-137 in Milk (consuming districts)	38
-4 Strontium-90 and Cesium-137 in Milk (powdered milk)	42
(10) Strontium-90 and Cesium-137 in Vegetables	42
(11) Strontium-90 and Cesium-137 in Tea (Japanese tea)	46
(12) Strontium-90 and Cesium-137 in Sea fish	47
(13) Strontium-90 and Cesium-137 in Freshwater fish	51
(14) Strontium-90 and Cesium-137 in Shellfish	53
(15) Strontium-90 and Cesium-137 in Seaweeds	55

Environmental and Dietary Materials*

(Japan Chemical Analysis Center)

1. Collection and pretreatment of samples

(1) Rain and dry fallout

Rain and dry fallout was collected monthly on a sampling tray, approximately 5000 cm² in area, which was filled with water to a depth of 1 cm at the beginning of every month.

The sample was filtered after strontium and cesium carriers were added. The tray was washed with 5ℓ of distilled water and the washing was combined to the filtrate. The sample was passed through a cation exchange column (500 mℓ of Dowex 50W X8, 50 ~ 100 mesh, Na form) at a rate of 80 mℓ/min.

(2) Airborne dust

Airborne dust was collected by an electrostatic precipitator or a filter air sampler for every three months at a rate of more than 3000 m³ per month. The sampling was done 1 to 1.5 meters above the ground.

(3) Service water and freshwater

Service water, 100ℓ each, was collected at an intake of the water-treatment plant and at the tap after water was left running for five minutes. Water, to which added carriers of strontium and cesium immediately after sampling, was vigorously stirred and filtered. The subsequent process was the same as that described in the section (1). Freshwater was treated in the same way as the service water.

(4) Soil

Soil was collected from the location in the spacious and flat area without past disturbance on the surface caused by duststorms, inflow and outflow due to precipitation, and so on. Any places located under trees in a forest, in a stony area or inside of river banks were avoided. Soil was taken from two layers of different depths, 0 ~ 5 cm and 5 ~ 20 cm. In the course of air-drying, lumps were crushed by hand, and roots of plants and pebbles were removed. The soil was then passed through a 2 mm sieve to remove small gravels.

(5) Sea water

Sea water was collected at the fixed stations where the effect of terrestrial fresh water from rivers was expected to be negligibly small. A special consideration was also given to weather conditions. The sampling was carried out when there was no rainfall for the last few days. To prevent contamination, water samples were collected at the bow of a sampling boat just before she stood still by scooping surface water using a polyethylene bucket. Immediately after the collection, the samples were acidified to a pH lower than 3 by adding concentrated hydrochloric acid in a ratio of 1 mℓ to 1 ℓ of sea water, and then stored in 20 ℓ polyethylene containers. The sampling equipments as well as containers were thoroughly rinsed with dilute hydrochloric acid and then with distilled water before use. Two hundred milliliters of sea water was also collected at the same stations for the determination of chlorinity.

(6) Sea sediments

Sediment was collected in the same area as that for the sea water sample, taking the following criteria into account:

- a. The depth of water exceeds 1 m at low tide.
- b. No significant sedimental movement is observed in the vicinity of concern.
- c. Mud, silt and fine sand are preferable.

A conventional sediment sampling device was used for collecting the top few centimeters of surface sediment. Approximately 4 kg of the sample in wet weight was spread on a large porcelain dish and dried in an electric oven at 105 to 110 °C to a constant weight.

(7) Total diet

A full one day ordinary diet including three meals, water, tea and other in-between snacks for five persons was collected as a sample of "total diet". The sample in a large stainless steel pan was carbonized carefully by direct application of gas flame, and was transferred to a porcelain dish and then ashed at 500°C in an electric muffle furnace.

(8) Rice

Polished rice was collected in producing districts at the harvest and in consuming areas when new crops were first put on sale. The sample was carbonized and ashed in a porcelain dish.

* Samples were sent to the Center from 32 contracted prefectures.

(9) Milk

Raw milk was collected in producing districts and commercial milk was purchased in consuming districts. Milk in a stainless steel pan or a porcelain dish was evaporated to dryness followed by carbonization and ashing.

(10) Vegetables

Spinach and Japanese radish were selected as the representatives for leaf vegetables and for non-starch roots, respectively. After removing soil, the edible part of vegetable sample was dried and carbonized in a stainless steel pan or a porcelain dish.

(11) Tea

Five hundred grams of manufactured green tea was collected, carbonized and ashed in a stainless steel pan or a porcelain dish.

(12) Fish, shellfish and seaweeds

a. Sea fish and freshwater fish

Fish was rinsed with water and blotted with a filter paper. Only the edible part was used in case of larger sized fish, and the whole part was used in case of smaller ones. Each sample was weighed and placed in a stainless steel pan or a porcelain dish. After carbonized, the sample was ashed in an electric muffle furnace.

b. Shellfish

Approximately 4 kg of shellfish including the shells was collected or purchased. After removing the shells, it was treated in the same way as that for the sea fish.

c. Seaweeds

Edible seaweeds were collected and rinsed with water to remove sand and other adhering matters on the surface. These were removed of excess water, weighed, dried and ashed.

Table 1 shows details of sample collection.

Table 1 Details of sample collection

Sample	Frequency of sampling	Quantity of sample
= Environmental materials =		
(1) Rain and dry fallout		
1 for domestic program	monthly	
2 for WHO program	monthly	
(2) Airborne dust	quarterly	>3000 m ³ /month
(3) Service water and freshwater		
1 Service water (source water)	semiyearly (June and December)	100 ℓ
2 Service water (tap water)	semiyearly (June and December)	100 ℓ
3 Freshwater	yearly (fishing season)	100 ℓ
(4) Soil		
1 0 ~ 5 cm	yearly (June or July)	4 kg
2 5 ~ 20 cm	yearly (June or July)	4 kg
(5) Sea water	yearly (July or August)	40 ℓ
(6) Sea sediments	yearly (July or August)	4 kg
= Dietary materials =		
(7) Total diet	semiyearly (June, November or December)	daily amount for 5 person
(8) Rice		
1 Producing districts	yearly (harvesting season)	5 kg (polished rice)
2 Consuming districts	yearly (harvesting season)	5 kg (polished rice)
(9) Milk		
1 Producing districts for WHO program	quarterly (February, May, August and November)	3 ℓ
2 Producing districts for domestic program	semiyearly (February and August)	3 ℓ

Sample	Frequency of sampling	Quantity of sample
3 Consuming districts	semiyearly (February and August)	3 ℓ
4 Powdered milk	semiyearly (April and October)	2 ~ 3 kg
(10) Vegetables		
1 Producing districts	yearly (harvesting season)	4 kg
2 Consuming districts	yearly (harvesting season)	4 kg
(11) Tea	yearly (the first harvesting season)	500 g (manufactured tea)
(12) Fish, shellfish, and seaweeds		
1 Sea fish	yearly (fishing season)	4 kg
2 Freshwater fish	yearly (fishing season)	4 kg
3 Shellfish	yearly (fishing season)	4 kg
4 Seaweeds	yearly (fishing season)	2 ~ 3 kg

2. Preparation of samples for analysis

(1) Rain, service water and freshwater

Strontium and cesium were eluted with hydrochloric acid from the cation exchange column. The residue of rain sample on the filter paper was ashed in an electric muffle furnace and the ash was dissolved in hydrochloric acid. The insoluble part was filtered and washed. The filtrate and the washings were combined to the previous eluate and used for radiochemical analysis.

(2) Soil

Air-dried soil was passed through a 20 mesh sieve. The sieved sample was heated, in the presence of strontium and cesium carriers, together with sodium hydroxide. The sample was then heated with hydrochloric acid and the insoluble part was filtered and washed. The combined solution of the filtrate and washings was used for radiochemical analysis.

(3) Sea sediments

After removal of pebbles, shells and other foreign matters, the sediment sample was dried in a hot-air oven and ground finely with a mortar. The sample was passed through a 20 mesh sieve. The further preparation of the sample was the same as that described in the section 2-(2).

(4) Rice

The ashed sample was pulverized with a porcelain mortar and passed through a 42 mesh sieve. The sieved sample to which both strontium and cesium carriers were added, was digested with hydrochloric

acid by heating. After the sample was heated again with nitric acid to dryness, strontium and cesium were extracted with hydrochloric acid and water. The insoluble part was filtered and washed. The filtrate and washings were combined for subsequent radiochemical analysis.

(5) Airborne dust, diet, milk, vegetable, fish and shellfish, seaweeds, tea, and others

These ashed samples were treated with the same procedure as that described in the section 2-(4).

3. Separation of strontium-90 and cesium-137

(1) Strontium-90

Sample solutions, prepared as in the foregoing sections 2-(1) through 2-(5), were neutralized with sodium hydroxide. After sodium carbonate was added, the precipitate of strontium and calcium carbonates was separated. The supernatant solution was retained for cesium-137 determination. The carbonates were dissolved in hydrochloric acid and calcium and strontium were precipitated as oxalates. The precipitate was dissolved in nitric acid and strontium was separated from calcium by successive fuming nitric acid separations. Iron scavenge was made after addition of ferric iron carrier followed by barium chromate separation after addition of barium carrier to remove radium, its daughters and lead. Strontium was recovered as carbonate, and the precipitate was dried and weighed to determine strontium recovery. The strontium carbonate was dissolved in hydrochloric acid and the iron carrier was added. The solution was allowed to stand

for two weeks for strontium-90 and yttrium-90 to attain equilibrium. The yttrium-90 was coprecipitated with ferric hydroxide and the precipitate was filtered off, washed and counted.

(2) Cesium-137

The supernatant separated from the strontium fraction in the solution was acidified with hydrochloric acid. While stirring the solution, cesium was adsorbed on ammonium molybdophosphate.

After filtered off and washed with dilute nitric acid, the precipitate was dissolved in 2.5N sodium hydroxide solution. Ammonia was removed completely from the solution by boiling. The solution was adjusted to pH 8.2 with hydrochloric acid and allowed to cool. Molybdenum hydroxide which came out in the solution, was filtered off and washed with water. In such circumstance that contamination by rubidium-87 was not negligible for the measurement of cesium-137, the following ion-exchange procedure was applied. A fixed amount of ferric chloride solution was added to the solution dissolved with 2.5N sodium hydroxide. Ammonia and molybdenum hydroxide were removed as described above. Ethylenediaminetetraacetic acid tetrasodium salt was added to the filtrate and washings. Cesium and rubidium were adsorbed on a cation exchange resin. Cesium was separated from rubidium by eluting with hydrochloric acid.

To this eluate or the filtrate and washings after removing molybdenum hydroxide, chloroplatinic acid solution was added to precipitate cesium. The precipitate was filtered onto a tared paper in a demountable filter and washed with water and then ethanol. After fixing the filter paper on a tared planchette and drying

it, the chemical yield of cesium was determined by weighing the precipitate with the planchette. Radioactivity from cesium-137 was measured for this precipitate.

4. Determination of stable strontium, calcium and potassium

A weighed amount of soil or sea sediment was treated under heating with sodium hydroxide and then with hydrochloric acid for extraction. A weighed aliquot of ashed samples of total diet, vegetables, milk, fish, shellfish or seaweeds was digested using hydrochloric acid or nitric acid, hydrofluoric acid being used when necessary. The extract was made up to an appropriate volume with dilute hydrochloric acid. The sample solution was analyzed for calcium by titration with standard potassium permanganate solution after separating calcium as oxalate. Atomic absorption spectroscopy was applied when appropriate. Stable strontium and potassium were determined by atomic absorption and flame emission spectrometry, respectively.

5. Counting

After the radiochemical separation, the mounted precipitates were counted for activity using low background beta counters normally for 60 min. Net sample counting rates were corrected for counter efficiency, recovery, self-absorption and decay to obtain the content of strontium-90 and cesium-137 radio activity per sample aliquot. From the results, concentrations of these nuclides in the original samples were calculated.

6. Results

(1)-1 Strontium-90 and Cesium-137 in Rain and dry fallout (for domestic program) (from Jan. 1979 to Dec. 1979)

— continued from No. 51 of this publication —

Table 1-1: Strontium-90 and Cesium-137 in Rain and dry fallout

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
January, 1979				
Sapporo, HOKKAIDO	37	140.0	0.031 ± 0.0016	0.053 ± 0.0017
Aomori, AOMORI	28	97.5	0.041 ± 0.0018	0.058 ± 0.0018
Sendai, MIYAGI	25	87.6	0.009 ± 0.0009	0.018 ± 0.0011
Yamagata, YAMAGATA	35	48.19	0.020 ± 0.0013	0.031 ± 0.0014
Ohkuma, FUKUSHIMA	29	8.0	0.007 ± 0.0009	0.007 ± 0.0009
Mito, IBARAGI	28	62.0	0.008 ± 0.0008	0.016 ± 0.0011
Shinjuku, TOKYO	31	66.5	0.009 ± 0.0011	0.018 ± 0.0011
Yokohama, KANAGAWA	29	28.8	0.009 ± 0.0010	0.019 ± 0.0011
Fukui, FUKUI	28	245.8	0.070 ± 0.0026	0.13 ± 0.003
Shizuoka, SHIZUOKA	27	67.0	0.010 ± 0.0010	0.020 ± 0.0012
Nagoya, AICHI	29	72.2	0.008 ± 0.0009	0.013 ± 0.0010
Kyoto, KYOTO	27	66.7	0.012 ± 0.0011	0.017 ± 0.0011
Kobe, HYOGO	29	46.4	0.009 ± 0.0011	0.013 ± 0.0011
Tottori, TOTTORI	28	102.50	0.041 ± 0.0017	0.059 ± 0.0018
Matsue, SHIMANE	32	96.5	0.043 ± 0.0018	0.073 ± 0.0020
Matsuyama, EHIME	36	74.5	0.021 ± 0.0014	0.033 ± 0.0014
Dazaifu, FUKUOKA	27	42.6	0.014 ± 0.0012	0.020 ± 0.0012
Saga, SAGA	34	54.5	0.011 ± 0.0011	0.017 ± 0.0012
Nagasaki, NAGASAKI	28	98.0	0.025 ± 0.0015	0.036 ± 0.0015
Naha, OKINAWA	27	109.5	0.008 ± 0.0010	0.012 ± 0.0010
February, 1979				
Sapporo, HOKKAIDO	29	72.5	0.023 ± 0.0014	0.033 ± 0.0014
Aomori, AOMORI	29	27.1	0.023 ± 0.0014	0.048 ± 0.0017
Sendai, MIYAGI	29	134.7	0.016 ± 0.0011	0.024 ± 0.0012
Yamagata, YAMAGATA	28	78.04	0.021 ± 0.0013	0.028 ± 0.0014
Ohkuma, FUKUSHIMA	29	166.6	0.019 ± 0.0014	0.032 ± 0.0015
Mito, IBARAGI	29	81.5	0.019 ± 0.0013	0.032 ± 0.0014
Shinjuku, TOKYO	28	26.9	0.022 ± 0.0014	0.038 ± 0.0016
Yokohama, KANAGAWA	28	90.4	0.022 ± 0.0014	0.037 ± 0.0015
Fukui, FUKUI	29	232.8	0.16 ± 0.004	0.22 ± 0.004
Shizuoka, SHIZUOKA	28	140.0	0.028 ± 0.0014	0.041 ± 0.0016

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Nagoya, AICHI	29	130.7	0.024 ± 0.0013	0.039 ± 0.0016
Kyoto, KYOTO	28	79.0	0.024 ± 0.0014	0.036 ± 0.0016
Kobe, HYOGO	28	81.0	0.022 ± 0.0014	0.038 ± 0.0016
Wakayama, WAKAYAMA	27	71.5	0.026 ± 0.0014	0.030 ± 0.0015
Tottori, TOTTORI	31	149.5	0.055 ± 0.0021	0.089 ± 0.0023
Matsue, SHIMANE	31	116.6	0.063 ± 0.0019	0.10 ± 0.002
Matsuyama, EHIME	29	88.5	0.022 ± 0.0013	0.032 ± 0.0014
Dazaifu, FUKUOKA	28	112.0	0.037 ± 0.0018	0.054 ± 0.0018
Saga, SAGA	27	99.3	0.035 ± 0.0016	0.054 ± 0.0018
Nagasaki, NAGASAKI	28	148.5	0.051 ± 0.0021	0.068 ± 0.0020
Naha, OKINAWA	29	119.5	0.032 ± 0.0017	0.049 ± 0.0017
March, 1979				
Sapporo, HOKKAIDO	33	110.5	0.041 ± 0.0018	0.061 ± 0.0018
Aomori, AOMORI	32	70.1	0.040 ± 0.0023	0.049 ± 0.0017
Sendai, MIYAGI	34	42.2	0.031 ± 0.0015	0.042 ± 0.0016
Yamagata, YAMAGATA	30	26.34	0.016 ± 0.0012	0.026 ± 0.0014
Ohkuma, FUKUSHIMA	33	49.0	0.026 ± 0.0015	0.041 ± 0.0016
Mito, IBARAGI	33	56.0	0.026 ± 0.0013	0.039 ± 0.0016
Shinjuku, TOKYO	31	89.2	0.038 ± 0.0015	0.059 ± 0.0020
Yokohama, KANAGAWA	31	91.7	0.037 ± 0.0017	0.056 ± 0.0018
Fukui, FUKUI	33	97.2	0.039 ± 0.0018	0.071 ± 0.0021
Shizuoka, SHIZUOKA	33	289.0	0.095 ± 0.0031	0.14 ± 0.003
Nagoya, AICHI	34	144.1	0.044 ± 0.0018	0.059 ± 0.0018
Kyoto, KYOTO	30	80.1	0.023 ± 0.0014	0.035 ± 0.0016
Kobe, HYOGO	31	86.4	0.023 ± 0.0013	0.030 ± 0.0014
Wakayama, WAKAYAMA	31	124.0	0.020 ± 0.0014	0.038 ± 0.0016
Tottori, TOTTORI	33	46.0	0.048 ± 0.0019	0.073 ± 0.0021
Matsue, SHIMANE	31	106.7	0.046 ± 0.0018	0.076 ± 0.0021
Hiroshima, HIROSHIMA	32	151.6	0.029 ± 0.0015	0.041 ± 0.0016
Matsuyama, EHIME	30	102.0	0.019 ± 0.0012	0.026 ± 0.0013
Dazaifu, FUKUOKA	31	158.9	0.037 ± 0.0016	0.055 ± 0.0019
Saga, SAGA	27	183.2	0.038 ± 0.0018	0.053 ± 0.0018
Kagoshima, KAGOSHIMA	33	291	0.009 ± 0.0009	0.011 ± 0.0010
Naha, OKINAWA	31	180.0	0.043 ± 0.0018	0.074 ± 0.0020
April, 1979				
Sapporo, HOKKAIDO	30	37.0	0.022 ± 0.0013	0.036 ± 0.0015
Aomori, AOMORI	30	67.7	0.049 ± 0.0019	0.077 ± 0.0021
Sendai, MIYAGI	30	127.5	0.036 ± 0.0018	0.052 ± 0.0018
Yamagata, YAMAGATA	31	58.5	0.043 ± 0.0017	0.066 ± 0.0019
Ohkuma, FUKUSHIMA	31	96.0	0.053 ± 0.0017	0.078 ± 0.0021

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Mito, IBARAGI	30	111.5	0.048 ± 0.0019	0.064 ± 0.0020
Shinjuku, TOKYO	30	114	0.040 ± 0.0018	0.073 ± 0.0020
Yokohama, KANAGAWA	33	96.3	0.040 ± 0.0018	0.066 ± 0.0019
Fukui, FUKUI	30	136.6	0.042 ± 0.0016	0.068 ± 0.0020
Shizuoka, SHIZUOKA	28	218.0	0.055 ± 0.0019	0.091 ± 0.0023
Nagoya, AICHI	29	217	0.063 ± 0.0022	0.087 ± 0.0022
Kyoto, KYOTO	31	163.6	0.051 ± 0.0019	0.083 ± 0.0025
Kobe, HYOGO	32	177.4	0.068 ± 0.0024	0.11 ± 0.002
Wakayama, WAKAYAMA	28	92.5	0.038 ± 0.0016	0.049 ± 0.0017
Tottori, TOTTORI	28	57.25	0.033 ± 0.0016	0.063 ± 0.0021
Matsue, SHIMANE	32	82.4	0.042 ± 0.0018	0.063 ± 0.0020
Hiroshima, HIROSHIMA	34	136.5	0.046 ± 0.0030	0.076 ± 0.0023
Matsuyama, EHIME	33	94.5	0.037 ± 0.0016	0.058 ± 0.0019
Dazaifu, FUKUOKA	31	102.3	0.030 ± 0.0014	0.050 ± 0.0020
Saga, SAGA	28	174.1	0.028 ± 0.0014	0.051 ± 0.0023
Nagasaki, NAGASAKI	29	183.0	0.082 ± 0.0025	0.11 ± 0.002
May, 1979				
Sapporo, HOKKAIDO	32	38.0	0.018 ± 0.0011	0.030 ± 0.0014
Aomori, AOMORI	32	132.0	0.042 ± 0.0016	0.065 ± 0.0024
Sendai, MIYAGI	32	30.8	0.026 ± 0.0015	0.044 ± 0.0017
Yamagata, YAMAGATA	31	101.60	0.049 ± 0.0018	0.065 ± 0.0019
Ohkuma, FUKUSHIMA	32	250.0	0.057 ± 0.0019	0.087 ± 0.0022
Mito, IBARAGI	32	223.0	0.066 ± 0.0020	0.10 ± 0.003
Shinjuku, TOKYO	31	210	0.051 ± 0.0022	0.087 ± 0.0027
Yokohama, KANAGAWA	35	140.3	0.038 ± 0.0016	0.068 ± 0.0021
Fukui, FUKUI	32	160.9	0.044 ± 0.0018	0.068 ± 0.0024
Shizuoka, SHIZUOKA	31	220.0	0.060 ± 0.0020	0.090 ± 0.0022
Nagoya, AICHI	32	166.8	0.022 ± 0.0013	0.036 ± 0.0020
Kyoto, KYOTO	30	153.2	0.057 ± 0.0021	0.083 ± 0.0028
Kobe, HYOGO	32	99.4	0.042 ± 0.0016	0.058 ± 0.0018
Wakayama, WAKAYAMA	28	190.5	0.080 ± 0.0024	0.12 ± 0.003
Tottori, TOTTORI	32	36.75	0.040 ± 0.0016	0.073 ± 0.0030
Matsue, SHIMANE	31	103.4	0.058 ± 0.0022	0.077 ± 0.0021
Hiroshima, HIROSHIMA	29	96.2	0.029 ± 0.0014	0.041 ± 0.0016
Matsuyama, EHIME	32	42.5	0.016 ± 0.0011	0.025 ± 0.0013
Dazaifu, FUKUOKA	30	92.2	0.011 ± 0.0009	0.022 ± 0.0015
Saga, SAGA	30	157.5	0.013 ± 0.0010	0.019 ± 0.0013
Nagasaki, NAGASAKI	31	84.5	0.013 ± 0.0010	0.022 ± 0.0014
Naha, OKINAWA	30	209.0	0.029 ± 0.0014	0.050 ± 0.0022

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
June, 1979				
Yamagata, YAMAGATA	31	145.70	0.040 ± 0.0017	0.053 ± 0.0018
Ohkuma, FUKUSHIMA	32	69.0	0.029 ± 0.0015	0.047 ± 0.0017
Shizuoka, SHIZUOKA	31	144.5	0.013 ± 0.0012	0.025 ± 0.0013
Nagoya, AICHI	32	267	0.035 ± 0.0016	0.057 ± 0.0018
Kyoto, KYOTO	32	292.9	0.023 ± 0.0015	0.035 ± 0.0015
Kobe, HYOGO	32	189.7	0.023 ± 0.0013	0.040 ± 0.0016
Kagoshima, KAGOSHIMA	31	385	0.006 ± 0.0008	0.016 ± 0.0011
Sapporo, HOKKAIDO	32	48.0	0.017 ± 0.0010	0.023 ± 0.0013
Aomori, AOMORI	32	197.2	0.023 ± 0.0013	0.036 ± 0.0015
Sendai, MIYAGI	32	73.5	0.031 ± 0.0014	0.046 ± 0.0017
Mito, IBARAGI	32	76	0.019 ± 0.0012	0.032 ± 0.0014
Shinjuku, TOKYO	30	58	0.010 ± 0.0010	0.014 ± 0.0011
Yokohama, KANAGAWA	29	73.8	0.011 ± 0.0010	0.014 ± 0.0010
Fukui, FUKUI	32	215.2	0.031 ± 0.0015	0.050 ± 0.0017
Wakayama, WAKAYAMA	33	534.0	0.023 ± 0.0015	0.047 ± 0.0017
Tottori, TOTTORI	31	144.3	0.037 ± 0.0020	0.057 ± 0.0021
Matsue, SHIMANE	31	165.1	0.020 ± 0.0013	0.035 ± 0.0015
Hiroshima, HIROSHIMA	35	317.9	0.028 ± 0.0014	0.041 ± 0.0017
Matsuyama, EHIME	32	444.5	0.013 ± 0.0012	0.028 ± 0.0014
Dazaifu, FUKUOKA	30	587.4	0.026 ± 0.0015	0.043 ± 0.0017
Saga, SAGA	31	607.1	0.033 ± 0.0015	0.045 ± 0.0017
Nagasaki, NAGASAKI	32	570.0	0.050 ± 0.0019	0.076 ± 0.0021
Naha, OKINAWA	33	232.0	0.011 ± 0.0011	0.024 ± 0.0013
July, 1979				
Sapporo, HOKKAIDO	31	85.5	0.023 ± 0.0013	0.038 ± 0.0015
Aomori, AOMORI	31	76.4	0.017 ± 0.0013	0.018 ± 0.0012
Sendai, MIYAGI	31	210.9	0.023 ± 0.0014	0.036 ± 0.0014
Yamagata, YAMAGATA	30	216.1	0.025 ± 0.0013	0.058 ± 0.0020
Ohkuma, FUKUSHIMA	32	199.5	0.036 ± 0.0015	0.060 ± 0.0019
Mito, IBARAGI	31	127.5	0.015 ± 0.0011	0.024 ± 0.0013
Shinjuku, TOKYO	32	113.4	0.025 ± 0.0014	0.034 ± 0.0015
Yokohama, KANAGAWA	30	128	0.022 ± 0.0013	0.038 ± 0.0016
Fukui, FUKUI	30	51.5	0.009 ± 0.0009	0.017 ± 0.0011
Shizuoka, SHIZUOKA	31	207	0.020 ± 0.0013	0.033 ± 0.0014
Nagoya, AICHI	21	105	0.017 ± 0.0010	0.026 ± 0.0013
Kyoto, KYOTO	30	34.9	0.006 ± 0.0009	0.008 ± 0.0010
Kobe, HYOGO	31	51.2	0.008 ± 0.0009	0.012 ± 0.0010
Wakayama, WAKAYAMA	34	96.5	0.007 ± 0.0010	0.012 ± 0.0010
Tottori, TOTTORI	33	33.0	0.018 ± 0.0014	0.014 ± 0.0011


Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Matsue, SHIMANE	32	121.9	0.014 ± 0.0010	0.021 ± 0.0012
Hiroshima, HIROSHIMA	31	124.94	0.013 ± 0.0011	0.011 ± 0.0010
Matsuyama, EHIME	30	72.0	0.007 ± 0.0010	0.014 ± 0.0011
Dazaifu, FUKUOKA	32	158.4	0.009 ± 0.0009	0.016 ± 0.0010
Saga, SAGA	32	312.4	0.016 ± 0.0012	0.034 ± 0.0015
Nagasaki, NAGASAKI	31	263.0	0.011 ± 0.0011	0.018 ± 0.0011
Naha, OKINAWA	31	105.0	0.0011±0.0007	0.005 ± 0.0008
August, 1979				
Sapporo, HOKKAIDO	32	41.5	0.006 ± 0.0009	0.007 ± 0.0008
Aomori, AOMORI	32	112.2	0.013 ± 0.0010	0.017 ± 0.0011
Sendai, MIYAGI	32	137.7	0.012 ± 0.0011	0.020 ± 0.0012
Yamagata, YAMAGATA	32	126.6	0.007 ± 0.0008	0.011 ± 0.0009
Ohkuma, FUKUSHIMA	32	79.0	0.011 ± 0.0010	0.016 ± 0.0013
Mito, IBARAGI	32	81	0.009 ± 0.0010	0.010 ± 0.0010
Shinjuku, TOKYO	32	83	0.009 ± 0.0009	0.016 ± 0.0011
Yokohama, KANAGAWA	32	100.0	0.006 ± 0.0007	0.010 ± 0.0009
Fukui, FUKUI	33	125.2	0.006 ± 0.0009	0.014 ± 0.0010
Shizuoka, SHIZUOKA	31	270.5	0.006 ± 0.0009	0.009 ± 0.0008
Nagoya, AICHI	34	123	0.011 ± 0.0009	0.016 ± 0.0011
Kyoto, KYOTO	33	139.6	0.014 ± 0.0010	0.015 ± 0.0011
Kobe, HYOGO	34	41.3	0.0033±0.0007	0.006 ± 0.0008
Wakayama, WAKAYAMA	29	112.5	0.006 ± 0.0009	0.009 ± 0.0010
Tottori, TOTTORI	31	53.0	0.015 ± 0.0013	0.010 ± 0.0009
Matsue, SHIMANE	32	85.6	0.010 ± 0.0009	0.015 ± 0.0011
Hiroshima, HIROSHIMA	32	55.3	0.014 ± 0.0010	0.007 ± 0.0008
Matsuyama, EHIME	32	126.5	0.004 ± 0.0006	0.004 ± 0.0007
Dazaifu, FUKUOKA	32	134	0.003 ± 0.0005	0.005 ± 0.0008
Saga, SAGA	32	245.0	0.004 ± 0.0009	0.005 ± 0.0008
Nagasaki, NAGASAKI	32	390.5	0.003 ± 0.0008	0.005 ± 0.0007
Naha, OKINAWA	31	388.5	0.004 ± 0.0007	0.010 ± 0.0009
September, 1979				
Sapporo, HOKKAIDO	31	138.5	0.011 ± 0.0009	0.014 ± 0.0010
Aomori, AOMORI	31	111.7	0.007 ± 0.0009	0.011 ± 0.0009
Sendai, MIYAGI	31	155.9	0.007 ± 0.0008	0.013 ± 0.0010
Yamagata, YAMAGATA	31	96.0	0.005 ± 0.0007	0.008 ± 0.0009
Ohkuma, FUKUSHIMA	31	111	0.009 ± 0.0008	0.012 ± 0.0010
Mito, IBARAGI	31	177	0.006 ± 0.0010	0.010 ± 0.0010
Shinjuku, TOKYO	31	140	0.010 ± 0.0010	0.012 ± 0.0010
Yokohama, KANAGAWA	33	91.1	0.007 ± 0.0008	0.011 ± 0.0010
Fukui, FUKUI	32	231.2	0.011 ± 0.0009	0.023 ± 0.0013
Shizuoka, SHIZUOKA	31	33.5	0.007 ± 0.0009	0.014 ± 0.0011

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Nagoya, AICHI	29	301	0.009 ± 0.0010	0.017 ± 0.0011
Kyoto, KYOTO	31	146.8	0.006 ± 0.0008	0.008 ± 0.0008
Kobe, HYOGO	29	108.3	0.005 ± 0.0007	0.006 ± 0.0008
Wakayama, WAKAYAMA	29	131	0.005 ± 0.0007	0.006 ± 0.0009
Tottori, TOTTORI	31	172.0	0.015 ± 0.0011	0.020 ± 0.0012
Matsue, SHIMANE	32	284.4	0.009 ± 0.0009	0.015 ± 0.0011
Hiroshima, HIROSHIMA	30	178.8	0.009 ± 0.0009	0.008 ± 0.0010
Matsuyama, EHIME	32	163.5	0.003 ± 0.0006	0.006 ± 0.0008
Dazaifu, FUKUOKA	32	68.5	0.006 ± 0.0008	0.006 ± 0.0008
Saga, SAGA	31	121.8	0.003 ± 0.0006	0.004 ± 0.0008
Nagasaki, NAGASAKI	31	47.0	0.003 ± 0.0007	0.004 ± 0.0007
Naha, OKINAWA	32	127.0	0.002 ± 0.0007	0.004 ± 0.0007
October, 1979				
Sapporo, HOKKAIDO	32	194.5	0.011 ± 0.0009	0.017 ± 0.0011
Aomori, AOMORI	32	202.0	0.013 ± 0.0011	0.023 ± 0.0014
Sendai, MIYAGI	32	145.1	0.012 ± 0.0010	0.019 ± 0.0012
Yamagata, YAMAGATA	32	77.5	0.005 ± 0.0008	0.009 ± 0.0010
Ohkuma, FUKUSHIMA	32	368.0	0.011 ± 0.0010	0.016 ± 0.0011
Mito, IBARAGI	32	261.5	0.009 ± 0.0009	0.011 ± 0.0010
Shinjuku, TOKYO	32	338.4	0.012 ± 0.0010	0.017 ± 0.0011
Yokohama, KANAGAWA	30	219.9	0.010 ± 0.0009	0.014 ± 0.0011
Fukui, FUKUI	31	114.4	0.010 ± 0.0009	0.018 ± 0.0012
Shizuoka, SHIZUOKA	32	270.0	0.006 ± 0.0008	0.011 ± 0.0010
Nagoya, AICHI	32	159	0.005 ± 0.0008	0.015 ± 0.0011
Kyoto, KYOTO	32	79.6	0.002 ± 0.0006	0.004 ± 0.0007
Kobe, HYOGO	31	78.8	0.004 ± 0.0008	0.008 ± 0.0009
Wakayama, WAKAYAMA	32	98.5	0.003 ± 0.0006	0.002 ± 0.0007
Tottori, TOTTORI	32	177.50	0.015 ± 0.0012	0.016 ± 0.0011
Matsue, SHIMANE	31	129.3	0.009 ± 0.0010	0.018 ± 0.0012
Hiroshima, HIROSHIMA	31	52.9	0.013 ± 0.0015	0.004 ± 0.0007
Matsuyama, EHIME	32	193.0	0.003 ± 0.0008	0.002 ± 0.0007
Dazaifu, FUKUOKA	31	102.7	0.004 ± 0.0008	0.004 ± 0.0008
Saga, SAGA	32	73.3	0.001 ± 0.0006	0.001 ± 0.0007
Nagasaki, NAGASAKI	32	91.0	0.004 ± 0.0007	0.002 ± 0.0007
Naha, OKINAWA	32	363.5	0.003 ± 0.0010	0.005 ± 0.0009
November, 1979				
Sapporo, HOKKAIDO	31	102.0	0.007 ± 0.0008	0.016 ± 0.0011
Aomori, AOMORI	33	183.6	0.013 ± 0.0011	0.026 ± 0.0013
Sendai, MIYAGI	33	132.7	0.006 ± 0.0008	0.008 ± 0.0009
Yamagata, YAMAGATA	30	108.5	0.004 ± 0.0007	0.006 ± 0.0009
Ohkuma, FUKUSHIMA	31	161.0	0.005 ± 0.0009	0.009 ± 0.0009

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Mito, IBARAGI	31	163.5	0.007 ± 0.0008	0.009 ± 0.0009
Shinjuku, TOKYO	31	182	0.007 ± 0.0008	0.013 ± 0.0011
Yokohama, KANAGAWA	31	155.4	0.008 ± 0.0009	0.013 ± 0.0010
Fukui, FUKUI	33	273.9	0.024 ± 0.0014	0.036 ± 0.0015
Shizuoka, SHIZUOKA	31	228	0.005 ± 0.0008	0.006 ± 0.0008
Nagoya, AICHI	31	118	0.004 ± 0.0007	0.007 ± 0.0008
Kyoto, KYOTO	33	109.1	0.007 ± 0.0014	0.009 ± 0.0013
Kobe, HYOGO	31	72.9	0.003 ± 0.0006	0.007 ± 0.0009
Wakayama, WAKAYAMA	31	84	0.004 ± 0.0008	0.008 ± 0.0010
Tottori, TOTTORI	33	201.0	0.023 ± 0.0014	0.031 ± 0.0015
Matsue, SHIMANE	31	115.9	0.014 ± 0.0012	0.021 ± 0.0012
Hiroshima, HIROSHIMA	33	109.4	0.005 ± 0.0008	0.005 ± 0.0008
Matsuyama, EHIME	30	132.5	0.004 ± 0.0007	0.006 ± 0.0009
Dazaufi, FUKUOKA	31	88.2	0.007 ± 0.0009	0.011 ± 0.0010
Saga, SAGA	34	65.7	0.004 ± 0.0007	0.003 ± 0.0007
Nagasaki, NAGASAKI	31	91.5	0.006 ± 0.0008	0.009 ± 0.0009
Naha, OKINAWA	34	177.5	0.004 ± 0.0008	0.006 ± 0.0009
December, 1979				
Sapporo, HOKKAIDO	27	41.0	0.006 ± 0.0009	0.012 ± 0.0010
Aomori, AOMORI	34	115.2	0.014 ± 0.0013	0.022 ± 0.0012
Sendai, MIYAGI	34	12.3	0.003 ± 0.0006	0.003 ± 0.0007
Yamagata, YAMAGATA	26	45.5	0.006 ± 0.0009	0.006 ± 0.0008
Ohkuma, FUKUSHIMA	35	9.6	0.002 ± 0.0007	0.003 ± 0.0007
Mito, IBARAGI	36	43	0.003 ± 0.0008	0.006 ± 0.0008
Shinjuku, TOKYO	32	51	0.003 ± 0.0006	0.006 ± 0.0008
Yokohama, KANAGAWA	36	40.2	0.002 ± 0.0007	0.005 ± 0.0008
Fukui, FUKUI	34	195.8	0.021 ± 0.0013	0.036 ± 0.0015
Shizuoka, SHIZUOKA	31	60	0.002 ± 0.0006	0.005 ± 0.0007
Nagoya, AICHI	38	25.8	0.004 ± 0.0010	0.007 ± 0.0009
Kyoto, KYOTO	36	70.3	0.005 ± 0.0007	0.005 ± 0.0008
Kobe, HYOGO	28	20.2	0.002 ± 0.0007	0.005 ± 0.0008
Wakayama, WAKAYAMA	40	72.1	0.003 ± 0.0008	0.008 ± 0.0009
Tottori, TOTTORI	34	116.25	0.016 ± 0.0013	0.021 ± 0.0012
Matsue, SHIMANE	30	15.5	0.030 ± 0.0016	0.020 ± 0.0013
Hiroshima, HIROSHIMA	33	75.6	0.009 ± 0.0009	0.005 ± 0.0008
Matsuyama, EHIME	29	49.0	0.003 ± 0.0007	0.002 ± 0.0007
Fukuoka, FUKUOKA	37	82.0	0.006 ± 0.0008	0.011 ± 0.0010
Saga, SAGA	36	81.9	0.005 ± 0.0009	0.008 ± 0.0009
Nagasaki, NAGASAKI	35	123	0.007 ± 0.0011	0.014 ± 0.0010
Naha, OKINAWA	35	88.0	0.003 ± 0.0008	0.006 ± 0.0008

Figure 1-1 Sampling Locations of Rain and dry fallout

- | | |
|---------------|---------------|
| 1. Sapporo | 19. Dazaifu |
| 2. Aomori | 20. Saga |
| 3. Sendai | 21. Nagasaki |
| 4. Yamagata | 22. Kagoshima |
| 5. Ohkuma | 23. Naha |
| 6. Mito | |
| 7. Shinjuku | |
| 8. Yokohama | |
| 9. Fukui | |
| 10. Shizuoka | |
| 11. Nagoya | |
| 12. Kyoto | |
| 13. Kobe | |
| 14. Wakayama | |
| 15. Tottori | |
| 16. Matsue | |
| 17. Hiroshima | |
| 18. Matsuyama | |



(1)-2 Strontium-90 and Cesium-137 in Rain and dry fallout (for WHO program)
(from Jan. 1979 to Dec. 1979)

— continued from No. 51 of this publication —

Table 1-2: Strontium-90 and Cesium-137 in Rain and dry fallout

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
January, 1979				
Akita, AKITA	27	95.44	0.028 ± 0.0016	0.053 ± 0.0018
Niigata, NIIGATA	28	115.80	0.039 ± 0.0016	0.065 ± 0.0020
Kanazawa, ISHIKAWA	29	194.5	0.090 ± 0.0025	0.14 ± 0.003
Nagano, NAGANO	29	16.0	0.007 ± 0.0010	0.010 ± 0.0009
Osaka, OSAKA	32	53	0.009 ± 0.0009	0.011 ± 0.0010
Okayama, OKAYAMA	27	54.7	0.007 ± 0.0009	0.012 ± 0.0010
Yamaguchi, YAMAGUCHI	30	57.5	0.017 ± 0.0012	0.025 ± 0.0013
Kochi, KOCHI	27	91.1	0.016 ± 0.0012	0.016 ± 0.0011
Kagoshima, KAGOSHIMA	29	121.4	0.014 ± 0.0012	0.019 ± 0.0011
February, 1979				
Akita, AKITA	29	90.71	0.035 ± 0.0019	0.049 ± 0.0017
Niigata, NIIGATA	29	111.64	0.053 ± 0.0018	0.076 ± 0.0021
Kanazawa, ISHIKAWA	29	253.5	0.12 ± 0.003	0.19 ± 0.003
Nagano, NAGANO	29	10.0	0.007 ± 0.0009	0.011 ± 0.0009
Osaka, OSAKA	28	70.78	0.014 ± 0.0013	0.027 ± 0.0015
Okayama, OKAYAMA	28	104.4	0.010 ± 0.0014	0.023 ± 0.0013
Yamaguchi, YAMAGUCHI	29	120.5	0.047 ± 0.0020	0.076 ± 0.0021
Kochi, KOCHI	27	158.4	0.037 ± 0.0016	0.062 ± 0.0019
Kagoshima, KAGOSHIMA	28	38.2	0.017 ± 0.0014	0.028 ± 0.0014
March, 1979				
Akita, AKITA	33	89.58	0.045 ± 0.0017	0.071 ± 0.0021
Niigata, NIIGATA	33	57.66	0.046 ± 0.0016	0.075 ± 0.0021
Kanazawa, ISHIKAWA	33	83.0	0.045 ± 0.0019	0.079 ± 0.0025
Nagano, NAGANO	33	41.5	0.011 ± 0.0010	0.014 ± 0.0010
Osaka, OSAKA	30	106	0.017 ± 0.0012	0.027 ± 0.0013
Okayama, OKAYAMA	32	96.0	0.018 ± 0.0012	0.025 ± 0.0013
Yamaguchi, YAMAGUCHI	31	200.5	0.044 ± 0.0019	0.058 ± 0.0018
Kochi, KOCHI	31	279.8	0.090 ± 0.0025	0.12 ± 0.003
Nagasaki, NAGASAKI	33	198.0	0.036 ± 0.0015	0.057 ± 0.0019
April, 1979				
Akita, AKITA	30	158.3	0.086 ± 0.0024	0.13 ± 0.003
Niigata, NIIGATA	31	91.97	0.042 ± 0.0018	0.066 ± 0.0020
Kanazawa, ISHIKAWA	30	105.5	0.052 ± 0.0019	0.076 ± 0.0021
Nagano, NAGANO	30	74.5	0.020 ± 0.0014	0.033 ± 0.0014
Osaka, OSAKA	31	178	0.051 ± 0.0016	0.075 ± 0.0020


Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Okayama, OKAYAMA	29	71.2	0.022 ± 0.0012	0.033 ± 0.0014
Yamaguchi, YAMAGUCHI	30	155.0	0.044 ± 0.0018	0.066 ± 0.0020
Kochi, KOCHI	30	283.1	0.087 ± 0.0023	0.13 ± 0.003
Kagoshima, KAGOSHIMA	30	115.5	0.015 ± 0.0013	0.016 ± 0.0011
May, 1979				
Akita, AKITA	32	84.08	0.036 ± 0.0015	0.052 ± 0.0017
Niigata, NIIGATA	31	56.68	0.025 ± 0.0013	0.039 ± 0.0019
Kanazawa, ISHIKAWA	35	131.5	0.072 ± 0.0023	0.11 ± 0.002
Nagano, NAGANO	32	76.5	0.038 ± 0.0017	0.049 ± 0.0019
Osaka, OSAKA	31	140	0.042 ± 0.0018	0.068 ± 0.020
Okayama, OKAYAMA	31	84.9	0.025 ± 0.0013	0.049 ± 0.0018
Yamaguchi, YAMAGUCHI	31	159.0	0.029 ± 0.0014	0.043 ± 0.0019
Kochi, KOCHI	31	191.8	0.030 ± 0.0014	0.034 ± 0.0015
Kagoshima, KAGOSHIMA	32	111	0.014 ± 0.0011	0.035 ± 0.0015
June, 1979				
Akita, AKITA	31	240.03	0.075 ± 0.0023	0.12 ± 0.003
Kanazawa, ISHIKAWA	29	111.5	0.028 ± 0.0015	0.045 ± 0.0016
Niigata, NIIGATA	32	77.4	0.034 ± 0.0018	0.061 ± 0.0020
Nagano, NAGANO	32	95.5	0.027 ± 0.0015	0.040 ± 0.0016
Osaka, OSAKA	32	160	0.023 ± 0.0013	0.033 ± 0.0014
Okayama, OKAYAMA	32	84.9	0.017 ± 0.0014	0.021 ± 0.0012
Yamaguchi, YAMAGUCHI	32	514.5	0.037 ± 0.0016	0.030 ± 0.0014
Kochi, KOCHI	30	380.2	0.038 ± 0.0015	0.048 ± 0.0017
Kagoshima, KAGOSHIMA	31	385	0.006 ± 0.0008	0.016 ± 0.0011
July, 1979				
Akita, AKITA	32	133.1	0.027 ± 0.0015	0.046 ± 0.0016
Niigata, NIIGATA	31	306.55	0.014 ± 0.0012	0.025 ± 0.0013
Kanazawa, ISHIKAWA	31	166	0.0077 ± 0.0012	0.012 ± 0.0012
Nagano, NAGANO	31	153.5	0.013 ± 0.0011	0.023 ± 0.0013
Osaka, OSAKA	31	58	0.007 ± 0.0009	0.008 ± 0.0009
Okayama, OKAYAMA	30	260.6	0.003 ± 0.0007	0.011 ± 0.0010
Yamaguchi, YAMAGUCHI	31	136.0	0.007 ± 0.0009	0.012 ± 0.0010
Kochi, KOCHI	32	153.1	0.004 ± 0.0008	0.005 ± 0.0008
Kagoshima, KAGOSHIMA	32	256.8	0.001 ± 0.0006	0.003 ± 0.0007
August, 1979				
Akita, AKITA	32	166.91	0.014 ± 0.0013	0.022 ± 0.0012
Niigata, NIIGATA	32	123.49	0.010 ± 0.0010	0.015 ± 0.0011
Kanazawa, ISHIKAWA	34	203.5	0.010 ± 0.0009	0.012 ± 0.0009
Nagano, NAGANO	32	162.0	0.003 ± 0.0007	0.008 ± 0.0009
Osaka, OSAKA	32	65	0.005 ± 0.0008	0.006 ± 0.0008

Location	Duration (Days)	Precipitation (mm)	⁹⁰ Sr (mCi/km ²)	¹³⁷ Cs (mCi/km ²)
Okayama, OKAYAMA	32	29.5	0.005 ± 0.0006	0.004 ± 0.0008
Yamaguchi, YAMAGUCHI	32	149.0	0.009 ± 0.0008	0.037 ± 0.0015
Matsuyama, EHIME	32	126.5	0.004 ± 0.0006	0.004 ± 0.0007
Kochi, KOCHI	32	338.8	0.007 ± 0.0008	0.012 ± 0.0010
Kagoshima, KAGOSHIMA	33	213.5	0.004 ± 0.0007	0.006 ± 0.0008
September, 1979				
Akita, AKITA	31	240.25	0.015 ± 0.0011	0.020 ± 0.0012
Niigata, NIIGATA	31	176.73	0.008 ± 0.0008	0.015 ± 0.0011
Kanazawa, ISHIKAWA	28	230.5	0.013 ± 0.0010	0.017 ± 0.0012
Nagano, NAGANO	31	63.5	0.003 ± 0.0007	0.003 ± 0.0007
Osaka, OSAKA	31	234	0.005 ± 0.0007	0.008 ± 0.0009
Okayama, OKAYAMA	31	69.5	0.003 ± 0.0006	0.004 ± 0.0007
Yamaguchi, YAMAGUCHI	31	161.0	0.010 ± 0.0010	0.014 ± 0.0010
Matsuyama, EHIME	32	163.5	0.003 ± 0.0006	0.006 ± 0.0008
Kochi, KOCHI	31	372.8	0.015 ± 0.0010	0.021 ± 0.0012
Kagoshima, KAGOSHIMA	31	193.5	0.004 ± 0.0008	0.007 ± 0.0009
October, 1979				
Akita, AKITA	32	182.75	0.011 ± 0.0010	0.023 ± 0.0012
Niigata, NIIGATA	32	83.8	0.011 ± 0.0011	0.017 ± 0.0011
Kanazawa, ISHIKAWA	31	154	0.013 ± 0.0011	0.025 ± 0.0013
Nagano, NAGANO	32	120.0	0.005 ± 0.0008	0.005 ± 0.0007
Osaka, OSAKA	31	124	0.002 ± 0.0006	0.004 ± 0.0007
Okayama, OKAYAMA	32	51.3	0.003 ± 0.0008	0.001 ± 0.0007
Yamaguchi, YAMAGUCHI	32	99.5	0.009 ± 0.0009	0.009 ± 0.0009
Kochi, KOCHI	32	202.4	0.007 ± 0.0008	0.005 ± 0.0008
Kagoshima, KAGOSHIMA	31	151	0.002 ± 0.0007	0.003 ± 0.0007
November, 1979				
Akita, AKITA	31	170.57	0.005 ± 0.0008	0.016 ± 0.0011
Niigata, NIIGATA	31	130.08	0.012 ± 0.0010	0.018 ± 0.0012
Kanazawa, ISHIKAWA	32	280	0.027 ± 0.0013	0.048 ± 0.0017
Nagano, NAGANO	31	84.5	0.002 ± 0.0007	0.003 ± 0.0008
Osaka, OSAKA	31	92	0.004 ± 0.0006	0.004 ± 0.0008
Okayama, OKAYAMA	31	193.6	0.003 ± 0.0006	0.004 ± 0.0008
Yamaguchi, YAMAGUCHI	31	83.0	0.010 ± 0.0010	0.013 ± 0.0010
Kochi, KOCHI	31	231.9	0.008 ± 0.0008	0.009 ± 0.0009
Kagoshima, KAGOSHIMA	32	107.5	0.002 ± 0.0006	0.007 ± 0.0008
December, 1979				
Akita, AKITA	31	172.1	0.006 ± 0.0007	0.027 ± 0.0013
Kanazawa, ISHIKAWA	36	188.5	0.031 ± 0.0017	0.048 ± 0.0017
Nagano, NAGANO	35	12.5	0.003 ± 0.0006	0.003 ± 0.0007
Osaka, OSAKA	31	34	0.003 ± 0.0008	0.004 ± 0.0007
Okayama, OKAYAMA	37	98.4	0.004 ± 0.0008	0.004 ± 0.0007

Location	Duration (Days)	Precipitation (mm)	^{90}Sr (mCi/km ²)	^{137}Cs (mCi/km ²)
Yamaguchi, YAMAGUCHI	36	91.0	0.008 ± 0.0009	0.015 ± 0.0011
Kochi, KOCHI	38	3.8	0.009 ± 0.0010	0.007 ± 0.0009
Kagoshima, KAGOSHIMA	33	73.5	0.014 ± 0.0009	0.008 ± 0.0008

Figure 1-2 Sampling Locations of Rain and dry fallout

1. Akita
2. Niigata
3. Kanazawa
4. Nagano
5. Osaka
6. Okayama
7. Yamaguchi
8. Matsuyama
9. Kochi
10. Nagasaki
11. Kagoshima



(2) Strontium-90 and Cesium-137 in Airborne dust
(from Nov. 1978 to Jul. 1979)

— continued from No. 51 of this publication —


Table 2: Strontium-90 and Cesium-137 in Airborne dust

Location	Sampling period	Absorption volume (m ³)	⁹⁰ Sr (10 ⁻³ pCi/m ³)	¹³⁷ Cs (10 ⁻³ pCi/m ³)
November ~ December 1978				
Ohkuma, FUKUSHIMA	11 ~ 12	12,735	0.3 ± 0.04	0.7 ± 0.03
Mito, IBARAGI	10 ~ 12	11,023	0.4 ± 0.06	0.4 ± 0.03
Niigata, NIIGATA	10 ~ 12	13,996.6	0.5 ± 0.04	0.8 ± 0.04
Fukui, FUKUI	10 ~ 12	22,100	0.6 ± 0.03	1.1 ± 0.03
Shizuoka, SHIZUOKA	10 ~ 11	10,488	0.4 ± 0.05	0.5 ± 0.03
Shizuoka, SHIZUOKA	12	10,350	0.3 ± 0.05	0.4 ± 0.03
Osaka, OSAKA	10 ~ 12	9,072	0.4 ± 0.06	0.6 ± 0.04
Tottori, TOTTORI	10 ~ 12	11,803	0.5 ± 0.06	0.7 ± 0.04
Hiroshima, HIROSHIMA	10 ~ 12	10,800	0.1 ± 0.04	0.2 ± 0.02
Nagasaki, NAGASAKI	11 ~ 12	13,259	0.5 ± 0.04	0.7 ± 0.03
January ~ March 1979				
Ohkuma, FUKUSHIMA	2 ~ 3	12,034	0.4 ± 0.04	0.6 ± 0.04
Mito, IBARAGI	1 ~ 3	10,379	0.2 ± 0.04	0.3 ± 0.03
Niigata, NIIGATA	1 ~ 3	12,699.4	0.6 ± 0.06	1.1 ± 0.04
Fukui, FUKUI	1 ~ 3	18,324	1.0 ± 0.05	1.7 ± 0.04
Shizuoka, SHIZUOKA	1 ~ 2	12,062	0.4 ± 0.05	0.6 ± 0.03
Shizuoka, SHIZUOKA	3	11,411	0.6 ± 0.04	0.9 ± 0.04
Nagoya, AICHI	1 ~ 3	10,740	0.7 ± 0.06	1.0 ± 0.05
Osaka, OSAKA	1 ~ 3	7,776	0.5 ± 0.05	0.7 ± 0.05
Tottori, TOTTORI	1 ~ 3	11,074.8	0.9 ± 0.05	1.3 ± 0.05
Hiroshima, HIROSHIMA	1 ~ 3	10,800	0.1 ± 0.03	0.1 ± 0.03
Nagasaki, NAGASAKI	1 ~ 3	8,161	0.9 ± 0.07	1.5 ± 0.06
April ~ June 1979				
Mito, IBARAGI	4 ~ 6	10,368	0.4 ± 0.04	0.5 ± 0.03
Niigata, NIIGATA	4 ~ 6	10,888.4	1.1 ± 0.06	1.6 ± 0.05
Fukui, FUKUI	4 ~ 6	19,249	1.2 ± 0.05	1.9 ± 0.04
Nagoya, AICHI	4 ~ 6	11,919	0.7 ± 0.05	1.0 ± 0.04
Osaka, OSAKA	4 ~ 6	7,776	0.5 ± 0.06	0.8 ± 0.05
Tottori, TOTTORI	4 ~ 6	12,577	0.9 ± 0.05	1.3 ± 0.05
Hiroshima, HIROSHIMA	4 ~ 6	10,800	0.2 ± 0.03	0.3 ± 0.03
Nagasaki, NAGASAKI	4 ~ 6	9,005	1.3 ± 0.07	1.9 ± 0.06
Ohkuma, FUKUSHIMA	4 ~ 6	12,415	0.4 ± 0.03	0.7 ± 0.04
Shizuoka, SHIZUOKA	4 ~ 6	17,252	0.6 ± 0.03	0.9 ± 0.03

Location	Sampling period	Absorption volume (m ³)	⁹⁰ Sr (10 ⁻³ pCi/m ³)	¹³⁷ Cs (10 ⁻³ pCi/m ³)
July ~ September 1979				
Ohkuma, FUKUSHIMA	7 ~ 8	12,380	0.2 ± 0.03	0.3 ± 0.02
Mito, IBARAGI	7 ~ 9	10,368	0.1 ± 0.03	0.2 ± 0.03
Niigata, NIIGATA	7 ~ 9	10,712.6	0.2 ± 0.03	0.3 ± 0.03
Fukui, FUKUI	7 ~ 9	29,521	0.3 ± 0.02	0.5 ± 0.2
Shizuoka, SHIZUOKA	7 ~ 9	16,906	0.2 ± 0.02	0.4 ± 0.02
Nagoya, AICHI	7 ~ 9	10,874	0.3 ± 0.04	0.4 ± 0.03
Osaka, OSAKA	7 ~ 9	9,072	0.2 ± 0.03	0.3 ± 0.03
Tottori, TOTTORI	7 ~ 9	13,375	0.3 ± 0.03	0.4 ± 0.03
Hiroshima, HIROSHIMA	7 ~ 9	10,800	0.1 ± 0.03	0.2 ± 0.03
Nagasaki, NAGASAKI	7 ~ 9	10,962	0.2 ± 0.03	0.5 ± 0.03

Figure 2 Sampling Locations of Airborne dust

1. Ohkuma, Fukushima
2. Mito, Ibaragi
3. Niigata
4. Fukui
5. Shizuoka
6. Nagoya
7. Osaka
8. Tottori
9. Hiroshima
10. Nagasaki



(3) Strontium-90 and Cesium-137 in Service water
(from Dec. 1978 to Dec. 1979)

— continued from No. 50 of this publication —

Table 3: Strontium-90 and Cesium-137 in Service water


Location	pH	⁹⁰ Sr (pCi/ℓ)	¹³⁷ Cs (pCi/ℓ)
(Source Water)			
December, 1978			
Nagoya, AICHI	6.8	0.09 ± 0.006	0.01 ± 0.004
Moriguchi, OSAKA	8.0	0.24 ± 0.009	0.02 ± 0.004
January, 1979			
Sapporo, HOKKAIDO	7.2	0.12 ± 0.007	0.01 ± 0.003
Kanamachi, TOKYO	7.5	0.07 ± 0.005	0.01 ± 0.004
Keage, KYOTO	7.80	0.33 ± 0.01	0.01 ± 0.003
Fukuoka, FUKUOKA	7.55	0.12 ± 0.006	0.01 ± 0.003
June, 1979			
Sapporo, HOKKAIDO	6.9	0.10 ± 0.008	0.01 ± 0.003
Yokohama, KANAGAWA	7.6	0.02 ± 0.003	0.001 ± 0.003
Nagoya, AICHI	7.1	0.10 ± 0.007	0.01 ± 0.003
Moriguchi, OSAKA	7.4	0.30 ± 0.010	0.01 ± 0.004
Kanamachi, TOKYO	6.8	0.18 ± 0.008	0.03 ± 0.004
Fukuoka, FUKUOKA	6.75	0.12 ± 0.007	0.003 ± 0.003
July, 1979			
Sapporo, HOKKAIDO	6.6	0.48 ± 0.013	0.01 ± 0.003
August, 1979			
Keage, KYOTO	7.25	0.35 ± 0.012	0.02 ± 0.004
December, 1979			
Tsukui-gun, KANAGAWA	7.8	0.02 ± 0.004	0.002 ± 0.003
(Tap Water)			
December, 1978			
Wakkanai, HOKKAIDO	6.7	0.49 ± 0.012	0.01 ± 0.003
Aomori, AOMORI	6.9	0.07 ± 0.005	0.01 ± 0.003
Sendai, MIYAGI	7.3	0.12 ± 0.007	0.001 ± 0.003
Fukushima, FUKUSHIMA	—	0.16 ± 0.007	0.001 ± 0.003
Yokohama, KANAGAWA	7.1	0.03 ± 0.004	0.01 ± 0.003
Niigata, NIIGATA	7.13	0.20 ± 0.009	0.01 ± 0.003
Kanazawa, ISHIKAWA	7.3	0.16 ± 0.007	0.02 ± 0.004
Nagano, NAGANO	7.5	0.04 ± 0.004	0.01 ± 0.003
Nagoya, AICHI	6.6	0.11 ± 0.007	0.01 ± 0.004
Osaka, OSAKA	6.6	0.21 ± 0.009	0.003 ± 0.003

Location	pH	⁹⁰ Sr (pCi/ℓ)	¹³⁷ Cs (pCi/ℓ)
Tottori, TOTTORI	7.4	0.09 ± 0.006	0.003 ± 0.003
Matsue, SHIMANE	6.9	0.19 ± 0.009	0.002 ± 0.003
Ube, YAMAGUCHI	7.0	0.12 ± 0.006	0.01 ± 0.003
Kochi, KOCHI	7.3	0.11 ± 0.006	0.01 ± 0.003
Fukuoka, FUKUOKA	6.9	0.12 ± 0.007	0.005 ± 0.003
Saga, SAGA	7.1	0.08 ± 0.006	0.000 ± 0.003
Nagasaki, NAGASAKI	7.0	0.11 ± 0.007	0.01 ± 0.004
January, 1979			
Kanamachi, TOKYO	7.0	0.07 ± 0.005	0.02 ± 0.004
Kyoto, KYOTO	7.68	0.33 ± 0.01	0.01 ± 0.003
Kagoshima, KAGOSHIMA	6.7	0.004 ± 0.003	0.000 ± 0.003
Naha, OKINAWA	7.3	0.18 ± 0.008	0.01 ± 0.003
February, 1979			
Shizuoka, SHIZUOKA	6.9	0.005 ± 0.003	0.004 ± 0.003
Wakayama, WAKAYAMA	6.75	0.08 ± 0.005	0.02 ± 0.004
June, 1979			
Sendai, MIYAGI	7.5	0.09 ± 0.006	0.003 ± 0.003
Akita, AKITA	6.5	0.29 ± 0.009	0.02 ± 0.004
Yamagata, YAMAGATA	6.7	0.12 ± 0.007	0.02 ± 0.004
Fukushima, FUKUSHIMA	—	0.25 ± 0.009	0.004 ± 0.003
Mito, IBARAGI Ⅱ	5.2	0.08 ± 0.006	0.01 ± 0.003
Yokohama, KANAGAWA	7.2	0.03 ± 0.004	0.003 ± 0.003
Kanazawa, ISHIKAWA	6.8	0.16 ± 0.007	0.01 ± 0.003
Nagano, NAGANO	7.5	0.05 ± 0.004	0.02 ± 0.004
Nagoya, AICHI	6.7	0.12 ± 0.007	0.01 ± 0.004
Osaka, OSAKA	6.4	0.19 ± 0.008	0.01 ± 0.003
Kobe, HYOGO	7.1	0.22 ± 0.008	0.02 ± 0.003
Ube, YAMAGUCHI	6.8	0.15 ± 0.007	0.00 ± 0.003
Kochi, KOCHI	7.3	0.09 ± 0.006	0.003 ± 0.003
Saga, SAGA	7.4	0.10 ± 0.006	0.005 ± 0.003
Aomori, AOMORI	7.2	0.09 ± 0.006	0.02 ± 0.003
Kanamachi, TOKYO	7.0	0.15 ± 0.007	0.01 ± 0.003
Niigata, NIIGATA	7.38	0.22 ± 0.009	0.02 ± 0.003
Fukui, FUKUI	7.1	0.001 ± 0.003	0.001 ± 0.003
Shizuoka, SHIZUOKA	7.1	0.00 ± 0.003	0.00 ± 0.003
Tottori, TOTTORI	7.7	0.11 ± 0.007	0.00 ± 0.003
Matsue, SHIMANE	6.5	0.24 ± 0.010	0.01 ± 0.003
Okayama, OKAYAMA	6.5	0.03 ± 0.004	0.00 ± 0.003
Hiroshima, HIROSHIMA	7.0	0.12 ± 0.007	0.004 ± 0.003
Fukuoka, FUKUOKA	6.73	0.15 ± 0.008	0.01 ± 0.003
Nagasaki, NAGASAKI	—	0.12 ± 0.007	0.01 ± 0.003

Location	pH	^{90}Sr (pCi/l)	^{137}Cs (pCi/l)
July, 1979			
Matsuyama, EHIME	7.1	0.09 ± 0.007	0.003 ± 0.003
Kagoshima, KAGOSHIMA	7.1	0.00 ± 0.003	0.002 ± 0.003
August, 1979			
Kyoto, KYOTO	7.10	0.31 ± 0.010	0.01 ± 0.004
Wakayama, WAKAYAMA	6.8	0.11 ± 0.006	0.01 ± 0.003
Naha, OKINAWA	7.4	0.16 ± 0.007	0.003 ± 0.003
December, 1979			
Yamagata, YAMAGATA	6.7	0.12 ± 0.007	0.01 ± 0.003
Mito, IBARAGI	6.5	0.06 ± 0.005	0.00 ± 0.003
Yokohama, KANAGAWA	6.7	0.03 ± 0.004	0.002 ± 0.003
Niigata, NIIGATA	7.42	0.15 ± 0.008	0.01 ± 0.003
Kanazawa, ISHIKAWA	7.2	0.15 ± 0.008	0.01 ± 0.003
Nagano, NAGANO	7.3	0.06 ± 0.005	0.004 ± 0.003
Kobe, HYOGO	6.9	0.22 ± 0.009	0.00 ± 0.003
Matsuyama, EHIME	7.1	0.07 ± 0.005	0.01 ± 0.003
May, 1979			
Mito, IBARAGI	8.9	0.32 ± 0.011	0.04 ± 0.004
July, 1979			
Barato-lake, HOKKAIDO	7.3	0.19 ± 0.008	0.04 ± 0.004
Akita, AKITA	6.5	0.16 ± 0.009	0.03 ± 0.004
October, 1979			
Fukushima, FUKUSHIMA	—	0.11 ± 0.006	0.01 ± 0.003
November, 1979			
Niigata, NIIGATA	7.18	0.23 ± 0.008	0.03 ± 0.004

Figure 3 Sampling Locations of Service water

- | | |
|----------------|---------------|
| 1. Wakkanai | 21. Matsue |
| 2. Sapporo | 22. Okayama |
| 3. Aomori | 23. Hiroshima |
| 4. Sendai | 24. Ube |
| 5. Akita | 25. Matsuyama |
| 6. Yamagata | 26. Kochi |
| 7. Fukushima | 27. Fukuoka |
| 8. Mito | 28. Saga |
| 9. Kanamachi | 29. Nagasaki |
| 10. Tsukui-gun | 30. Kagoshima |
| 11. Niigata | 31. Naha |
| 12. Kanazawa | |
| 13. Nagano | |
| 14. Shizuoka | |
| 15. Nagoya | |
| 16. Keage | |
| 17. Moriguchi | |
| 18. Wakayama | |
| 19. Kobe | |
| 20. Tottori | |



(4) Strontium-90 and Cesium-137 in Soil
(from Jul. 1978 to Sep. 1979)

— continued from No. 50 of this publication —


Table 4: Strontium-90 and Cesium-137 in Soil

Location	Sampling Depth (cm)	Air Soil (%)	Sr (%)	⁹⁰ Sr		¹³⁷ Cs	
				(pCi/Kg)	(mCi/Km ²)	(pCi/Kg)	(mCi/Km ²)
July, 1978							
Hagi, YAMAGUCHI	0 ~ 5	97.1	0.002	120 ± 7	3.3 ± 0.19	380 ± 11	11 ± 0.3
"	5 ~ 20	96.9	0.001	86 ± 6	13 ± 1.0	130 ± 7	20 ± 1.1
Naha, OKINAWA	0 ~ 5	94.7	0.023	93 ± 6	4.7 ± 0.30	350 ± 1	18 ± 0.6
"	5 ~ 20	94.8	0.024	88 ± 6	24 ± 1.6	180 ± 9	49 ± 2.3
August, 1978							
Tokai (Naka-gun), IBARAGI	0 ~ 5	91.5	0.004	710 ± 19	33 ± 0.9	3,100 ± 40	140 ± 2
"	5 ~ 20	92.8	0.005	560 ± 14	50 ± 1.0	280 ± 10	25 ± 0.9
May, 1979							
Atsumi-gun, AICHI	0 ~ 5	97.5	0.001	90 ± 5.6	2.4 ± 0.15	430 ± 11	11 ± 0.3
"	5 ~ 20	98.6	0.001	54 ± 4.8	7.5 ± 0.67	340 ± 10	48 ± 1.4
June, 1979							
Tsuyama, OKAYAMA	0 ~ 5	92.8	0.003	62 ± 5.3	1.4 ± 0.12	98 ± 6.4	2.2 ± 0.14
"	5 ~ 20	92.6	0.003	41 ± 4.6	2.5 ± 0.28	75 ± 5.8	45 ± 0.35
July, 1979							
Aomori, AOMORI	0 ~ 5	91.6	0.003	84 ± 5.9	2.7 ± 0.19	110 ± 7	3.4 ± 0.21
"	5 ~ 20	91.4	0.005	19 ± 3.7	2.4 ± 0.48	3.0 ± 3.4	0.39 ± 0.4
Yamagata, YAMAGATA	0 ~ 5	96.7	0.003	300 ± 10	24 ± 0.8	410 ± 11	33 ± 0.9
"	5 ~ 20	97.1	0.002	120 ± 6	14 ± 0.7	140 ± 7	16 ± 0.8
Katsusika, TOKYO	0 ~ 5	95.9	0.007	190 ± 8	22 ± 0.9	310 ± 10	36 ± 1.1
"	5 ~ 20	97.7	0.006	170 ± 7	42 ± 1.9	340 ± 10	87 ± 2.5
Kashiwazaki, NIIGATA	0 ~ 5	98.5	0.004	170 ± 8	15 ± 0.7	670 ± 14	59 ± 1.2
"	5 ~ 20	92.3	0.004	300 ± 10	40 ± 1.4	530 ± 13	73 ± 1.7
Kanazawa, ISHIKAWA	0 ~ 5	97.4	0.003	220 ± 9	10 ± 0.4	570 ± 13	27 ± 0.6
"	5 ~ 20	97.0	0.003	130 ± 7	17 ± 1.0	180 ± 8	24 ± 1.1
Nagano, NAGANO	0 ~ 5	97.4	0.007	180 ± 8	11 ± 0.5	450 ± 11	28 ± 0.7
"	5 ~ 20	97.5	0.009	110 ± 6	20 ± 1.2	150 ± 7	28 ± 1.3
Gotenba, SHIZUOKA	0 ~ 5	99.0	0.011	120 ± 6	2.8 ± 0.13	320 ± 10	7.2 ± 0.22
"	5 ~ 20	99.0	0.013	60 ± 4.6	4.6 ± 0.35	180 ± 8	14 ± 0.6
Miyazu, KYOTO	0 ~ 5	98.5	0.001	200 ± 8	5.4 ± 0.21	2,800 ± 30	75 ± 0.7
"	5 ~ 20	98.9	0.001	190 ± 8	35 ± 1.5	190 ± 8	35 ± 1.4
Iwakura, TOTTORI	0 ~ 5	97.6	0.002	130 ± 7	5.5 ± 0.29	740 ± 14	32 ± 0.6
"	5 ~ 20	96.9	0.003	160 ± 7	27 ± 1.2	310 ± 10	51 ± 1.6
Ota, SHIMANE	0 ~ 5	94.6	0.004	120 ± 8	4.9 ± 0.32	930 ± 17	39 ± 0.7
"	5 ~ 20	94.3	0.003	75 ± 5.9	13 ± 1.0	860 ± 16	150 ± 3

Location	Sampling Depth (cm)	Air Soil (%)	Sr (%)	⁹⁰ Sr		¹³⁷ Cs	
				(pCi/Kg)	(mCi/Km ²)	(pCi/Kg)	(mCi/Km ²)
Hiroshima, HIROSHIMA	0 ~ 5	98.8	0.003	16 ± 3.2	1.3 ± 0.25	24 ± 4.2	1.8 ± 0.32
"	5 ~ 20	99.0	0.002	21 ± 3.5	4.2 ± 0.70	59 ± 5.1	12 ± 1.0
Matsuyama, EHIME	0 ~ 5	97.7	0.001	81 ± 5.3	2.8 ± 0.18	210 ± 8	7.4 ± 0.29
"	5 ~ 20	97.6	0.001	86 ± 5.3	6.2 ± 0.38	290 ± 9	21 ± 0.7
Kochi, KOCHI	0 ~ 5	95.7	0.006	400 ± 12	21 ± 0.6	1,000 ± 20	54 ± 0.9
"	5 ~ 20	95.7	0.008	210 ± 9	25 ± 1.1	190 ± 8	23 ± 1.0
Fukuoka, FUKUOKA	0 ~ 5	99.0	0.002	500 ± 12	18 ± 0.4	340 ± 10	13 ± 0.4
"	5 ~ 20	99.0	0.002	120 ± 6	16 ± 0.8	44 ± 4.8	5.7 ± 0.62
Minamitakaki-gun, NAGASAKI	0 ~ 5	88.4	0.003	460 ± 15	10 ± 0.3	2,700 ± 30	60 ± 0.6
"	5 ~ 20	82.8	0.003	470 ± 13	46 ± 1.3	1,600 ± 20	160 ± 2
August, 1979							
Sapporo, HOKKAIDO	0 ~ 5	90.7	0.007	600 ± 13	21 ± 0.5	1,400 ± 20	51 ± 0.7
"	5 ~ 20	91.3	0.007	210 ± 8	41 ± 1.6	140 ± 7	29 ± 1.4
Shibata-gun, MIYAGI	0 ~ 5	92.1	0.003	240 ± 9	4.0 ± 0.15	910 ± 17	15 ± 0.3
"	5 ~ 20	91.0	0.003	80 ± 6.3	5.6 ± 0.44	170 ± 8	12 ± 0.6
Fukushima, FUKUSHIMA	0 ~ 5	90.0	0.003	180 ± 9	5.0 ± 0.24	330 ± 11	8.8 ± 0.29
"	5 ~ 20	88.6	0.003	38 ± 4.8	3.4 ± 0.44	18 ± 4.3	1.7 ± 0.39
Tokai (Naka-gun), IBARAGI	0 ~ 5	89.8	0.003	470 ± 13	21 ± 0.6	2,600 ± 30	110 ± 1
"	5 ~ 20	89.5	0.005	750 ± 18	66 ± 1.6	150 ± 8	13 ± 0.7
Fukui, FUKUI	0 ~ 5	97.4	0.004	180 ± 8	9.8 ± 0.42	850 ± 15	45 ± 0.8
"	5 ~ 20	97.6	0.004	160 ± 8	22 ± 1.1	680 ± 14	96 ± 2.0
Kobe, HYOGO	0 ~ 5	96.4	0.001	69 ± 5.2	2.2 ± 0.16	270 ± 9	8.4 ± 0.28
"	5 ~ 20	97.4	0.001	59 ± 5.1	3.9 ± 0.34	77 ± 5.5	5.1 ± 0.37
Wakayama, WAKAYAMA	0 ~ 5	99.3	0.002	18 ± 3.6	1.0 ± 0.19	30 ± 4.0	1.6 ± 0.21
"	5 ~ 20	99.3	0.002	19 ± 3.1	4.4 ± 0.74	30 ± 4.0	7.0 ± 0.93
Hagi, YAMAGUCHI	0 ~ 5	99.0	0.002	20 ± 3.3	0.9 ± 0.15	67 ± 5.3	3.0 ± 0.24
"	5 ~ 20	98.9	0.001	29 ± 3.7	6.1 ± 0.78	72 ± 5.3	15 ± 1.1
Saga, SAGA	0 ~ 5	99.0	0.002	66 ± 5.5	3.3 ± 0.28	16 ± 3.9	0.8 ± 0.20
"	5 ~ 20	99.1	0.002	16 ± 3.4	3.5 ± 0.73	4.9 ± 3.4	1.0 ± 0.74
Ibusuki-gun, KAGOSHIMA	0 ~ 5	96.5	0.011	160 ± 7	9.7 ± 0.41	840 ± 16	50 ± 0.9
"	5 ~ 20	95.7	0.011	300 ± 10	48 ± 1.6	580 ± 13	92 ± 2.1
September, 1979							
Yokohama, KANAGAWA	0 ~ 5	95.6	0.007	520 ± 13	13 ± 0.3	1,200 ± 20	30 ± 0.5
"	5 ~ 20	91.6	0.006	400 ± 12	41 ± 1.3	700 ± 15	71 ± 1.5

Figure 4 Sampling Locations of Soil

- | | |
|---------------------|----------------------|
| 1. Sapporo | 21. Hiroshima |
| 2. Aomori | 22. Hagi |
| 3. Shibata-gun | 23. Matsuyama |
| 4. Yamagata | 24. Kochi |
| 5. Fukushima | 25. Fukuoka |
| 6. Tokai (Naka-gun) | 26. Saga |
| 7. Katsushika | 27. Minamitakaki-gun |
| 8. Yokohama | 28. Ibusuki-gun |
| 9. Kashiwazaki | 29. Naha |
| 10. Kanazawa | |
| 11. Fukui | |
| 12. Nagano | |
| 13. Gotenba | |
| 14. Atsumi-gun | |
| 15. Miyazu | |
| 16. Wakayama | |
| 17. Kobe | |
| 18. Iwakura | |
| 19. Ota | |
| 20. Tsuyama | |



(5) Strontium-90 and Cesium-137 in Sea water
(from Jul. 1979 to Nov. 1979)


— continued from No. 50 of this publication —

Table 5: Strontium-90 and Cesium-137 in Sea water

Location	Cl (%)	Sample volume analyzed (ℓ)	⁹⁰ Sr (pCi/ℓ)	¹³⁷ Cs (pCi/ℓ)
July, 1979				
off-Niigata-port, NIIGATA	17.5	41.5	0.14 ± 0.014	—
Ise-bay, AICHI	15.5	40	0.15 ± 0.014	0.11 ± 0.013
off-Ajisu, YAMAGUCHI	18.0	39	0.13 ± 0.015	0.15 ± 0.013
Moji-port, FUKUOKA	17.1	38	0.13 ± 0.014	0.12 ± 0.013
off-Tomari-port, HOKKAIDO	18.6	35	0.14 ± 0.016	0.13 ± 0.014
August, 1979				
off-Tomari-port, HOKKAIDO	18.6	35	0.14 ± 0.016	0.13 ± 0.014
Mutsu-bay, AOMORI	17.2	43	0.14 ± 0.015	0.13 ± 0.012
off-Soma, FUKUSHIMA	15.1	40	0.11 ± 0.012	0.14 ± 0.012
Odawa-bay, KANAGAWA	17.1	40	0.12 ± 0.015	0.17 ± 0.014
Sendai, KAGOSHIMA	18	37	0.13 ± 0.015	0.16 ± 0.014
September, 1979				
Osaka-bay, OSAKA	13.1	40	0.16 ± 0.015	0.12 ± 0.012
November, 1979				
White beach, OKINAWA	19.3	40	0.10 ± 0.013	0.13 ± 0.013

Figure 5 Sampling Locations of Sea water

1. off Tomari-port
2. Mutsu-bay
3. off Soma
4. off Niigata-port
5. Odawa-bay
6. Ise-bay
7. Osaka-bay
8. off Ajisu
9. Moji-port
10. Sendai
11. White beach



(6) Strontium-90 and Cesium-137 in Sea sediments
(from Apr. 1979 to Nov. 1979)


— continued from No. 50 of this publication —

Table 6: Strontium-90 and Cesium-137 in Sea sediments

Location	Depth (m)	⁹⁰ Sr (pCi/kg)	¹³⁷ Cs (pCi/kg)
April, 1979			
off-Tokai (0.50 km), IBARAGI	5	0.2 ± 3.2	16 ± 3.6
off-Niigata-port, NIIGATA	30	7 ± 3.8	200 ± 8
Ise-bay, AICHI	18	0 ± 3.5	160 ± 7
Ajisu, YAMAGUCHI	5	10 ± 3.4	150 ± 7
in Moji-port, FUKUOKA	8.5	0.2 ± 3.0	90 ± 5.8
August, 1979			
Tomari, HOKKAIDO	5	0 ± 3.1	98 ± 6.1
Mutsu-bay, AOMORI	10	16 ± 4.3	250 ± 9
Soma, FUKUSHIMA	1	0 ± 2.6	8 ± 3.6
Odawa-bay, KANAGAWA	7	0 ± 3.2	130 ± 7
Sendai, KAGOSHIMA	4.5	0.6 ± 3.1	15 ± 3.8
September, 1979			
Osaka-port, OSAKA	10	6 ± 3.1	180 ± 8
November, 1979			
White-beach, OKINAWA	14	5 ± 3.2	49 ± 5.0

Figure 6 Sampling Locations of Sea sediments

- 1. Tomari
- 2. Mutsu-bay
- 3. Soma
- 4. off Niigata-port
- 5. off Tokai (0.50 km)
- 6. Odawa-bay
- 7. Ise-bay
- 8. Osaka-port
- 9. Ajisu
- 10. in Moji-port
- 11. Sendai
- 12. White-beach



(7) Strontium-90 and Cesium-137 in Total diet
(from Nov. 1978 to Dec. 1979)

— continued from No. 50 of this publication —

Table 7: Strontium-90 and Cesium-137 in Total diet


Location	Ash (g/p/d)	Ca (mg/p/d)	K (mg/p/d)	⁹⁰ Sr		¹³⁷ Cs	
				pCi/p/d	S.U.	pCi/p/d	C.U.
November, 1978							
Nishikanbara-gun, NIIGATA	20.1	582	2,620	5.8 ± 0.51	10 ± 0.9	6.1 ± 0.40	2.3 ± 0.15
Matsue, SHIMANE	23.3	882	3,080	7.7 ± 0.58	8.7 ± 0.66	10 ± 0.5	3.3 ± 0.18
Okayama, OKAYAMA	20.5	655	2,030	4.7 ± 0.48	7.2 ± 0.73	4.1 ± 0.35	2.0 ± 0.17
Saga, SAGA	15.9	739	1,730	3.6 ± 0.38	4.9 ± 0.51	2.8 ± 0.27	1.6 ± 0.15
December, 1978							
Fukushima, FUKUSHIMA	22.0	863	3,040	7.2 ± 0.63	8.3 ± 0.74	5.7 ± 0.43	1.9 ± 0.14
Mito, IBARAGI	23.2	880	2,540	6.5 ± 0.60	7.3 ± 0.68	5.3 ± 0.43	2.1 ± 0.17
Shinjuku, TOKYO	15.9	668	1,500	3.4 ± 0.38	5.0 ± 0.57	3.8 ± 0.29	2.5 ± 0.20
Hiratsuka, KANAGAWA	14.9	524	1,920	3.6 ± 0.35	6.9 ± 0.67	5.2 ± 0.33	2.7 ± 0.17
Kanazawa, ISHIKAWA	18.4	519	2,010	4.0 ± 0.45	7.8 ± 0.87	4.8 ± 0.36	2.4 ± 0.18
Nagano, NAGANO	20.0	1,050	2,170	4.5 ± 0.45	4.3 ± 0.43	4.8 ± 0.36	2.2 ± 0.17
Kyoto, KYOTO	19.5	1,540	2,850	7.3 ± 0.58	4.8 ± 0.37	6.5 ± 0.46	2.3 ± 0.16
Wakayama, WAKAYAMA	38.2	1,750	4,350	7.0 ± 0.84	4.0 ± 0.48	6.5 ± 0.62	1.5 ± 0.14
Yamaguchi, YAMAGUCHI	19.8	944	2,100	4.7 ± 0.44	5.0 ± 0.47	3.6 ± 0.34	1.7 ± 0.16
Matsuyama, EHIME	20.7	789	2,320	4.8 ± 0.45	6.1 ± 0.57	3.9 ± 0.35	1.7 ± 0.15
Nagasaki, NAGASAKI	15.5	671	1,860	4.4 ± 0.36	6.5 ± 0.54	3.4 ± 0.28	1.8 ± 0.15
Shibushi, KAGOSHIMA	19.0	774	2,220	4.3 ± 0.46	5.5 ± 0.59	7.9 ± 0.45	3.6 ± 0.20
January, 1979							
Hiroshima, HIROSHIMA	12.6	484	1,300	2.7 ± 0.30	5.5 ± 0.62	2.2 ± 0.22	1.7 ± 0.17
February, 1979							
Naha, OKINAWA	14.8	413	1,370	2.1 ± 0.29	5.1 ± 0.70	3.7 ± 0.29	2.7 ± 0.21
May, 1979							
Wakayama, WAKAYAMA	32.5	2,460	3,310	2.5 ± 0.45	1.0 ± 0.18	4.4 ± 0.39	1.3 ± 0.12
June, 1979							
Sapporo, HOKKAIDO	18.5	70.5	2,490	6.2 ± 0.57	8.8 ± 0.81	5.2 ± 0.37	2.1 ± 0.15
Aomori, AOMORI	18.5	624	1,850	11 ± 0.7	18 ± 1.1	8.3 ± 0.42	4.5 ± 0.23
Sendai, MIYAGI	15.3	518	1,810	2.7 ± 0.40	5.3 ± 0.77	4.6 ± 0.31	2.5 ± 0.17
Akita, AKITA	20.3	582	1,900	7.8 ± 0.63	13 ± 1.1	7.2 ± 0.42	3.8 ± 0.22
Fukushima, FUKUSHIMA	16.6	59.9	2,510	3.8 ± 0.43	6.3 ± 0.71	3.7 ± 0.34	1.5 ± 0.13

Location	Ash	Ca	K	⁹⁰ Sr		¹³⁷ Cs	
	(g/p/d)	(mg/p/d)	(mg/p/d)	pCi/p/d	S.U.	pCi/p/d	C.U.
Mito, IBARAGI	17.7	862	2,500	4.1 ± 0.47	4.8 ± 0.54	4.2 ± 0.33	1.7 ± 0.13
Shinjuku, TOKYO	16.1	559	1,650	3.6 ± 0.42	6.4 ± 0.74	3.5 ± 0.30	2.1 ± 0.18
Nishikanbara-gun, NIIGATA	22.2	818	2,800	5.7 ± 0.58	7.0 ± 0.71	4.2 ± 0.38	1.5 ± 0.14
Nishikanbara-gun, NIIGATA	21.6	493	2,070	3.0 ± 0.83	6.1 ± 1.7	4.1 ± 0.50	2.0 ± 0.24
Fukui, FUKUI	12.4	169	1,480	2.0 ± 0.30	12 ± 1.8	2.5 ± 0.22	1.7 ± 0.15
Nagano, NAGANO	15.4	654	1,980	2.8 ± 0.38	4.4 ± 0.58	3.0 ± 0.27	1.5 ± 0.13
Nagoya, AICHI	18.2	681	2,340	4.1 ± 0.51	6.0 ± 0.76	5.6 ± 0.37	2.4 ± 0.16
Kyoto, KYOTO	20.4	890	3,070	3.4 ± 0.42	3.8 ± 0.47	3.4 ± 0.37	1.1 ± 0.12
Neyagawa, OSAKA	16.7	453	2,260	2.9 ± 0.37	6.4 ± 0.82	3.1 ± 0.28	1.4 ± 0.12
Matsue, SHIMANE	19.9	888	2,480	4.6 ± 0.57	5.2 ± 0.65	5.7 ± 0.39	2.3 ± 0.16
Okayama, OKAYAMA	14.5	578	1,820	5.7 ± 0.43	9.8 ± 0.74	3.6 ± 0.26	2.0 ± 0.15
Yamaguchi, YAMAGUCHI	16.8	499	1,880	3.3 ± 0.51	6.7 ± 1.02	2.4 ± 0.36	1.3 ± 0.19
Matsuyama, EHIME	14.7	679	2,340	3.1 ± 0.37	4.5 ± 0.54	3.5 ± 0.27	1.5 ± 0.12
Kochi, KOCHI	15.4	592	2,200	4.4 ± 0.50	7.4 ± 0.85	3.9 ± 0.40	1.8 ± 0.18
Kasuga, FUKUOKA	18.5	787	2,450	2.4 ± 0.45	3.0 ± 0.57	4.5 ± 0.35	1.8 ± 0.14
Saga, SAGA	13.1	592	1,540	2.6 ± 0.35	4.4 ± 0.59	2.7 ± 0.23	1.8 ± 0.15
Nagasaki, NAGASAKI	11.4	528	1,380	2.2 ± 0.25	4.2 ± 0.48	3.2 ± 0.23	2.3 ± 0.17
Shibushi, KAGOSHIMA	16.4	596	1,930	4.2 ± 0.42	7.0 ± 0.70	4.0 ± 0.31	2.1 ± 0.16
Kakogawa, HYOGO	8.63	485	1,190	2.2 ± 0.28	4.6 ± 0.58	3.6 ± 0.23	3.0 ± 0.19
July, 1979							
Yamagata, YAMAGATA	14.9	384	1,940	2.9 ± 0.37	7.6 ± 0.97	2.9 ± 0.26	1.5 ± 0.13
Hiratsuka, KANAGAWA	15.6	505	2,290	2.7 ± 0.39	5.4 ± 0.76	3.9 ± 0.30	1.7 ± 0.13
Kanazawa, ISHIKAWA	14.0	400	1,850	2.8 ± 0.30	7.0 ± 0.76	2.7 ± 0.25	1.5 ± 0.14
Iwami-gun, TOTTORI	15.4	381	1,790	4.4 ± 0.61	12 ± 1.6	2.8 ± 0.36	1.6 ± 0.20
Matsue, SHIMANE	19.9	888	2,480	4.6 ± 0.57	5.2 ± 0.65	5.7 ± 0.39	2.3 ± 0.16
September, 1979							
Hiroshima, HIROSHIMA	16.7	498	1,720	2.1 ± 0.34	4.1 ± 0.67	2.7 ± 0.27	1.6 ± 0.16
October, 1979							
Mito, IBARAGI	20.9	704	2,610	5.0 ± 0.54	7.1 ± 0.76	8.8 ± 0.51	3.4 ± 0.20
November, 1979							
Sendai, MIYAGI	20.4	599	2,090	2.7 ± 0.43	4.5 ± 0.73	5.6 ± 0.40	2.7 ± 0.19
Hiratsuka, KANAGAWA	16.0	645	2,250	4.0 ± 0.40	6.2 ± 0.62	4.5 ± 0.33	2.0 ± 0.14
Shizuoka, SHIZUOKA	20.8	1,010	2,490	5.3 ± 0.59	5.3 ± 0.59	5.5 ± 0.38	2.2 ± 0.15
Nagoya, AICHI	16.5	889	2,260	3.2 ± 0.39	3.6 ± 0.44	4.9 ± 0.33	2.2 ± 0.15
Wakayama, WAKAYAMA	26.3	2,370	3,090	5.8 ± 0.58	2.4 ± 0.25	4.7 ± 0.43	1.5 ± 0.14
Iwami-gun, TOTTORI	16.0	440	2,060	4.3 ± 0.44	9.7 ± 1.01	5.4 ± 0.33	2.6 ± 0.16
Okayama, OKAYAMA	17.8	597	2,170	3.8 ± 0.44	6.4 ± 0.74	3.1 ± 0.31	1.4 ± 0.14
Kochi, KOCHI	15.1	705	2,120	3.9 ± 0.38	5.5 ± 0.54	2.4 ± 0.27	1.1 ± 0.13
Fukuoka, FUKUOKA	15.9	531	2,000	1.8 ± 0.31	3.7 ± 0.62	5.8 ± 0.34	2.9 ± 0.17
Saga, SAGA	18.4	600	1,930	2.9 ± 0.38	4.8 ± 0.63	1.6 ± 0.25	0.8 ± 0.13

Location	Ash	Ca	K	⁹⁰ Sr		¹³⁷ Cs	
	(g/p/d)	(mg/p/d)	(mg/p/d)	pCi/p/d	S.U.	pCi/p/d	C.U.
Shibushi, KAGOSHIMA	18.1	813	2,050	4.8 ± 0.45	6.0 ± 0.55	3.9 ± 0.33	1.9 ± 0.16
December, 1979							
Yamagata, YAMAGATA	16.4	453	1,580	2.3 ± 0.32	5.1 ± 0.71	4.7 ± 0.31	3.0 ± 0.20
Nagano, NAGANO	21.6	1,130	2,690	4.0 ± 0.57	3.5 ± 0.50	4.4 ± 0.37	1.6 ± 0.14

Figure 7 Sampling Locations of Total diet

- | | |
|----------------------|---------------|
| 1. Sapporo | 21. Matsue |
| 2. Aomori | 22. Okayama |
| 3. Sendai | 23. Hiroshima |
| 4. Akita | 24. Yamaguchi |
| 5. Yamagata | 25. Matsuyama |
| 6. Fukushima | 26. Kochi |
| 7. Mito | 27. Fukuoka |
| 8. Shinjuku | 28. Saga |
| 9. Hiratsuka | 29. Nagasaki |
| 10. Nishikanbara-gun | 30. Shibushi |
| 11. Kanazawa | 31. Naha |
| 12. Fukui | |
| 13. Nagano | |
| 14. Shizuoka | |
| 15. Nagoya | |
| 16. Neyagawa | |
| 17. Kyoto | |
| 18. Kakogawa | |
| 19. Wakayama | |
| 20. Iwami-gun | |



(8) Strontium-90 and Cesium-137 in Rice
(from Dec. 1978 to Dec. 1979)

— continued from No. 51 of this publication —


Table 8: Strontium-90 and Cesium-137 in Rice

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
December, 1978							
Fukushima, FUKUSHIMA	0.441	0.0047	0.093	0.4 ± 0.16	8.2 ± 3.3	3.3 ± 0.21	3.6 ± 0.23
January, 1979							
Hirosaki, AOMORI	0.444	0.0051	0.097	0.8 ± 0.16	15 ± 3.2	4.3 ± 0.22	4.4 ± 0.23
Nagoya, AICHI	0.427	0.0051	0.086	0.7 ± 0.15	13 ± 2.9	12 ± 0.3	15 ± 0.4
Kobe, HYOGO	0.410	0.0052	0.087	0.9 ± 0.14	17 ± 2.7	4.6 ± 0.21	5.3 ± 0.25
Akashi, HYOGO	0.426	0.0054	0.090	0.3 ± 0.13	5.0 ± 2.3	1.2 ± 0.14	1.4 ± 0.15
Tottori, TOTTORI	0.390	0.0049	0.079	1.0 ± 0.15	20 ± 3.1	2.4 ± 0.16	3.1 ± 0.20
Matsue, SHIMANE	0.428	0.0055	0.088	1.2 ± 0.18	22 ± 3.3	11 ± 0.3	13 ± 0.4
Okayama, OKAYAMA	0.503	0.0051	0.12	1.2 ± 0.19	24 ± 3.7	4.1 ± 0.22	3.5 ± 0.19
Saga, SAGA	0.628	0.0064	0.14	0.9 ± 0.22	13 ± 3.4	2.5 ± 0.22	1.8 ± 0.16
Nagasaki, NAGASAKI	0.423	0.0054	0.089	0.8 ± 0.17	15 ± 3.1	3.2 ± 0.18	3.6 ± 0.21
Kagoshima, KAGOSHIMA	0.496	0.0066	0.098	0.9 ± 0.19	13 ± 2.8	2.4 ± 0.21	2.5 ± 0.21
September, 1979							
Kanazawa, ISHIKAWA	0.546	0.0106	0.22	1.4 ± 0.67	14 ± 6.3	1.9 ± 0.56	0.9 ± 0.26
Minami-Azumi-gun, NAGANO	0.467	0.0070	0.11	0.6 ± 0.25	7.9 ± 3.5	0.7 ± 0.23	0.6 ± 0.21
Yamagata, YAMAGATA	0.421	0.0056	0.12	0.9 ± 0.28	17 ± 5.0	4.0 ± 0.31	3.5 ± 0.27
Mito, IBARAGI	0.414	0.0047	0.093	0.7 ± 0.31	14 ± 6.6	2.6 ± 0.29	2.8 ± 0.31
Mito, IBARAGI	0.372	0.0051	0.090	0.2 ± 0.27	4.2 ± 5.2	2.6 ± 0.27	2.9 ± 0.30
Shinjuku, TOKYO	0.424	0.0051	0.11	0.6 ± 0.22	12 ± 4.3	3.6 ± 0.30	3.4 ± 0.28
Niigata, NIIGATA	0.350	0.0050	0.088	1.0 ± 0.25	21 ± 4.9	13 ± 0.20	1.5 ± 0.22
Nishikanbara-gun, NIIGATA	0.328	0.0044	0.10	0.7 ± 0.21	15 ± 4.9	1.7 ± 0.20	1.7 ± 0.20
Wakayama, WAKAYAMA	0.372	0.0051	0.099	1.0 ± 0.24	19 ± 4.8	1.2 ± 0.21	1.2 ± 0.21
November, 1979							
Akita, AKITA	0.448	0.0050	0.099	1.6 ± 0.29	32 ± 5.9	7.8 ± 0.41	7.9 ± 0.42
Yokohama, KANAGAWA	0.450	0.0057	0.095	0.4 ± 0.25	7.1 ± 4.5	1.4 ± 0.25	1.4 ± 0.27
Fukui, FUKUI	0.402	0.0054	0.097	0.5 ± 0.22	8.3 ± 4.1	1.5 ± 0.23	1.6 ± 0.24
Shizuoka, SHIZUOKA	0.433	0.0062	0.11	0.9 ± 0.26	14 ± 4.2	7.3 ± 0.39	6.8 ± 0.37
Kyoto, KYOTO	0.453	0.0075	0.11	0.4 ± 0.26	5.9 ± 3.4	1.2 ± 0.25	1.1 ± 0.23
Osaka, OSAKA	0.404	0.0057	0.12	0.3 ± 0.26	5.9 ± 4.6	2.1 ± 0.25	1.8 ± 0.20

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
December, 1979							
Kurihara-gun, MIYAGI	0.390	0.0050	0.096	0.4 ± 0.26	7.4 ± 5.3	1.1 ± 0.22	1.1 ± 0.24
Kobe, HYOGO	0.394	0.0055	0.088	1.1 ± 0.26	21 ± 4.8	4.6 ± 0.32	5.2 ± 0.36
Akashi, HYOGO	0.510	0.0071	0.11	0.2 ± 0.28	2.4 ± 4.0	0.8 ± 0.27	0.7 ± 0.24
Tottori, TOTTORI	0.337	0.0053	0.079	0.5 ± 0.21	8.6 ± 4.0	5.1 ± 0.29	6.5 ± 0.36
Okayama, OKAYAMA	0.607	0.0069	0.17	0.0 ± 0.33	0.0 ± 4.8	2.4 ± 0.35	1.4 ± 0.21
Hiroshima, HIROSHIMA	0.349	0.0049	0.082	0.6 ± 0.23	13 ± 4.6	2.3 ± 0.24	2.8 ± 0.29
Kochi, KOCHI	0.420	0.0053	0.10	0.9 ± 0.25	17 ± 4.7	2.7 ± 0.27	2.6 ± 0.27
Saga, SAGA	0.644	0.0072	0.16	0.0 ± 0.32	0.0 ± 45	2.8 ± 0.37	1.7 ± 0.23
Kagoshima, KAGOSHIMA	0.481	0.0045	0.099	0.9 ± 0.37	21 ± 8.2	4.3 ± 0.41	4.4 ± 0.41

Figure 8 Sampling Locations of Rice

- | | |
|----------------------|---------------|
| 1. Hirosaki | 21. Tottori |
| 2. Akita | 22. Matsue |
| 3. Kurihara-gun | 23. Okayama |
| 4. Yamagata | 24. Hiroshima |
| 5. Fukushima | 25. Kochi |
| 6. Mito | 26. Saga |
| 7. Shinjuku | 27. Nagasaki |
| 8. Yokohama | 28. Kagoshima |
| 9. Niigata | |
| 10. Nishikanbara-gun | |
| 11. Kanazawa | |
| 12. Fukui | |
| 13. Minami-Azumi-gun | |
| 14. Shizuoka | |
| 15. Nagoya | |
| 16. Osaka | |
| 17. Kyoto | |
| 18. Kobe | |
| 19. Akashi | |
| 20. Wakayama | |



(9)-1 Strontium-90 and Cesium-137 in Milk (producing districts for WHO program)
(from Nov. 1978 to Nov. 1979)

— continued from No. 51 of this publication —


Table 9-1: Strontium-90 and Cesium-137 in Milk

Location	Ash (g/p/d)	Ca (mg/p/d)	K (mg/p/d)	⁹⁰ Sr		¹³⁷ Cs	
				pCi/p/d	S.U.	pCi/p/d	C.U.
November, 1978							
Nishikanbara-gun, NIIGATA	7.34	1.25	1.53	2.6 ± 0.33	2.1 ± 0.26	4.2 ± 0.31	2.7 ± 0.20
Katsuyama, FUKUI	8.88	1.28	1.99	3.1 ± 0.39	2.4 ± 0.30	14 ± 0.5	7.0 ± 0.26
Kasuya-gun, FUKUOKA	7.75	1.26	1.68	1.9 ± 0.29	1.5 ± 0.23	2.3 ± 0.27	1.3 ± 0.16
Shobara, HIROSHIMA	6.21	0.983	1.40	1.7 ± 0.23	1.7 ± 0.24	3.2 ± 0.24	2.3 ± 0.17
December, 1978							
Sapporo, HOKKAIDO	7.35	1.19	1.57	3.4 ± 0.35	2.9 ± 0.30	11 ± 0.4	7.0 ± 0.27
Yatsuka-gun, SHIMANE	7.63	1.05	1.58	0.93 ± 0.29	0.89 ± 0.28	5.2 ± 0.32	3.3 ± 0.20
January, 1979							
Toyono-gun, OSAKA	7.33	1.09	1.51	1.1 ± 0.28	0.98 ± 0.25	1.9 ± 0.23	1.2 ± 0.15
February, 1979							
Sapporo, HOKKAIDO	7.32	1.17	1.65	3.0 ± 0.35	2.5 ± 0.30	7.4 ± 0.35	4.5 ± 0.21
Hachijyo-Island, TOKYO	6.99	1.08	1.56	7.9 ± 0.44	7.3 ± 0.41	30 ± 0.7	19 ± 0.4
Nishikanbara-gun, NIIGATA	7.65	1.23	1.71	2.8 ± 0.30	2.3 ± 0.25	4.5 ± 0.32	2.6 ± 0.19
Katsuyama, FUKUI	7.14	1.09	1.63	4.6 ± 0.39	4.2 ± 0.36	12 ± 0.4	7.6 ± 0.27
Shobara, HIROSHIMA	6.77	1.03	1.53	2.3 ± 0.28	2.2 ± 0.27	3.9 ± 0.27	2.6 ± 0.18
Kochi, KOCHI	7.63	1.26	1.48	2.9 ± 0.37	2.3 ± 0.29	2.4 ± 0.24	1.6 ± 0.16
Kasuya-gun, FUKUOKA	7.71	1.25	1.64	1.3 ± 0.25	1.0 ± 0.20	2.4 ± 0.24	1.5 ± 0.15
March, 1979							
Aira-gun, KAGOSHIMA	7.37	1.18	1.61	1.6 ± 0.28	1.4 ± 0.24	5.7 ± 0.32	3.5 ± 0.20
Yatsuka-gun, SHIMANE	7.19	1.10	1.59	2.6 ± 0.37	2.4 ± 0.34	5.4 ± 0.31	3.4 ± 0.19
Hachijyo-Island, TOKYO	6.91	1.05	1.64	5.9 ± 0.40	5.6 ± 0.38	36 ± 0.7	22 ± 0.4
Toyono-gun, OSAKA	7.10	1.05	1.49	1.6 ± 0.27	1.5 ± 0.25	2.1 ± 0.22	1.4 ± 0.15
Kochi, KOCHI	7.24	1.18	1.41	6.8 ± 0.44	5.7 ± 0.37	3.7 ± 0.30	2.6 ± 0.21
Kasuya-gun, FUKUOKA	7.50	1.16	1.61	2.0 ± 0.29	1.7 ± 0.25	4.3 ± 0.29	2.7 ± 0.18
June, 1979							
Nishikanbara-gun, NIIGATA	7.38	1.17	1.64	3.7 ± 0.35	3.1 ± 0.30	8.2 ± 0.37	5.0 ± 0.23
Katsuyama, FUKUI	6.78	0.985	1.69	3.8 ± 0.34	3.8 ± 0.34	6.1 ± 0.31	3.6 ± 0.19
Aira-gun, KAGOSHIMA	7.01	1.06	1.57	3.2 ± 0.32	3.0 ± 0.30	3.6 ± 0.26	2.3 ± 0.17
Sapporo, HOKKAIDO	7.58	1.24	1.70	3.1 ± 0.31	2.5 ± 0.25	9.4 ± 0.40	5.5 ± 0.23
Yatsuka-gun, SHIMANE	7.06	1.10	1.58	4.1 ± 0.33	3.7 ± 0.30	5.4 ± 0.31	3.4 ± 0.20

Location	Ash	Ca	K	⁹⁰ Sr		¹³⁷ Cs	
	(g/p/d)	(mg/p/d)	(mg/p/d)	pCi/p/d	S.U.	pCi/p/d	C.U.
July, 1979							
Shobara, HIROSHIMA	6.12	0.993	1.43	1.2 ± 0.25	1.2 ± 0.25	1.5 ± 0.22	1.0 ± 0.16
August, 1979							
Nishikanbara-gun, NIIGATA	6.98	1.03	1.49	1.8 ± 0.23	1.8 ± 0.23	2.9 ± 0.24	1.9 ± 0.16
Katsuyama, FUKUI	6.80	1.04	1.72	2.9 ± 0.31	2.8 ± 0.30	5.2 ± 0.29	3.0 ± 0.17
Toyono-gun, OSAKA	6.89	1.04	1.48	1.4 ± 0.25	1.3 ± 0.25	2.9 ± 0.25	2.0 ± 0.17
Yatsuka-gun, SHIMANE	7.27	1.09	1.56	2.4 ± 0.27	2.2 ± 0.25	7.1 ± 0.35	4.5 ± 0.23
Kasuya-gun, FUKUOKA	7.41	1.09	1.72	1.4 ± 0.25	1.3 ± 0.23	1.8 ± 0.23	1.1 ± 0.13
Kochi, KOCHI	7.30	1.03	1.55	4.3 ± 0.37	4.1 ± 0.35	1.7 ± 0.22	1.1 ± 0.14
September, 1979							
Sapporo, HOKKAIDO	7.69	1.23	1.72	2.7 ± 0.29	2.2 ± 0.23	5.4 ± 0.33	3.2 ± 0.19
Shobara, HIROSHIMA	6.56	1.00	1.48	0.9 ± 0.19	0.9 ± 0.18	2.4 ± 0.23	1.6 ± 0.15
Aira-gun, KAGOSHIMA	6.98	1.07	1.58	2.6 ± 0.28	2.5 ± 0.26	3.8 ± 0.26	2.4 ± 0.17
November, 1979							
Hachijyo-Island, TOKYO	6.84	1.08	1.50	9.6 ± 0.50	8.9 ± 0.47	58 ± 0.9	39 ± 0.6
Nishikanbara-gun, NIIGATA	7.82	1.17	1.51	1.8 ± 0.26	1.5 ± 0.22	2.6 ± 0.26	1.7 ± 0.17
Toyono-gun, OSAKA	7.02	1.05	1.49	1.7 ± 0.24	1.7 ± 0.23	2.0 ± 0.22	1.3 ± 0.15
Kochi, KOCHI	7.44	1.10	1.41	2.6 ± 0.29	2.3 ± 0.26	1.7 ± 0.22	1.2 ± 0.16
Aira-gun, KAGOSHIMA	6.73	1.05	1.55	1.8 ± 0.27	1.7 ± 0.26	4.0 ± 0.28	2.6 ± 0.18

Figure 9-1 Sampling location of Milk

1. Sapporo
2. Nishikanbara-gun
3. Hachijyo-Island
4. Katsuyama
5. Toyono-gun
6. Yatsuka-gun
7. Shobara
8. Kochi
9. Kasuya-gun
10. Aira-gun



(9)-2 Strontium-90 and Cesium-137 in Milk (producing districts for domestic program)
(from Feb. 1979 to Oct. 1979)

— continued from No. 51 of this publication —

Table 9-2: Strontium-90 and Cesium-137 in Milk

Location	Ash (g/p/d)	Ca (mg/p/d)	K (mg/p/d)	⁹⁰ Sr		¹³⁷ Cs	
				pCi/p/d	S.U.	pCi/p/d	C.U.
February, 1979							
Aomori, AOMORI	7.11	1.10	1.56	2.3 ± 0.33	2.1 ± 0.30	11 ± 0.4	6.8 ± 0.26
Hakui-gun, ISHIKAWA	6.52	1.08	1.55	2.5 ± 0.31	2.3 ± 0.29	10 ± 0.4	6.5 ± 0.27
Akashi, HYOGO	7.10	1.12	1.62	1.9 ± 0.30	1.7 ± 0.27	1.8 ± 0.21	1.1 ± 0.13
Matsuyama, EHIME	7.28	1.18	1.58	1.4 ± 0.25	1.2 ± 0.21	2.2 ± 0.23	1.4 ± 0.15
June, 1979							
Saga-gun, SAGA	7.12	1.07	1.56	1.6 ± 0.25	1.5 ± 0.23	3.2 ± 0.26	2.0 ± 0.17
August, 1979							
Aomori, AOMORI	7.06	1.04	1.54	20 ± 0.6	19 ± 0.6	12 ± 0.4	7.7 ± 0.27
Mito, IBARAGI	7.34	1.14	1.45	2.8 ± 0.31	2.4 ± 0.27	2.5 ± 0.25	1.7 ± 0.17
Hakui-gun, ISHIKAWA	6.73	0.934	1.76	1.9 ± 0.24	2.1 ± 0.26	7.1 ± 0.34	4.0 ± 0.19
Himeji, HYOGO	6.51	1.02	1.47	1.7 ± 0.27	1.7 ± 0.26	2.3 ± 0.21	1.5 ± 0.14
Matsuyama, EHIME	7.47	1.09	1.64	1.0 ± 0.23	0.9 ± 0.21	2.1 ± 0.23	1.3 ± 0.14
October, 1979							
Saga-gun, SAGA	7.44	1.11	1.58	1.6 ± 0.28	1.4 ± 0.25	2.1 ± 0.24	1.3 ± 0.15

(9)-3 Strontium-90 and Cesium-137 in Milk (consuming districts)
(from Oct. 1978 to Oct. 1979)

— continued from No. 51 of this publication —

Table 9-3: Strontium-90 and Cesium-137 in Milk


Location	Ash (g/p/d)	Ca (mg/p/d)	K (mg/p/d)	⁹⁰ Sr		¹³⁷ Cs	
				pCi/p/d	S.U.	pCi/p/d	C.U.
October, 1978							
Kyoto, KYOTO	7.10	1.05	1.53	1.6 ± 0.27	1.5 ± 0.26	2.8 ± 0.25	1.8 ± 0.17
December, 1978							
Sendai, MIYAGI	7.29	1.10	1.65	1.4 ± 0.30	1.3 ± 0.27	4.7 ± 0.30	2.9 ± 0.18
Matsue, SHIMANE	7.11	1.08	1.52	2.1 ± 0.31	1.9 ± 0.29	3.9 ± 0.27	2.6 ± 0.18

Location	Ash	Ca	K	⁹⁰ Sr		¹³⁷ Cs	
	(g/p/d)	(mg/p/d)	(mg/p/d)	pCi/p/d	S.U.	pCi/p/d	C.U.
January, 1979							
Osaka, OSAKA	7.35	1.09	1.60	1.5 ± 0.30	1.3 ± 0.28	2.9 ± 0.27	1.8 ± 0.17
Wakayama, WAKAYAMA	7.82	1.07	1.80	2.1 ± 0.33	1.9 ± 0.31	2.3 ± 0.25	1.3 ± 0.14
Hiroshima, HIROSHIMA	6.56	0.999	1.45	1.0 ± 0.22	1.0 ± 0.22	2.7 ± 0.22	1.9 ± 0.15
Nagasaki, NAGASAKI	7.10	1.07	1.59	1.7 ± 0.25	1.6 ± 0.23	3.2 ± 0.26	2.0 ± 0.16
February, 1979							
Akita, AKITA	7.56	1.04	1.45	3.1 ± 0.34	3.0 ± 0.33	5.6 ± 0.33	3.8 ± 0.23
Yamagata, YAMAGATA	6.31	0.935	1.41	1.9 ± 0.26	2.1 ± 0.28	3.5 ± 0.25	2.5 ± 0.18
Fukushima, FUKUSHIMA	9.47	1.44	2.16	2.4 ± 0.36	1.6 ± 0.25	6.2 ± 0.38	2.9 ± 0.17
Yokohama, KANAGAWA	6.88	1.03	1.48	0.96 ± 0.24	0.93 ± 0.23	4.1 ± 0.28	2.8 ± 0.19
Niigata, NIIGATA	8.07	1.15	1.69	3.1 ± 0.35	2.7 ± 0.30	6.8 ± 0.37	4.1 ± 0.22
Fukui, FUKUI	7.13	1.11	1.53	1.3 ± 0.26	1.2 ± 0.23	3.5 ± 0.27	2.3 ± 0.18
Nagano, NAGANO	6.62	0.989	1.47	1.2 ± 0.23	1.3 ± 0.23	2.9 ± 0.23	2.0 ± 0.16
Shizuoka, SHIZUOKA	7.03	1.10	1.57	3.3 ± 0.35	3.0 ± 0.31	11 ± 0.4	6.7 ± 0.25
Nagoya, AICHI	7.06	1.04	1.52	1.7 ± 0.30	1.7 ± 0.29	8.5 ± 0.36	5.6 ± 0.24
Yonago, TOTTORI	6.90	1.08	1.48	2.0 ± 0.27	1.9 ± 0.25	10 ± 0.40	6.8 ± 0.27
Okayama, OKAYAMA	7.27	1.11	1.62	1.7 ± 0.28	1.5 ± 0.25	2.7 ± 0.25	1.7 ± 0.16
Matsuyama, EHIME	7.25	1.07	1.61	2.0 ± 0.27	1.9 ± 0.25	2.4 ± 0.24	1.5 ± 0.15
Kochi, KOCHI	7.33	1.12	1.62	2.6 ± 0.34	2.3 ± 0.31	3.7 ± 0.27	2.3 ± 0.17
Tsukushino, FUKUOKA	7.29	1.10	1.63	1.5 ± 0.28	1.4 ± 0.26	3.0 ± 0.27	1.9 ± 0.16
Kagoshima, KAGOSHIMA	7.02	1.10	1.59	2.4 ± 0.30	2.1 ± 0.27	8.6 ± 0.38	5.4 ± 0.24
Katsuren, OKINAWA	6.71	0.840	1.50	0.69 ± 0.23	0.83 ± 0.27	2.5 ± 0.24	1.6 ± 0.16
March, 1979							
Shinjuku, TOKYO	7.15	1.12	1.67	1.5 ± 0.26	1.4 ± 0.23	4.6 ± 0.30	2.8 ± 0.18
Yamaguchi, YAMAGUCHI	6.79	1.29	1.56	1.6 ± 0.31	1.2 ± 0.24	3.4 ± 0.30	2.2 ± 0.19
June, 1979							
Matsue, SHIMANE	7.13	1.08	1.56	2.4 ± 0.29	2.2 ± 0.27	7.8 ± 0.37	5.0 ± 0.23
Kyoto, KYOTO	6.98	1.04	1.54	1.4 ± 0.26	1.4 ± 0.25	2.9 ± 0.27	1.9 ± 0.18
July, 1979							
Akita, AKITA	8.04	1.02	1.72	3.1 ± 0.37	3.1 ± 0.36	6.7 ± 0.35	3.9 ± 0.20
Kagoshima, KAGOSHIMA	6.92	1.02	1.55	2.3 ± 0.27	2.3 ± 0.26	7.8 ± 0.34	5.0 ± 0.22
August, 1979							
Sapporo, HOKKAIDO	7.10	1.08	1.56	2.9 ± 0.32	2.7 ± 0.30	8.1 ± 0.36	5.2 ± 0.23
Sendai, MIYAGI	7.02	1.08	1.58	1.1 ± 0.22	1.1 ± 0.20	3.4 ± 0.26	2.2 ± 0.17
Yamagata, YAMAGATA	6.35	0.918	1.44	1.7 ± 0.25	1.9 ± 0.27	3.7 ± 0.25	2.6 ± 0.17
Fukushima, FUKUSHIMA	7.23	1.07	1.64	1.8 ± 0.27	1.7 ± 0.25	4.7 ± 0.32	2.9 ± 0.19
Niigata, NIIGATA	7.65	1.14	1.69	2.3 ± 0.28	2.0 ± 0.24	4.4 ± 0.29	2.6 ± 0.17
Fukui, FUKUI	6.99	1.06	1.59	1.0 ± 0.26	0.9 ± 0.24	2.7 ± 0.23	1.7 ± 0.15
Nagano, NAGANO	7.03	1.07	1.62	1.5 ± 0.23	1.4 ± 0.22	5.0 ± 0.28	3.1 ± 0.17
Shizuoka, SHIZUOKA	7.04	1.06	1.66	0.6 ± 0.25	0.5 ± 0.23	2.1 ± 0.21	1.2 ± 0.13

Location	Ash	Ca	K	⁹⁰ Sr		¹³⁷ Cs	
	(g/p/d)	(mg/p/d)	(mg/p/d)	pCi/p/d	S.U.	pCi/p/d	C.U.
Nagoya, AICHI	6.70	0.989	1.47	1.5 ± 0.23	1.5 ± 0.23	2.2 ± 0.23	1.5 ± 0.15
Osaka, OSAKA	7.03	1.05	1.59	1.4 ± 0.24	1.4 ± 0.23	3.6 ± 0.26	2.3 ± 0.16
Wakayama, WAKAYAMA	6.75	1.04	1.58	1.2 ± 0.21	1.1 ± 0.21	2.3 ± 0.21	1.4 ± 0.13
Yonago, TOTTORI	6.78	1.03	1.57	2.7 ± 0.27	2.7 ± 0.26	8.7 ± 0.35	5.6 ± 0.23
Okayama, OKAYAMA	6.76	1.02	1.55	1.2 ± 0.23	1.2 ± 0.22	2.4 ± 0.23	1.5 ± 0.15
Hiroshima, HIROSHIMA	6.87	1.04	1.60	1.2 ± 0.23	1.2 ± 0.22	1.8 ± 0.21	1.1 ± 0.13
Matsuyama, EHIME	7.10	1.03	1.54	1.1 ± 0.24	1.1 ± 0.23	2.1 ± 0.23	1.4 ± 0.15
Kochi, KOCHI	6.96	1.06	1.58	1.4 ± 0.27	1.3 ± 0.26	1.6 ± 0.19	1.0 ± 0.12
Tsukushino, FUKUOKA	7.29	1.10	1.68	1.5 ± 0.26	1.4 ± 0.24	4.0 ± 0.28	2.4 ± 0.17
September, 1979							
Yamaguchi, YAMAGUCHI	7.19	1.08	1.61	1.9 ± 0.26	1.7 ± 0.24	2.7 ± 0.26	1.7 ± 0.16
Nagasaki, NAGASAKI	6.76	1.02	1.53	1.0 ± 0.20	1.0 ± 0.20	8.9 ± 0.38	5.8 ± 0.25
October, 1979							
Yokohama, KANAGAWA	7.17	1.07	1.55	1.8 ± 0.30	1.7 ± 0.28	16 ± 0.6	10 ± 0.4

Figure 9-2, -3 Sampling location of Milk

- | | |
|---------------|----------------|
| 1. Aomori | 21. Wakayama |
| 2. Mito | 22. Yonago |
| 3. Hakui-gun | 23. Matsue |
| 4. Akashi | 24. Okayama |
| 5. Matsuyama | 25. Hiroshima |
| 6. Saga-gun | 26. Yamaguchi |
| 7. Sapporo | 27. Kochi |
| 8. Akita | 28. Tsukushino |
| 9. Sendai | 29. Nagasaki |
| 10. Yamagata | 30. Kagoshima |
| 11. Fukushima | 31. Katsuren |
| 12. Shinjuku | |
| 13. Yokohama | |
| 14. Niigata | |
| 15. Fukui | |
| 16. Nagano | |
| 17. Shizuoka | |
| 18. Nagoya | |
| 19. Kyoto | |
| 20. Osaka | |



(9)-4 Strontium-90 and Cesium-137 in Milk (powdered milk)
(from May, 1979 to Dec. 1979)

— continued from No. 51 of this publication —

Table 9-4: Strontium-90 and Cesium-137 in Milk

Manufacturer	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
May, 1979							
*Morinaga	8.23	1.27	1.85	33 ± 1.3	2.6 ± 0.10	69 ± 1.5	3.7 ± 0.08
Morinaga	2.57	0.326	0.604	17 ± 0.8	5.2 ± 0.23	96 ± 1.4	16 ± 0.2
Yukijirushi	2.40	0.362	0.478	14 ± 0.7	3.8 ± 0.18	51 ± 1.0	11 ± 0.2
Meiji	3.07	0.467	0.682	22 ± 0.8	4.7 ± 0.18	83 ± 1.3	12 ± 0.2
Wakodo	2.39	0.323	0.590	6.5 ± 0.51	2.0 ± 0.16	28 ± 0.8	4.8 ± 0.13
July, 1979							
*Meiji	7.93	1.23	1.79	74 ± 1.8	6.0 ± 0.15	130 ± 2	7.2 ± 0.11
December, 1979							
*Morinaga	8.28	1.25	1.96	32 ± 1.3	2.5 ± 0.10	60 ± 1.4	3.0 ± 0.07
Morinaga	2.58	0.325	0.655	12 ± 0.6	3.6 ± 0.19	83 ± 1.3	13 ± 0.2
Yukijirushi	2.41	0.328	0.494	12 ± 0.6	3.6 ± 0.19	48 ± 0.9	9.7 ± 0.19
Meiji	2.50	0.418	0.588	14 ± 0.7	3.3 ± 0.17	65 ± 1.1	11 ± 0.2
Wakodo	2.34	0.321	0.625	4.9 ± 0.48	1.5 ± 0.15	15 ± 0.5	2.5 ± 0.09
*Meiji	8.02	1.23	1.93	55 ± 1.7	4.5 ± 0.14	290 ± 3	15 ± 0.2

* Skim milk

(10) Strontium-90 and Cesium-137 in Vegetables
(from Nov. 1978 to Dec. 1979)

— continued from No. 51 of this publication —

Table 10: Strontium-90 and Cesium-137 in Vegetables

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
Japanese radish							
November, 1978							
Sannohe, AOMORI	0.567	0.031	0.252	19 ± 0.6	60 ± 1.9	2.9 ± 0.25	1.2 ± 0.10
Fukushima, FUKUSHIMA	0.439	0.028	0.170	2.9 ± 0.25	10 ± 0.9	0.4 ± 0.11	0.3 ± 0.06
Niigata, NIIGATA	0.438	0.021	0.194	2.5 ± 0.22	12 ± 1.1	0.7 ± 0.12	0.4 ± 0.06
Fukui, FUKUI	0.545	0.030	0.249	4.5 ± 0.32	15 ± 1.1	0.3 ± 0.14	0.1 ± 0.06
Okayama, OKAYAMA	0.668	0.032	0.276	1.5 ± 0.26	4.7 ± 0.79	0.3 ± 0.16	0.1 ± 0.06


Location	Component			⁹⁰ Sr				¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg		S.U.		pCi/kg	C.U.
Kaimon, KAGOSHIMA	0.666	0.041	0.284	14	± 0.6	34	± 1.4	4.9 ± 0.30	1.7 ± 0.10
December, 1978									
Wakayama, WAKAYAMA	0.764	0.043	0.293	5.6	± 0.42	13	± 1.0	0.5 ± 0.19	0.2 ± 0.06
Tottori, TOTTORI	0.622	0.028	0.282	11	± 0.5	41	± 1.8	0.5 ± 0.17	0.2 ± 0.06
Saga, SAGA	0.547	0.032	0.259	7.3	± 0.40	23	± 1.3	0.6 ± 0.16	0.2 ± 0.06
January, 1979									
Kochi, KOCHI	0.561	0.048	0.223	17	± 0.6	35	± 1.2	0.7 ± 0.15	0.3 ± 0.07
Nagasaki, NAGASAKI	0.465	0.035	0.172	6.4	± 0.32	18	± 0.9	—	—
February, 1979									
Yokohama, KANAGAWA	0.449	0.023	0.195	5.3	± 0.32	23	± 1.4	0.1 ± 0.11	0.1 ± 0.06
Hiroshima, HIROSHIMA	0.559	0.033	0.236	5.4	± 0.37	16	± 1.1	0.4 ± 0.13	0.2 ± 0.06
Otsu-gun, YAMAGUCHI	0.599	0.036	0.279	7.8	± 0.44	22	± 1.2	0.0 ± 0.14	0.0 ± 0.05
Katsuren, OKINAWA	0.637	0.035	0.260	2.5	± 0.31	7.1	± 0.88	0.4 ± 0.16	0.1 ± 0.06
May, 1979									
Atsumi-gun, AICHI	0.457	0.015	0.209	2.8	± 0.25	18	± 1.6	0.2 ± 0.13	0.1 ± 0.06
July, 1979									
Ota, SHIMANEANE	0.715	0.019	0.418	20	± 0.7	110	± 4	2.9 ± 0.27	0.7 ± 0.07
September, 1979									
Ishikari-gun, HOKKAIDO	0.609	0.023	0.286	71	± 1.2	310	± 5	0.8 ± 0.18	0.3 ± 0.06
October, 1979									
Sendai, MIYAGI	0.889	0.070	0.369	7.9	± 0.52	11	± 0.7	0.4 ± 0.20	0.1 ± 0.05
Yamagata, YAMAGATA	0.588	0.024	0.273	12	± 0.8	49	± 3.3	0.4 ± 0.27	0.1 ± 0.10
November, 1979									
Shinjuku, TOKYO	0.448	0.028	0.172	3.8	± 0.47	14	± 1.7	0.2 ± 0.21	0.1 ± 0.12
Niigata, NIIGATA	0.469	0.014	0.208	2.8	± 0.26	20	± 1.9	1.0 ± 0.14	0.5 ± 0.07
Saku, NAGANO	0.511	0.032	0.183	5.1	± 0.54	16	± 1.7	0.9 ± 0.26	0.5 ± 0.14
Gotenba, SHIZUOKA	0.520	0.023	0.241	23	± 0.6	99	± 2.7	5.5 ± 0.25	2.3 ± 0.10
Osaka, OSAKA	0.513	0.020	0.226	9.7	± 0.49	49	± 2.5	0.4 ± 0.15	0.2 ± 0.07
Okayama, OKAYAMA	0.607	0.025	0.266	5.5	± 0.37	22	± 1.5	0.5 ± 0.14	0.2 ± 0.05
Fukuoka, FUKUOKA	0.677	0.033	0.299	5.7	± 0.40	17	± 1.2	0.4 ± 0.15	0.1 ± 0.05
Kagoshima, KAGOSHIMA	0.623	0.035	0.244	6.5	± 0.48	18	± 1.4	5.4 ± 0.30	0.2 ± 0.12
Akashi, HYOGO	0.554	0.023	0.227	7.3	± 0.36	32	± 1.6	0.4 ± 0.13	0.2 ± 0.06
Spinach									
November, 1978									
Fukushima, FUKUSHIMA	1.53	0.056	0.657	9.3	± 0.71	17	± 1.3	1.6 ± 0.32	0.2 ± 0.05
Fukui, FUKUI	1.53	0.067	0.685	5.8	± 0.56	8.7	± 0.83	1.1 ± 0.29	0.2 ± 0.04
Okayama, OKAYAMA	1.54	0.137	0.596	6.1	± 0.61	4.4	± 0.44	1.3 ± 0.29	0.2 ± 0.05
Matsuyama, EHIME	1.68	0.056	0.639	3.4	± 0.50	6.1	± 0.89	2.1 ± 0.34	0.3 ± 0.05
Matsuyama, EHIME	1.48	0.059	0.626	3.1	± 0.44	5.2	± 0.73	1.1 ± 0.27	0.2 ± 0.04

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
Kagoshima, KAGOSHIMA	1.91	0.169	0.772	52 ± 1.5	31 ± 0.9	30 ± 0.9	3.9 ± 0.12
December, 1978							
Saga, SAGA	1.75	0.096	0.697	9.5 ± 0.71	9.9 ± 0.74	2.2 ± 0.35	0.3 ± 0.05
January, 1979							
Kochi, KOCHI	1.49	0.075	0.608	16 ± 0.9	22 ± 1.2	2.2 ± 0.31	0.4 ± 0.05
Nagasaki, NAGASAKI	1.65	0.088	0.600	12 ± 0.9	13 ± 1.0	3.1 ± 0.48	0.5 ± 0.08
February, 1979							
Yokohama, KANAGAWA	1.41	0.048	0.604	7.5 ± 0.58	16 ± 1.2	1.2 ± 0.29	0.2 ± 0.05
Hiroshima, HIROSHIMA	1.28	0.039	0.533	4.3 ± 0.53	11 ± 1.4	1.7 ± 0.35	0.3 ± 0.07
Otsu, YAMAGUCHI	1.45	0.069	0.632	21 ± 1.0	30 ± 1.5	3.4 ± 0.37	0.5 ± 0.06
Katsuren, OKINAWA	1.43	0.055	0.402	0.00 ± 0.46	0.00 ± 0.84	0.7 ± 0.38	0.2 ± 0.09
April, 1979							
Sendai, MIYAGI	1.73	0.105	0.598	11 ± 0.9	10 ± 0.9	3.1 ± 0.49	0.5 ± 0.08
May, 1979							
Atsumi-gun, AICHI	1.17	0.064	0.455	2.9 ± 0.44	4.5 ± 0.69	1.1 ± 0.30	0.2 ± 0.07
June, 1979							
Niigata, NIIGATA	1.92	0.056	0.884	3.3 ± 0.70	5.9 ± 1.3	1.1 ± 0.46	0.1 ± 0.05
July, 1979							
Ota, SHIMANE	1.56	0.037	0.735	11 ± 0.9	29 ± 2.5	3.2 ± 0.50	0.4 ± 0.07
September, 1979							
Ishikari-gun, HOKKAIDO	1.78	0.117	0.754	13 ± 0.7	11 ± 0.6	1.6 ± 0.28	0.2 ± 0.04
October, 1979							
Yamagata, YAMAGATA	1.36	0.048	0.582	4.2 ± 0.62	8.7 ± 1.3	1.4 ± 0.35	0.2 ± 0.06
November, 1979							
Nagano, NAGANO	1.96	0.116	0.707	8.9 ± 0.84	7.7 ± 0.72	0.9 ± 0.36	0.1 ± 0.05
Shizuoka, SHIZUOKA	1.07	0.026	0.514	7.9 ± 0.61	30 ± 2.3	0.8 ± 0.27	0.2 ± 0.05
Kyoto, KYOTO	1.53	0.119	0.436	9.6 ± 1.1	8.1 ± 0.94	2.4 ± 0.52	0.6 ± 0.12
Osaka, OSAKA	1.39	0.095	0.620	1.5 ± 0.53	1.6 ± 0.55	0.03 ± 0.32	0.01 ± 0.05
Tottori, TOTTORI	1.36	0.083	0.677	14 ± 1.1	17 ± 1.4	3.3 ± 0.48	0.5 ± 0.07
Okayama, OKAYAMA	1.30	0.058	0.540	3.7 ± 0.48	6.3 ± 0.82	0.6 ± 0.20	0.1 ± 0.04
Fukuoka, FUKUOKA	1.63	0.074	0.681	21 ± 0.9	28 ± 1.1	6.4 ± 0.43	0.9 ± 0.06
Kagoshima, KAGOSHIMA	0.866	0.064	0.305	3.1 ± 0.45	4.9 ± 0.70	23 ± 0.7	7.5 ± 0.22
December, 1979							
Shinjuku, TOKYO	1.20	0.073	0.405	4.4 ± 0.68	6.1 ± 0.94	8.3 ± 0.53	2.1 ± 0.13

Location	Component			⁹⁰ Sr		¹³⁷ Cs		
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.	
Cabbage								
June, 1979								
Akita, AKITA	0.653	0.047	0.277	14 ± 0.6	30 ± 1.2	1.4 ± 0.20	0.5 ± 0.07	
November, 1979								
Osaka, OSAKA	0.539	0.043	0.201	5.5 ± 0.56	13 ± 1.3	0.9 ± 0.28	0.5 ± 0.14	
Onion								
July, 1979								
Osaka, OSAKA	0.326	0.012	0.140	2.1 ± 0.29	18 ± 2.5	0.4 ± 0.18	0.3 ± 0.13	
Chinese cabbage								
Wakayama, WAKAYAMA	1.15	0.128	0.329	13 ± 0.8	10 ± 0.6	1.0 ± 0.27	0.3 ± 0.08	

Figure 10 Sampling Locations of Vegetables

- 1. Ishikari-gun
- 2. Sannohe
- 3. Sendai
- 4. Yamagata
- 5. Fukushima
- 6. Shinjuku
- 7. Yokohama
- 8. Niigata
- 9. Fukui
- 10. Saku
- 11. Gotenba
- 12. Atsumi-gun
- 13. Osaka
- 14. Wakayama
- 15. Akashi
- 16. Tottori
- 17. Ota ane
- 18. Okayama
- 19. Hiroshima
- 20. Otsu-gun
- 21. Matsuyama
- 22. Kochi
- 23. Fukuoka
- 24. Saga
- 25. Nagasaki
- 26. Kagoshima
- 27. Katsuren



(11) Strontium-90 and Cesium-137 in Tea (Japanese tea)
(from May, 1979 to Jun. 1979)


— continued from No. 51 of this publication —

Table 11: Strontium-90 and Cesium 137 in Tea

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
May, 1979							
Uji, KYOTO	4.70	0.292	1.83	48 ± 3.3	16 ± 1.1	49 ± 2.5	2.6 ± 0.14
June, 1979							
Yosa-gun, KYOTO	5.40	0.391	2.01	120 ± 6	30 ± 1.4	180 ± 5	9.1 ± 0.25
Chiran (Kawabe-gun), KAGOSHIMA	4.87	0.289	1.99	63 ± 3.7	22 ± 1.3	230 ± 5	11 ± 0.3
Miyanojyo (Satsuma-gun), KAGOSHIMA	5.69	0.367	2.09	61 ± 4.1	17 ± 1.1	140 ± 4	6.5 ± 0.21

Figure 11 Sampling Locations of Tea

1. Yosa-gun
2. Uji
3. Miyanojyo (Satsuma-gun)
4. Chiran (Kawabe-gun)



(12) Strontium-90 and Cesium-137 in Sea fish
(from Nov. 1978 to Jan. 1980)

— Continued from No. 51 of this publication —

Table 12: Strontium-90 and Cesium-137 in Sea fish


Location	Sampling Date	Component			⁹⁰ Sr		¹³⁷ Cs	
		Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
Limanda yokohamae								
Niigata, NIIGATA	Nov. 1978	3.81	22.2	10.7	1.0 ± 0.29	0.1 ± 0.03	4.8 ± 0.36	1.2 ± 0.09
Niigata, NIIGATA	Nov. 1979	3.40	28.4	9.24	1.9 ± 0.38	0.2 ± 0.04	2.8 ± 0.37	0.9 ± 0.15
Pleuronectidae								
Fukui, FUKUI	Nov. 1978	1.01	3.66	31.7	0.0 ± 0.30	0.0 ± 0.78	6.8 ± 0.43	2.1 ± 0.13
Hiroshima-bay, HIROSHIMA	Dec. 1978	2.64	22.9	13.0	1.0 ± 0.33	0.2 ± 0.05	4.3 ± 0.39	1.3 ± 0.11
Mutsu, AOMORI	Nov. 1979	1.21	4.82	29.5	1.1 ± 0.45	1.8 ± 0.75	4.7 ± 0.41	1.3 ± 0.11
Fukui, FUKUI	Nov. 1979	1.10	5.45	29.8	0.8 ± 0.41	1.4 ± 0.68	1.8 ± 0.7	5.5 ± 0.20
Sendai, MIYAGI	Feb. 1979	3.44	21.3	7.34	1.1 ± 0.39	0.2 ± 0.05	4.3 ± 0.41	1.7 ± 0.16
Sardinops melanosticta								
Nagano, NAGANO	Dec. 1978	2.24	28.2	9.53	0.3 ± 0.25	0.05 ± 0.04	4.3 ± 0.33	2.0 ± 0.15
Nagano, NAGANO	Dec. 1979	3.37	23.3	8.88	0.0 ± 0.27	0.0 ± 0.03	3.6 ± 0.35	1.1 ± 0.11
Pneumatophorus japonicus								
Tottori, TOTTORI	Feb. 1979	0.638	5.42	24.0	0.0 ± 0.25	0.0 ± 0.62	5.1 ± 0.32	2.9 ± 0.18
Matsuyama, EHIME	Aug. 1979	1.26	4.23	31.0	0.1 ± 0.21	0.2 ± 0.39	8.7 ± 0.42	2.2 ± 0.11
Osaka, OSAKA	Nov. 1979	2.31	22.2	11.9	0.1 ± 0.30	0.01 ± 0.06	7.1 ± 0.47	2.6 ± 0.17
Kyoto, KYOTO	Nov. 1979	2.77	19.3	14.0	0.0 ± 0.26	0.0 ± 0.05	9.3 ± 0.47	2.2 ± 0.11
Mugil cephalus								
Okayama, OKAYAMA	Nov. 1978	1.16	1.75	31.2	0.6 ± 0.44	2.5 ± 1.9	5.1 ± 0.44	1.3 ± 0.11
Katsuren, OKINAWA	Mar. 1979	6.59	19.8	5.63	2.0 ± 0.4	0.2 ± 0.03	4.5 ± 0.46	1.2 ± 0.12
Okayama, OKAYAMA	Nov. 1979	1.18	4.89	31.4	1.6 ± 0.56	2.7 ± 0.94	4.5 ± 0.41	1.2 ± 0.11
Ariake, SAGA	Sep. 1979	1.49	8.77	25.9	0.6 ± 0.29	0.4 ± 0.20	4.7 ± 0.40	1.1 ± 0.09
Hexagrammos otakii								
Yamaguchi, YAMAGUCHI	Mar. 1979	3.33	22.1	6.29	1.0 ± 0.34	0.1 ± 0.04	4.1 ± 0.40	1.6 ± 0.15
Stolephorus japonicus								
Kagoshima, KAGOSHIMA	Dec. 1978	2.92	20.6	11.4	0.5 ± 0.30	0.1 ± 0.05	11 ± 0.5	3.1 ± 0.14
Kagoshima, KAGOSHIMA	Dec. 1979	2.87	20.0	10.6	0.6 ± 0.29	0.1 ± 0.05	7.8 ± 0.45	2.3 ± 0.14
Arctoscopus japonicus								
	Dec. 1978	2.37	19.9	13.2	0.6 ± 0.29	0.1 ± 0.06	3.8 ± 0.36	1.2 ± 0.11
Sebastes inermis								
Fukushima, FUKUSHIMA	Mar. 1979	4.36	20.0	6.19	0.3 ± 0.34	0.04 ± 0.04	10 ± 0.5	3.8 ± 0.20
Naha, OKINAWA	Feb. 1979	3.32	17.5	8.04	0.5 ± 0.34	0.1 ± 0.06	6.4 ± 0.45	2.4 ± 0.17

Location	Sampling Date	Component			⁹⁰ Sr		¹³⁷ Cs	
		Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
<i>Gerres macrosoma</i> Katsuren, OKINAWA	Mar. 1979	4.13	18.2	6.61	0.9 ± 0.25	0.1 ± 0.03	6.2 ± 0.42	2.3 ± 0.15
<i>Scarots rubroviolaceus</i> Katsuren, OKINAWA	Mar. 1979	5.99	19.5	5.24	1.2 ± 0.31	0.1 ± 0.03	10 ± 0.6	3.1 ± 0.19
<i>Clupanodon thrissa</i> Katsuren, OKINAWA	Mar. 1979	3.83	20.3	7.46	0.8 ± 0.25	0.1 ± 0.03	3.2 ± 0.35	1.1 ± 0.12
<i>Lethrinus xanthochilus</i> Naha, OKINAWA	Mar. 1979	5.48	19.8	5.63	1.7 ± 0.31	0.2 ± 0.03	11 ± 0.6	3.7 ± 0.19
<i>Therapon theraps</i> Katsuren, OKINAWA	Mar. 1979	5.08	19.6	5.34	1.0 ± 0.28	0.1 ± 0.03	4.9 ± 0.42	1.8 ± 0.15
<i>Trachurus trachurus</i> Kobe, HYOGO	July 1979	1.43	9.48	24.2	0.2 ± 0.24	0.1 ± 0.18	9.4 ± 0.40	2.7 ± 0.1
Wakayama, WAKAYAMA	Sept. 1979	3.51	27.8	9.48	0.9 ± 0.33	0.1 ± 0.03	6.4 ± 0.44	1.6 ± 0.1
Odawara, KANAGAWA	Oct. 1979	3.41	24.4	11.6	1.0 ± 0.36	0.1 ± 0.04	6.4 ± 0.46	1.6 ± 0.1
Shizuoka, SHIZUOKA	Nov. 1979	3.45	27.1	9.59	2.7 ± 0.42	0.3 ± 0.04	9.5 ± 0.52	2.9 ± 0.16
<i>Decapterus</i> Miyake-Island, TOKYO	Sept. 1979	1.56	9.97	22.9	0.0 ± 0.29	0.0 ± 0.19	7.9 ± 0.47	2.2 ± 0.13
<i>Sebastes marmoratus</i> Hamada, SHIMANE	Aug. 1979	6.39	32.5	5.69	1.7 ± 0.45	0.1 ± 0.02	6.6 ± 0.52	1.8 ± 0.14
<i>Katsuwonus pelamis</i> Tosa, KOCHI	May 1979	1.29	1.36	26.4	0.4 ± 0.30	2.0 ± 1.4	2.3 ± 0.6	5.6 ± 0.15
<i>Chrysophrys major</i> Fukuoka, FUKUOKA	July 1979	1.36	3.81	32.3	0.4 ± 0.24	0.7 ± 0.45	8.4 ± 0.43	1.9 ± 0.10
<i>Argyrosomus argentatus</i> Nagasaki, NAGASAKI	Aug. 1979	0.932	5.69	27.1	0.0 ± 0.26	0.0 ± 0.49	7.0 ± 0.41	2.7 ± 0.16
<i>Oncorhynchus keta</i> Uraga-cho, HOKKAIDO	Oct. 1979	1.36	6.38	28.7	0.3 ± 0.28	0.3 ± 0.32	6.3 ± 0.39	1.6 ± 0.10
<i>Sillago sihame</i> Chita-gun, AICHI	June 1979	3.31	24.9	9.84	1.0 ± 0.30	0.1 ± 0.04	5.3 ± 0.34	1.6 ± 0.10
<i>Gadus macrocephalus</i> Oga, AKITA	Jan. 1980	1.53	4.25	24.9	1.1 ± 0.44	1.7 ± 0.65	25 ± 0.7	6.3 ± 0.19
<i>Doryteuthis bleekeri</i> Yamagata, YAMAGATA	Sept. 1979	1.25	0.79	21.3	0.1 ± 0.20	0.5 ± 1.9	1.6 ± 0.25	0.6 ± 0.09
<i>Seriola quinqueradiata</i> Hakui-gun, ISHIKAWA	Sept. 1979	0.993	2.24	29.4	0.0 ± 0.69	0.0 ± 2.8	12 ± 0.8	3.7 ± 0.26

Scientific name	English name	Japanese name
<i>Limanda yokohamae</i>	A flatfish	Makogarei
Pleuronectidae	Flatfish	Karei
<i>Sardinops melanosticta</i>	Sardine	Iwashi
<i>Pneumatophorus japonicus</i>	Mackerel	Saba
<i>Mugil cephalus</i>	Gray mullet	Bora
<i>Hexagrammos otakii</i>	Rock-trout	Ainame
<i>Stolephorus japonicus</i>	Kibinago	Kibinago
<i>Arctoscopus japonicus</i>	Hatahata	Hatahata
<i>Sebastes inermis</i>	Jacopever	Kurogara (Mebaru)
<i>Gerres macrosoma</i>	Big mouth majarras	Okuchisagi
<i>Scarots rubroviolaceus</i>	Parrotfish	Nagabudai
<i>Clupanodon thrissa</i>	Dizzard shad	Konoshiro
<i>Lethrinus xanthochilus</i>	Rudder fish	Muneakakuchibi
<i>Therapon theraps</i>		Kotohiki
<i>Trachurus trachurus</i>	Saurel	Aji
Decapterus	House-mackerel	Muroaji
<i>Sebastes marmoratus</i>	Scorpion-fish	Kasago
<i>Katsuwonus pelamis</i>	Bonito	Katsuo
<i>Chrysophrys major</i>	Sea bream	Tai
<i>Argyrosomus argentatus</i>	White croaker	Ishimochi (Guchi)
<i>Oncorhynchus keta</i>	Salmon	Sake
<i>Sillago sihame</i>	Sillago	Kisu
<i>Gadus macrocephalus</i>	Alaska codfish	Madara
<i>Doryteuthis bleekeri</i>	A squid	Yariika
<i>Seriola quinqueradiata</i>	Yellow-tail	Buri (Hukuragi)

Figure 12 Sampling Locations of Sea fish

- | | |
|-------------------|-------------------|
| 1. Uruga-cho | 16. Kobe |
| 2. Mutsu | 17. Tottori |
| 3. Oga | 18. Hamada |
| 4. Sendai | 19. Okayama |
| 5. Fukushima | 20. Hiroshima-bay |
| 6. Odawara | 21. Yamaguchi |
| 7. Niigata | 22. Matsuyama |
| 8. Fukui | 23. Tosa |
| 9. Nagano | 24. Fukuoka |
| 10. Shizuoka | 25. Ariake |
| 11. Miyake-Island | 26. Nagasaki |
| 12. Chita-gun | 27. Kagoshima |
| 13. Osaka | 28. Naha |
| 14. Kyota | 29. Katsuren |
| 15. Wakayama | |



**(13) Strontium-90 and Cesium-137 in Freshwater fish
(from Nov. 1978 to Dec. 1979)**

— Continued from No. 51 of this publication —


Table 13: Strontium-90 and Cesium-137 in Freshwater fish

Sampling Location	Date	Ash (%)	Component (% by weight)		⁹⁰ Sr (pCi/l)		¹³⁷ Cs (pCi/l)	
			Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
<i>Carassius carassius cuvieri</i>								
Niigata, NIIGATA	Nov. 1978	4.89	26.3	5.91	120 ± 2	8.3 ± 0.16	12 ± 0.5	3.8 ± 0.17
<i>Carassius auratus</i>								
mikata, FUKUI	Dec. 1978	0.992	3.83	30.1	2.3 ± 0.42	5.3 ± 0.98	15 ± 0.6	4.4 ± 0.18
Ishikari, HOKKAIDO	July 1979	3.87	22.6	6.69	40 ± 1.0	4.5 ± 0.12	7.8 ± 0.45	3.0 ± 0.17
Niigata, NIIGATA	Nov. 1979	4.41	31.0	7.08	91 ± 1.7	6.4 ± 0.12	7.7 ± 0.48	2.3 ± 0.15
Mikata, FUKUI	Dec. 1979	1.05	6.26	25.3	3.1 ± 0.42	3.4 ± 0.47	4.9 ± 0.36	1.3 ± 0.10
<i>Hypomesus olidus</i>								
Suwa-lake, NAGANO	Dec. 1978	2.13	20.5	14.5	7.3 ± 0.46	1.7 ± 0.10	6.7 ± 0.38	2.1 ± 0.12
Suwa-lake, NAGANO	Dec. 1979	2.21	22.3	12.8	5.9 ± 0.52	1.2 ± 0.10	6.8 ± 0.44	2.4 ± 0.16
<i>Cyprinus carpio</i>								
Akita, AKITA	July 1979	3.31	24.7	8.94	63 ± 1.2	7.4 ± 0.14	20 ± 0.7	6.4 ± 0.22
Fukushima, FUKUSHIMA	Oct. 1979	3.11	28.1	9.94	36 ± 1.2	4.1 ± 0.14	11 ± 0.5	3.5 ± 0.17
Shobara, HIROSHIMA	Nov. 1979	1.37	20.8	18.4	14 ± 0.6	4.8 ± 0.22	10 ± 0.5	3.9 ± 0.18

Scientific name	English name	Japanese name
<i>Carassius carassius cuvieri</i>	Crucian carp	Herabuna
<i>Carassius auratus</i>	a crucian carp	Funa
<i>Hypomesus olidus</i>	Pond-smelt	Wakasagi
<i>Cyprinus carpio</i>	Carp	Koi

Figure 13 Sampling Locations of Freshwater fish

- 1. Ishikari
- 2. Akita
- 3. Fukushima
- 4. Niigata
- 5. Nagano
- 6. Mikata
- 7. Shobara



(14) Strontium-90 and Cesium-137 in Shellfish
(from Nov. 1978 to Nov. 1979)

– Continued from No. 51 of this publication –


Table 14: Strontium-90 and Cesium-137 in Shellfish

Location	Date	Component			⁹⁰ Sr		¹³⁷ Cs	
		Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
Pecten yessoensis								
Mutsu-bay, AOMORI	Nov. 1978	1.10	1.49	24.8	0.0 ± 0.30	0.0 ± 1.8	1.6 ± 0.30	0.6 ± 0.11
" "	Nov. 1979	1.27	2.00	19.9	0.0 ± 0.47	0.0 ± 1.8	2.1 ± 0.35	0.8 ± 0.13
Ostrea gigas								
Hiroshima-bay, HIROSHIMA	Jan. 1979	3.02	15.2	6.83	0.4 ± 0.34	0.1 ± 0.07	2.2 ± 0.35	1.0 ± 0.17
Venerupis philippinarum								
Kitatakaki-gun, NAGASAKI	July 1979	1.73	8.52	9.72	0.0 ± 0.48	0.0 ± 0.32	0.7 ± 0.41	0.4 ± 0.24
Chita-gun, AICHI	June 1979	1.31	4.77	18.6	0.4 ± 0.46	0.6 ± 0.73	1.5 ± 0.34	0.6 ± 0.14
Turbo cornutus								
Sakata, YAMAGATA	July 1979	1.65	6.44	14.0	0.2 ± 0.51	0.2 ± 0.48	2.0 ± 0.45	0.8 ± 0.19
Ryozu, NIIGATA	May 1979	2.00	2.43	12.0	0.0 ± 1.1	0.0 ± 2.3	2.2 ± 0.91	0.8 ± 0.35
Hakui-gun, ISHIKAWA	July 1979	2.88	5.12	8.08	0.5 ± 0.33	0.3 ± 0.22	2.1 ± 0.36	0.9 ± 0.15

Scientific name	English name	Japanese name
Pecten yessoensis	Scallop	Hotategai
Ostrea gigas	Ouster	Kaki
Venerupis philippinarum	Short-necked clam	Asari
Turbo cornutus	Wreath shell	Sazae

Figure 14 Sampling Locations of Shellfish

1. Mutsu-bay
2. Sakata
3. Ryozu
4. Hakui-gun
5. Chita-gun
6. Hiroshima-bay
7. Kitatakaki-gun



**(15) Strontium-90 and Cesium-137 in Seaweeds
(from Jan. 1979 to Jun. 1979)**

– Continued from No. 51 of this publication –


Table 15: Strontium-90 and Cesium-137 in Seaweeds

Location	Date	Component			⁹⁰ Sr		¹³⁷ Cs	
		Ash (%)	Ca (%)	K (%)	pCi/kg	S.U.	pCi/kg	C.U.
<i>Undaria pinnatifida</i>								
Chita-gun, AICHI	Jan. 1979	1.56	2.40	27.0	0.8 ± 0.27	2.0 ± 0.70	0.8 ± 0.20	0.2 ± 0.05
Hiroshima, HIROSHIMA	Feb. 1979	3.11	1.86	19.7	0.4 ± 0.33	0.7 ± 0.57	1.3 ± 0.27	0.2 ± 0.04
Shimabara, NAGASAKI	Jan. 1979	2.08	3.26	35.2	1.9 ± 0.41	2.8 ± 0.60	1.4 ± 0.29	0.2 ± 0.04
Sakai, YAMAGATA	June 1979	1.10	6.41	14.6	2.6 ± 0.38	3.6 ± 0.53	0.9 ± 0.23	0.5 ± 0.14
Ryozu, NIIGATA	May 1979	1.74	5.43	23.3	4.6 ± 0.43	4.8 ± 0.46	0.9 ± 0.23	0.2 ± 0.06
Hakui-gun, ISHIKAWA	April 1979	3.64	2.24	22.5	2.3 ± 0.35	2.8 ± 0.42	2.6 ± 0.30	0.3 ± 0.04

Scientific name	English name	Japanese name
<i>Undaria pinnatifida</i>	Wakame seaweed	Wakame

Figure 15 Sampling Locations of Seaweeds

1. Sakata
2. Ryozu
3. Haku-gun
4. Chita-gun
5. Hiroshima-bay
6. Shimabara



Edited by National Institute of Radiological Sciences, under the supervision of Science and Technology Agency of Japanese Government.