

ISSN 0441-2516
NIRS-RSD-67

**RADIOACTIVITY
SURVEY DATA
in Japan**

**NUMBER 67
Dec. 1983**

**National Institute of Radiological Sciences
Chiba, Japan**

Radioactivity Survey Data in Japan

Number 67

December 1983

Contents

	Page
Environmental and Dietary Materials	1
<i>(Japan Chemical Analysis Center)</i>	
1. Collection and pretreatment of samples	1
2. Preparation of samples for analysis	3
3. Separation of Strontium-90 and Cesium-137	3
4. Determination of stable strontium, calcium and potassium	4
5. Counting	4
6. Results	5
(1) Strontium-90 and Cesium-137 in Total diet	5
(2)-1 Strontium-90 and Cesium-137 in Rice	7
(producing districts)	
-2 Strontium-90 and Cesium-137 in Rice	8
(consuming districts)	
(3)-1 Strontium-90 and Cesium-137 in Milk	10
(producing districts for WHO program)	
-2 Strontium-90 and Cesium-137 in Milk	12
(consuming districts)	
-3 Strontium-90 and Cesium-137 in Milk	13
(producing districts for Domestic program)	
-4 Strontium-90 and Cesium-137 in Milk	15
(powder milk)	
(4)-1 Strontium-90 and Cesium-137 in Vegetables	16
(producing districts)	
-2 Strontium-90 and Cesium-137 in Vegetables	18
(consuming districts)	
(5) Strontium-90 and Cesium-137 in Tea	20
(Japanese tea)	
(6) Strontium-90 and Cesium-137 in Sea fish	21
(7) Strontium-90 and Cesium-137 in Freshwater fish	24
(8) Strontium-90 and Cesium-137 in Shellfish	26
(9) Strontium-90 and Cesium-137 in Seaweeds	28

Environmental and Dietary Materials*

(Japan Chemical Analysis Center)

1. Collection and pretreatment of samples

(1) Rain and dry fallout

Rain and dry fallout was collected monthly on a sampling tray, approximately 5000 cm² in area, which was filled with water to a depth of 1 cm at the beginning of every month.

The sample was filtered after strontium and cesium carriers were added. The tray was washed with 5ℓ of distilled water and the washing was combined to the filtrate. The sample was passed through a cation exchange column (500 mℓ of Dowex 50W X8, 50 ~ 100 mesh, Na form) at a rate of 80 mℓ/min.

(2) Airborne dust

Airborne dust was collected by an electrostatic precipitator or a filter air sampler for every three months at a rate of more than 3000 m³ per month. The sampling was done 1 to 1.5 meters above the ground.

(3) Service water and freshwater

Service water, 100ℓ each, was collected at an intake of the water-treatment plant and at the tap after water was left running for five minutes. Water, to which added carriers of strontium and cesium immediately after sampling, was vigorously stirred and filtered. The subsequent process was the same as that described in the section (1). Freshwater was treated in the same way as the service water.

(4) Soil

Soil was collected from the location in the spacious and flat area without past disturbance on the surface caused by duststorms, inflow and outflow due to precipitation, and so on. Any places located under trees in a forest, in a stony area or inside of river banks were avoided. Soil was taken from two layers of different depths, 0 ~ 5 cm and 5 ~ 20 cm. In the course of air-drying, lumps were crushed by hand, and roots of plants and pebbles were removed. The soil was then passed through a 2 mm sieve to remove small gravels.

(5) Sea water

Sea water was collected at the fixed stations where the effect of terrestrial fresh water from rivers was expected to be negligibly small. A special consideration was also given to weather conditions. The sampling was carried out when there was no rainfall for the last few days. To prevent contamination, water samples were collected at the bow of a sampling boat just before she stood still by scooping surface water using a polyethylene bucket. Immediately after the collection, the samples were acidified to a pH lower than 3 by adding concentrated hydrochloric acid in a ratio of 1 mℓ to 1 ℓ of sea water, and then stored in 20 ℓ polyethylene containers. The sampling equipments as well as containers were thoroughly rinsed with dilute hydrochloric acid and then with distilled water before use. Two hundred milliliters of sea water was also collected at the same stations for the determination of chlorinity.

(6) Sea sediments

Sediment was collected in the same area as that for the sea water sample, taking the following criteria into account:

- a. The depth of water exceeds 1 m at low tide.
- b. No significant sedimental movement is observed in the vicinity of concern.
- c. Mud, silt and fine sand are preferable.

A conventional sediment sampling device was used for collecting the top few centimeters of surface sediment. Approximately 4 kg of the sample in wet weight was spread on a large porcelain dish and dried in an electric oven at 105 to 110 °C to a constant weight.

(7) Total diet

A full one day ordinary diet including three meals, water, tea and other in-between snacks for five persons was collected as a sample of "total diet". The sample in a large stainless steel pan was carbonized carefully by direct application of gas flame, and was transferred to a porcelain dish and then ashed at 500 °C in an electric muffle furnace.

(8) Rice

Polished rice was collected in producing districts at the harvest and in consuming areas when new crops were first put on sale. The sample was carbonized and ashed in a porcelain dish.

* Samples were sent to the Center from 32 contracted prefectures.

(9) Milk

Raw milk was collected in producing districts and commercial milk was purchased in consuming districts. Milk in a stainless steel pan or a porcelain dish was evaporated to dryness followed by carbonization and ashing.

(10) Vegetables

Spinach and Japanese radish were selected as the representatives for leaf vegetables and for non-starch roots, respectively. After removing soil, the edible part of vegetable sample was dried and carbonized in a stainless steel pan or a porcelain dish.

(11) Tea

Five hundred grams of manufactured green tea was collected, carbonized and ashed in a stainless steel pan or a porcelain dish.

(12) Fish, shellfish and seaweeds

a. Sea fish and freshwater fish

Fish was rinsed with water and blotted with a filter paper. Only the edible part was used in case of larger sized fish, and the whole part was used in case of smaller ones. Each sample was weighed and placed in a stainless steel pan or a porcelain dish. After carbonized, the sample was ashed in an electric muffle furnace.

b. Shellfish

Approximately 4 kg of shellfish including the shells was collected or purchased. After removing the shells, it was treated in the same way as that for the sea fish.

c. Seaweeds

Edible seaweeds were collected and rinsed with water to remove sand and other adhering matters on the surface. These were removed of excess water, weighed, dried and ashed.

Table 1 shows details of sample collection.

Table 1 Details of sample collection

Sample	Frequency of sampling	Quantity of sample
= Environmental materials =		
(1) Rain and dry fallout		
1 for domestic program	monthly	
2 for WHO program	monthly	
(2) Airborne dust	quarterly	>3000 m ³ /month
(3) Service water and freshwater		
1 Service water (source water)	semiyearly (June and December)	100 ℓ
2 Service water (tap water)	semiyearly (June and December)	100 ℓ
3 Freshwater	yearly (fishing season)	100 ℓ
(4) Soil		
1 0 ~ 5 cm	yearly (June or July)	4 kg
2 5 ~ 20 cm	yearly (June or July)	4 kg
(5) Sea water	yearly (July or August)	40 ℓ
(6) Sea sediments	yearly (July or August)	4 kg
= Dietary materials =		
(7) Total diet	semiyearly (June, November or December)	daily amount for 5 person
(8) Rice		
1 producing districts	yearly (harvesting season)	5 kg (polished rice)
2 consuming districts	yearly (harvesting season)	5 kg (polished rice)
(9) Milk		
1 producing districts for WHO program	quarterly (February, May, August and November)	3 ℓ
2 producing districts for domestic program	semiyearly (February and August)	3 ℓ

Sample	Frequency of sampling	Quantity of sample
3 consuming districts	semiyearly (February and August)	3 ℓ
4 powdered milk	semiyearly (April and October)	2 ~ 3 kg
(10) Vegetables		
1 producing districts	yearly (harvesting season)	4 kg
2 consuming districts	yearly (harvesting season)	4 kg
(11) Tea	yearly (the first harvesting season)	500 g (manufactured tea)
(12) Fish, shellfish, and seaweeds		
1 Sea fish	yearly (fishing season)	4 kg
2 Freshwater fish	yearly (fishing season)	4 kg
3 Shellfish	yearly (fishing season)	4 kg
4 Seaweeds	yearly (fishing season)	2 ~ 3 kg

2. Preparation of samples for analysis

(1) Rain, service water and freshwater

Strontium and cesium were eluted with hydrochloric acid from the cation exchange column. The residue of rain sample on the filter paper was ashed in an electric muffle furnace and the ash was dissolved in hydrochloric acid. The insoluble part was filtered and washed. The filtrate and the washings were combined to the previous eluate and used for radiochemical analysis.

(2) Soil

Air-dried soil was passed through a 20 mesh sieve. The sieved sample was heated, in the presence of strontium and cesium carriers, together with sodium hydroxide. The sample was then heated with hydrochloric acid and the insoluble part was filtered and washed. The combined solution of the filtrate and washings was used for radiochemical analysis.

(3) Sea sediments

After removal of pebbles, shells and other foreign matters, the sediment sample was dried in a hot-air oven and ground finely with a mortar. The sample was passed through a 20 mesh sieve. The further preparation of the sample was the same as that described in the section 2-(2).

(4) Rice

The ashed sample was pulverized with a porcelain mortar and passed through a 42 mesh sieve. The sieved sample to which both strontium and cesium carriers were added, was digested with hydrochloric

acid by heating. After the sample was heated again with nitric acid to dryness, strontium and cesium were extracted with hydrochloric acid and water. The insoluble part was filtered and washed. The filtrate and washings were combined for subsequent radiochemical analysis.

(5) Airborne dust, diet, milk, vegetable, fish and shellfish, seaweeds, tea, and others

These ashed samples were treated with the same procedure as that described in the section 2-(4).

3. Separation of strontium-90 and cesium-137

(1) Strontium-90

Sample solutions, prepared as in the foregoing sections 2-(1) through 2-(5), were neutralized with sodium hydroxide. After sodium carbonate was added, the precipitate of strontium and calcium carbonates was separated. The supernatant solution was retained for cesium-137 determination. The carbonates were dissolved in hydrochloric acid and calcium and strontium were precipitated as oxalates. The precipitate was dissolved in nitric acid and strontium was separated from calcium by successive fuming nitric acid separations. Iron scavenge was made after addition of ferric iron carrier followed by barium chromate separation after addition of barium carrier to remove radium, its daughters and lead. Strontium was recovered as carbonate, and the precipitate was dried and weighed to determine strontium recovery. The strontium carbonate was dissolved in hydrochloric acid and the iron carrier was added. The solution was allowed to stand

for two weeks for strontium-90 and yttrium-90 to attain equilibrium. The yttrium-90 was coprecipitated with ferric hydroxide and the precipitate was filtered off, washed and counted.

(2) Cesium-137

The supernatant separated from the strontium fraction in the solution was acidified with hydrochloric acid. While stirring the solution, cesium was adsorbed on ammonium molybdophosphate.

After filtered off and washed with dilute nitric acid, the precipitate was dissolved in 2.5N sodium hydroxide solution. Ammonia was removed completely from the solution by boiling. The solution was adjusted to pH 8.2 with hydrochloric acid and allowed to cool. Molybdenum hydroxide which came out in the solution, was filtered off and washed with water. In such circumstance that contamination by rubidium-87 was not negligible for the measurement of cesium-137, the following ion-exchange procedure was applied. A fixed amount of ferric chloride solution was added to the solution dissolved with 2.5N sodium hydroxide. Ammonia and molybdenum hydroxide were removed as described above. Ethylenediaminetetraacetic acid tetrasodium salt was added to the filtrate and washings. Cesium and rubidium were adsorbed on a cation exchange resin. Cesium was separated from rubidium by eluting with hydrochloric acid.

To this eluate or the filtrate and washings after removing molybdenum hydroxide, chloroplatinic acid solution was added to precipitate cesium. The precipitate was filtered onto a tared paper in a demountable filter and washed with water and then ethanol. After fixing the filter paper on a tared planchette and drying

it, the chemical yield of cesium was determined by weighing the precipitate with the planchette. Radioactivity from cesium-137 was measured for this precipitate.

4. Determination of stable strontium, calcium and potassium

A weighed amount of soil or sea sediment was treated under heating with sodium hydroxide and then with hydrochloric acid for extraction. A weighed aliquot of ashed samples of total diet, vegetables, milk, fish, shellfish or seaweeds was digested using hydrochloric acid or nitric acid, hydrofluoric acid being used when necessary. The extract was made up to an appropriate volume with dilute hydrochloric acid. The sample solution was analyzed for calcium by titration with standard potassium permanganate solution after separating calcium as oxalate. Atomic absorption spectroscopy was applied when appropriate. Stable strontium and potassium were determined by atomic absorption and flame emission spectrometry, respectively.

5. Counting

After the radiochemical separation, the mounted precipitates were counted for activity using low background beta counters normally for 60 min. Net sample counting rates were corrected for counter efficiency, recovery, self-absorption and decay to obtain the content of strontium-90 and cesium-137 radio activity per sample aliquot. From the results, concentrations of these nuclides in the original samples were calculated.

6. Results

(1) Strontium-90 and Cesium-137 in Total diet (from Jun. 1983 to Dec. 1983)

— continued from No. 65 of this publication —


Table (1): Strontium-90 and Cesium-137 in Total diet

Location	Ash	Ca	K	⁹⁰ Sr		¹³⁷ Cs	
	(g p. d ⁻¹)	(mg p. d ⁻¹)	(mg p. d ⁻¹)	pCi p. ⁻¹ d. ⁻¹	S.U.	pCi p. ⁻¹ d. ⁻¹	C.U.
June, 1983							
Aomori, AOMORI	17.3	375	1960	3.1 ± 0.34	8.3 ± 0.91	3.9 ± 0.31	2.0 ± 0.16
Fukushima, FUKUSHIMA	16.3	553	1620	1.9 ± 0.28	3.4 ± 0.51	1.3 ± 0.20	0.8 ± 0.12
Hiratsuka, KANAGAWA	17.8	704	2420	2.7 ± 0.33	3.8 ± 0.47	3.0 ± 0.30	1.3 ± 0.12
Shizuoka, SHIZUOKA	19.5	1010	2730	3.6 ± 0.39	3.5 ± 0.39	2.5 ± 0.31	0.9 ± 0.11
Kyoto, KYOTO	17.4	883	2320	3.3 ± 0.38	3.8 ± 0.43	2.1 ± 0.31	0.9 ± 0.13
Neyagawa, OSAKA	17.0	450	2410	3.3 ± 0.34	7.4 ± 0.77	2.1 ± 0.24	0.9 ± 0.10
Matsue, SHIMANE	20.1	918	2680	3.9 ± 0.41	4.2 ± 0.44	2.8 ± 0.33	1.1 ± 0.12
Dazaifu, FUKUOKA	12.6	431	1840	1.6 ± 0.23	3.8 ± 0.54	1.8 ± 0.21	1.0 ± 0.11
Saga, SAGA	16.8	906	1990	2.1 ± 0.29	2.3 ± 0.32	1.5 ± 0.23	0.8 ± 0.12
Nagasaki, NAGASAKI	16.4	717	1840	1.9 ± 0.30	2.6 ± 0.41	2.0 ± 0.23	1.1 ± 0.13
Soo-gun, KAGOSHIMA	16.3	538	2070	2.0 ± 0.30	3.6 ± 0.56	2.7 ± 0.27	1.3 ± 0.13
July, 1983							
Yamagata, YAMAGATA	19.5	682	2590	1.4 ± 0.30	2.1 ± 0.44	1.9 ± 0.28	0.7 ± 0.11
Matsuyama, EHIME	11.3	381	1460	1.4 ± 0.21	3.8 ± 0.55	0.9 ± 0.13	0.6 ± 0.09
Naha, OKINAWA	16.2	336	1600	1.6 ± 0.29	4.7 ± 0.87	1.5 ± 0.21	0.9 ± 0.13
August, 1983							
Hiroshima, HIROSHIMA	10.5	355	995	1.5 ± 0.19	4.1 ± 0.55	1.4 ± 0.16	1.4 ± 0.16
Yamaguchi, YAMAGUCHI	17.8	641	1980	2.3 ± 0.31	3.6 ± 0.49	1.8 ± 0.23	0.9 ± 0.12
September, 1983							
Ojika-gun, MIYAGI	16.6	625	2310	4.5 ± 0.38	7.1 ± 0.60	3.5 ± 0.32	1.5 ± 0.14
November, 1983							
Akita, AKITA	14.2	602	1950	3.6 ± 0.33	6.0 ± 0.54	5.5 ± 0.32	2.8 ± 0.16
Nishikawa, NIIGATA	21.5	641	2660	3.8 ± 0.39	6.0 ± 0.62	1.5 ± 0.23	0.6 ± 0.09
Kanazawa, ISHIKAWA	22.3	603	2780	4.5 ± 0.38	7.5 ± 0.64	3.2 ± 0.32	1.1 ± 0.12
Shizuoka, SHIZUOKA	20.1	717	2680	3.3 ± 0.32	4.6 ± 0.45	3.5 ± 0.31	1.3 ± 0.12
Nagoya, AICHI	14.2	420	2060	2.7 ± 0.29	6.3 ± 0.69	1.2 ± 0.21	0.6 ± 0.10

Location	Ash	Ca	K	^{90}Sr		^{137}Cs	
	($\text{g}\cdot\text{p}^{-1}\cdot\text{d}^{-1}$)	($\text{mg}\cdot\text{p}^{-1}\cdot\text{d}^{-1}$)	($\text{mg}\cdot\text{p}^{-1}\cdot\text{d}^{-1}$)	$\text{pCi}\cdot\text{p}^{-1}\cdot\text{d}^{-1}$	S.U.	$\text{pCi}\cdot\text{p}^{-1}\cdot\text{d}^{-1}$	C.U.
Okayama, OKAYAMA	15.3	608	1940	3.6 ± 0.34	5.9 ± 0.56	2.1 ± 0.25	1.1 ± 0.13
Kochi, KOCHI	18.3	718	2560	5.4 ± 0.43	7.5 ± 0.61	3.0 ± 0.33	1.2 ± 0.13
Saga, SAGA	20.2	1390	2080	2.7 ± 0.31	1.9 ± 0.22	1.6 ± 0.25	0.7 ± 0.12
Soo-gun, KAGOSHIMA	17.8	653	2150	2.9 ± 0.38	4.5 ± 0.58	2.0 ± 0.24	0.9 ± 0.11
December, 1983							
Sapporo, HOKKAIDO	16.5	528	2290	4.4 ± 0.40	8.3 ± 0.76	3.8 ± 0.29	1.7 ± 0.13
Mito, IBARAGI	14.7	454	1850	2.5 ± 0.30	5.5 ± 0.66	1.7 ± 0.20	0.9 ± 0.11
Kyoto, KYOTO	18.5	1110	2420	2.7 ± 0.37	2.4 ± 0.33	2.3 ± 0.27	0.9 ± 0.11
Kakogawa, HYOGO	13.1	741	1830	2.5 ± 0.26	3.3 ± 0.35	2.0 ± 0.22	1.1 ± 0.12
Wakayama, WAKAYAMA	11.0	548	1480	2.3 ± 0.24	4.3 ± 0.44	1.5 ± 0.16	1.0 ± 0.11

Figure (1) Sampling Locations of Total diet

- | | |
|--------------|---------------|
| 1. Sapporo | 15. Wakayama |
| 2. Aomori | 16. Kakogawa |
| 3. Akita | 17. Okayama |
| 4. Ojika-gun | 18. Matsue |
| 5. Yamagata | 19. Hiroshima |
| 6. Fukushima | 20. Yamaguchi |
| 7. Mito | 21. Matsuyama |
| 8. Nishikawa | 22. Kochi |
| 9. Hiratsuka | 23. Dazaifu |
| 10. Shizuoka | 24. Saga |
| 11. Nagoya | 25. Nagasaki |
| 12. Kanazawa | 26. Soo-gun |
| 13. Kyoto | 27. Naha |
| 14. Neyagawa | |



**(2)-1 Strontium-90 and Cesium-137 in Rice (producing districts)
(from Oct. 1983 to Dec. 1983)**


— continued from No. 65 of this publication —

Table (2)-1: Strontium-90 and Cesium-137 in Rice

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/Kg)	K (g/Kg)	pCi/Kg	S.U.	pCi/Kg	C.U.
October, 1983							
Nishikanbara-gun, NIIGATA	0.386	0.054	0.786	0.6 ± 0.14	10.0 ± 2.5	0.2 ± 0.10	0.2 ± 0.12
Minamiazumi-gun, NAGANO	0.587	0.072	1.07	0.0 ± 0.17	0.0 ± 2.4	0.7 ± 0.14	0.7 ± 0.13
November, 1983							
Ishikari, HOKKAIDO	0.474	0.059	1.10	0.5 ± 0.14	8.0 ± 2.4	0.4 ± 0.12	0.4 ± 0.11
Toda-gun, MIYAGI	0.535	0.058	0.882	0.3 ± 0.15	5.0 ± 2.6	1.0 ± 0.16	1.1 ± 0.18
Mito, IBARAGI	0.460	0.063	0.699	0.7 ± 0.15	10.0 ± 2.4	7.2 ± 0.28	10.0 ± 0.40
December, 1983							
Fukushima, FUKUSHIMA	0.600	0.074	0.972	0.3 ± 0.18	4.0 ± 2.4	2.0 ± 0.21	2.0 ± 0.22

Figure (2)-1 Sampling Locations of Rice

1. Ishikari
2. Toda-gun
3. Fukushima
4. Mito
5. Nishikanbara-gun
6. Minamiazumi-gun



**(2)-2 Strontium-90 and Cesium-137 in Rice (consuming districts)
(from Sept. 1983 to Dec. 1983)**


— continued from No. 65 of this publication —

Table (2)-2: Strontium-90 and Cesium-137 in Rice

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
September, 1983							
Kanazawa, ISHIKAWA	0.477	0.065	1.010	0.30 ± 0.15	5.0 ± 2.3	1.0 ± 0.15	1.0 ± 0.14
October, 1983							
Mito, IBARAGI	0.424	0.057	0.729	0.40 ± 0.13	8.0 ± 2.3	0.8 ± 0.13	1.2 ± 0.18
Shinjuku, TOKYO	0.416	0.061	0.911	0.50 ± 0.14	9.0 ± 2.3	0.7 ± 0.12	0.8 ± 0.14
Niigata, NIIGATA	0.365	0.059	0.710	0.60 ± 0.13	10.0 ± 2.3	0.6 ± 0.11	0.9 ± 0.16
Wakayama, WAKAYAMA	0.402	0.058	0.829	0.50 ± 0.13	8.0 ± 2.3	1.0 ± 0.13	1.2 ± 0.15
Hiroshima, HIROSHIMA	0.368	0.056	0.710	0.60 ± 0.13	10.0 ± 2.3	0.4 ± 0.11	0.6 ± 0.15
Matsuyama, EHIME	0.519	0.075	0.900	0.40 ± 0.17	5.0 ± 2.3	4.7 ± 0.26	5.2 ± 0.29
November, 1983							
Sapporo, HOKKAIDO	0.609	0.062	1.290	0.40 ± 0.17	7.0 ± 2.8	0.8 ± 0.17	0.6 ± 0.13
Akita, AKITA	0.460	0.060	0.855	1.00 ± 0.16	16.0 ± 2.7	2.3 ± 0.19	2.7 ± 0.22
Yamagata, YAMAGATA	0.563	0.073	1.180	0.40 ± 0.17	5.0 ± 2.3	0.5 ± 0.15	0.5 ± 0.13
Shizuoka, SHIZUOKA	0.540	0.069	0.955	0.10 ± 0.17	1.0 ± 2.4	1.2 ± 0.16	1.3 ± 0.16
Kyoto, KYOTO	0.485	0.059	1.04	0.03 ± 0.16	1.0 ± 2.7	0.6 ± 0.10	0.5 ± 0.10
Osaka, OSAKA	0.413	0.058	0.911	0.20 ± 0.13	3.0 ± 2.2	0.3 ± 0.10	0.3 ± 0.11
Saga, SAGA	0.525	0.072	0.929	0.40 ± 0.18	6.0 ± 2.5	0.3 ± 0.09	0.3 ± 0.10
Kagoshima, KAGOSHIMA	0.541	0.069	0.914	0.10 ± 0.17	2.0 ± 2.4	3.9 ± 0.24	4.2 ± 0.26
December, 1983							
Yokohama, KANAGAWA	0.426	0.063	0.719	0.30 ± 0.14	5.0 ± 2.2	1.2 ± 0.13	1.6 ± 0.18
Nagoya, AICHI	0.514	0.076	0.997	0.03 ± 0.16	0.0 ± 2.1	0.4 ± 0.10	0.4 ± 0.10
Kochi, KOCHI	0.477	0.056	0.944	0.10 ± 0.14	1.0 ± 2.5	1.0 ± 0.12	1.0 ± 0.13

Figure (2)-2 Sampling Locations of Rice

- | | |
|-------------|---------------|
| 1. Sapporo | 10. Kanazawa |
| 2. Akita | 11. Kyoto |
| 3. Yamagata | 12. Osaka |
| 4. Mito | 13. Wakayama |
| 5. Shinjuku | 14. Hiroshima |
| 6. Yokohama | 15. Matsuyama |
| 7. Niigata | 16. Kochi |
| 8. Shizuoka | 17. Saga |
| 9. Nagoya | 18. Kagoshima |



(3)-1 Strontium-90 and Cesium-137 in Milk (producing districts for WHO program)
(from Apr. 1983 to Dec. 1983)


— continued from No. 65 of this publication —

Table (3)-1: Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (g/l)	Ca (g/l)	K (g/l)	pCi/l	S. U.	pCi/l	C. U.
April, 1983							
Yatsuka-gun, SHIMANE	7.27	1.18	1.53	2.3 ± 0.29	1.9 ± 0.24	2.0 ± 0.22	1.3 ± 0.15
May, 1983							
Kasuya-gun, FUKUOKA	7.38	1.28	1.64	0.8 ± 0.24	0.6 ± 0.19	5.4 ± 0.32	3.3 ± 0.19
June, 1983							
Yatsuka-gun, SHIMANE	7.27	1.15	1.62	3.1 ± 0.32	2.7 ± 0.28	6.0 ± 0.33	3.7 ± 0.20
July, 1983							
Takada-gun, HIROSHIMA	6.97	1.10	1.50	1.2 ± 0.23	1.1 ± 0.21	0.9 ± 0.19	0.6 ± 0.13
August, 1983							
Sapporo, HOKKAIDO	7.57	1.35	1.60	1.9 ± 0.26	1.4 ± 0.19	3.1 ± 0.26	1.9 ± 0.16
Nishikanbara-gun, NIIGATA	7.35	1.07	1.58	1.6 ± 0.26	1.5 ± 0.24	4.2 ± 0.29	2.7 ± 0.18
Hachijo-Island, TOKYO	7.10	1.05	1.51	6.9 ± 0.40	6.6 ± 0.38	50.0 ± 0.90	33.0 ± 0.60
Katsuyama, FUKUI	7.20	1.10	1.64	3.2 ± 0.32	2.9 ± 0.29	3.7 ± 0.29	2.2 ± 0.18
Toyono-gun, OSAKA	7.35	1.10	1.53	1.2 ± 0.24	1.1 ± 0.22	0.6 ± 0.14	0.4 ± 0.09
Kochi, KOCHI	6.86	1.06	1.58	2.3 ± 0.27	2.2 ± 0.26	2.0 ± 0.24	1.2 ± 0.15
Kasuya-gun, FUKUOKA	6.91	1.06	1.41	1.0 ± 0.22	0.9 ± 0.21	1.9 ± 0.19	1.3 ± 0.14
Aira-gun, KAGOSHIMA	7.03	1.14	1.53	1.5 ± 0.26	1.3 ± 0.23	2.0 ± 0.23	1.3 ± 0.15
October, 1983							
Takada-gun, HIROSHIMA	6.64	1.06	1.36	1.3 ± 0.22	1.2 ± 0.20	0.8 ± 0.18	0.6 ± 0.13
November, 1983							
Sapporo, HOKKAIDO	7.62	1.27	1.60	2.1 ± 0.25	1.7 ± 0.20	3.7 ± 0.30	2.3 ± 0.19
Hachijo-Island, TOKYO	7.21	1.17	1.29	13.0 ± 0.50	11.0 ± 0.40	91.0 ± 1.20	71.0 ± 0.90
Nishikanbara-gun, NIIGATA	7.91	1.21	1.52	1.5 ± 0.24	1.2 ± 0.20	1.4 ± 0.24	0.9 ± 0.16
Katsuyama, FUKUI	7.72	1.19	1.62	3.3 ± 0.32	2.8 ± 0.27	2.6 ± 0.27	1.6 ± 0.17
Toyono-gun, OSAKA	7.43	1.16	1.52	1.4 ± 0.22	1.2 ± 0.19	0.9 ± 0.20	0.6 ± 0.13
Yatsuka-gun, SHIMANE	7.70	1.41	1.43	2.5 ± 0.28	1.7 ± 0.20	2.6 ± 0.27	1.8 ± 0.19
Kochi, KOCHI	7.37	1.13	1.59	2.5 ± 0.27	2.2 ± 0.24	1.4 ± 0.22	0.9 ± 0.14
Aira-gun, KAGOSHIMA	7.43	1.18	1.54	1.6 ± 0.25	1.3 ± 0.21	3.3 ± 0.27	2.2 ± 0.18
December, 1983							
Takada-gun, HIROSHIMA	6.12	0.988	1.28	0.7 ± 0.18	0.7 ± 0.18	0.8 ± 0.17	0.6 ± 0.13

Figure (3)-1 Sampling Locations of Milk

- 1. Sapporo
- 2. Hachijo-Island
- 3. Nishikanbara-gun
- 4. Katsuyama
- 5. Toyono-gun
- 6. Yatsuka-gun
- 7. Takada-gun
- 8. Kochi
- 9. Kasuya-gun
- 10. Aira-gun



**(3)-2 Strontium-90 and Cesium-137 in Milk (producing districts for domestic program)
(from Jun. 1983 to Sept. 1983)**


— continued from No. 65 of this publication —

Table (3)-2: Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (g/l)	Ca (g/l)	K (g/l)	pCi/l	S.U.	pCi/l	C.U.
June, 1983							
Saga-gun, SAGA	7.19	1.12	1.51	0.9 ± 0.24	0.8 ± 0.21	1.1 ± 0.22	0.8 ± 0.14
August, 1983							
Mito, IBARAGI	7.17	1.15	1.51	1.7 ± 0.26	1.5 ± 0.23	0.7 ± 0.15	0.5 ± 0.10
Hakui-gun, ISHIKAWA	7.17	1.17	1.51	1.1 ± 0.23	0.9 ± 0.20	1.0 ± 0.21	0.6 ± 0.14
Mihara-gun, HYOGO	7.00	1.11	1.59	1.1 ± 0.23	1.0 ± 0.21	3.1 ± 0.26	2.0 ± 0.16
Matsuyama, EHIME	7.27	1.12	1.49	1.2 ± 0.24	1.1 ± 0.22	1.1 ± 0.17	0.7 ± 0.11
September, 1983							
Aomori, AOMORI	7.45	1.08	1.62	20.0 ± 0.70	19.0 ± 0.60	7.8 ± 0.38	4.8 ± 0.23

Figure (3)-2 Sampling Locations of Milk

1. Aomori
2. Mito
3. Hakui-gun
4. Mihara-gun
5. Matsuyama
6. Saga-gun



(3)-3 Strontium-90 and Cesium-137 in Milk (consuming districts)
(from May, 1983 to Dec. 1983)


— continued from No. 65 of this publication —

Table (3)-3: Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (g/l)	Ca (g/l)	K (g/l)	pCi/l	S.U.	pCi/l	C.U.
May, 1983							
Sendai, MIYAGI	7.39	1.14	1.63	0.6 ± 0.19	0.5 ± 0.17	1.3 ± 0.21	0.8 ± 0.13
Kyoto, KYOTO	6.77	1.04	1.47	1.6 ± 0.25	1.6 ± 0.24	1.4 ± 0.21	1.0 ± 0.14
July, 1983							
Katsuren, OKINAWA	7.07	1.08	1.52	0.9 ± 0.24	0.9 ± 0.22	0.7 ± 0.15	0.5 ± 0.14
August, 1983							
Sapporo, HOKKAIDO	7.20	1.16	1.47	2.6 ± 0.28	2.2 ± 0.24	3.9 ± 0.29	2.6 ± 0.20
Akita, AKITA	7.17	1.00	1.57	5.2 ± 0.37	5.2 ± 0.37	4.3 ± 0.30	2.7 ± 0.19
Yamagata, YAMAGATA	6.88	1.07	1.52	0.9 ± 0.21	0.9 ± 0.20	1.1 ± 0.20	0.8 ± 0.13
Shinjuku, TOKYO	6.71	1.05	1.43	1.0 ± 0.23	0.9 ± 0.22	0.9 ± 0.19	0.7 ± 0.13
Yokohama, KANAGAWA	6.77	1.10	1.45	0.8 ± 0.23	0.8 ± 0.21	1.4 ± 0.20	0.9 ± 0.14
Niigata, NIIGATA	7.66	1.13	1.62	1.2 ± 0.27	1.1 ± 0.24	1.9 ± 0.24	1.2 ± 0.15
Fukui, FUKUI	6.95	1.04	1.46	1.2 ± 0.21	1.1 ± 0.20	1.3 ± 0.22	0.9 ± 0.15
Nagano, NAGANO	6.55	0.993	1.51	1.1 ± 0.20	1.1 ± 0.20	1.2 ± 0.18	0.8 ± 0.12
Shizuoka, SHIZUOKA	7.09	1.07	1.51	2.0 ± 0.28	1.8 ± 0.26	3.3 ± 0.26	2.2 ± 0.18
Nagoya, AICHI	7.36	1.11	1.61	1.0 ± 0.21	0.9 ± 0.19	0.6 ± 0.18	0.4 ± 0.11
Osaka, OSAKA	7.20	1.11	1.55	1.3 ± 0.26	1.1 ± 0.24	1.5 ± 0.21	1.0 ± 0.14
Yonago, TOTTORI	6.92	1.05	1.50	1.4 ± 0.23	1.3 ± 0.22	4.0 ± 0.28	2.7 ± 0.18
Okayama, OKAYAMA	7.20	1.08	1.55	0.8 ± 0.20	0.7 ± 0.19	1.4 ± 0.21	0.9 ± 0.13
Hiroshima, HIROSHIMA	5.87	0.882	1.25	1.1 ± 0.19	1.2 ± 0.22	1.0 ± 0.16	0.8 ± 0.13
Yamaguchi, YAMAGUCHI	6.99	1.06	1.55	0.7 ± 0.24	0.7 ± 0.22	1.3 ± 0.20	0.8 ± 0.13
Matsuyama, EHIME	7.60	1.10	1.52	0.9 ± 0.26	0.8 ± 0.24	0.8 ± 0.20	0.5 ± 0.13
Kochi, KOCHI	7.15	1.14	1.52	3.0 ± 0.31	2.6 ± 0.27	6.3 ± 0.34	4.1 ± 0.22
Chikushino, FUKUOKA	7.16	1.09	1.57	1.6 ± 0.27	1.5 ± 0.25	2.1 ± 0.24	1.3 ± 0.15
Kagoshima, KAGOSHIMA	7.05	1.06	1.54	2.0 ± 0.26	1.9 ± 0.25	5.9 ± 0.32	3.8 ± 0.21

Figure (3)-3 Sampling Locations of Milk

- | | |
|--------------|----------------|
| 1. Sapporo | 12. Kyoto |
| 2. Akita | 13. Osaka |
| 3. Sendai | 14. Yonago |
| 4. Yamagata | 15. Okayama |
| 5. Shinjuku | 16. Hiroshima |
| 6. Yokohama | 17. Yamaguchi |
| 7. Niigata | 18. Matsuyama |
| 8. Fukui | 19. Kochi |
| 9. Nagano | 20. Chikushino |
| 10. Shizuoka | 21. Kagoshima |
| 11. Nagoya | 22. Katsuren |



(3)-4 Strontium-90 and Cesium-137 in Milk (powderd milk)

– continued from No. 65 of this publication –

Table (3)-4: Strontium-90 and Cesium-137 in Milk

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (%)	K (%)	pCi/Kg	S.U.	pCi/Kg	C.U.
November, 1983							
Yukijirushi	2.45	3.72	5.86	5.0 ± 0.43	1.3 ± 0.12	13.0 ± 0.60	2.2 ± 0.10
Wakodo	2.47	3.38	6.20	2.9 ± 0.39	0.8 ± 0.12	4.3 ± 0.34	0.7 ± 0.05
Meiji	2.42	3.80	5.20	3.8 ± 0.40	1.0 ± 0.11	11.0 ± 0.5	2.1 ± 0.10
Morinaga	2.41	3.45	5.66	4.2 ± 0.40	1.2 ± 0.12	12.0 ± 0.5	2.2 ± 0.10
*Meiji	8.11	12.7	17.0	35.0 ± 1.4	27 ± 0.11	160.0 ± 2.0	9.1 ± 0.14
*Morinaga	8.14	12.6	18.3	19.0 ± 1.0	1.5 ± 0.08	20.0 ± 0.9	1.1 ± 0.05

* Skim milk

**(4)-1 Strontium-90 and Cesium-137 in Vegetables (producing districts)
(from May, 1983 to Dec. 1983)**

— continued from No. 65 of this publication —


Table (4)-1: Strontium-90 and Cesium-137 in Vegetables

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
(Japanese radish)							
May, 1983							
Atsumi-gun, AICHI	0.837	0.279	3.67	4.1 ± 0.38	15.0 ± 1.40	0.10 ± 0.14	0.04 ± 0.04
July, 1983							
Oota, SHIMANE	0.472	0.210	1.87	14.0 ± 0.70	66.0 ± 3.40	9.90 ± 0.51	5.30 ± 0.27
September, 1983							
Ishikari, HOKKAIDO	0.459	0.217	1.68	15.0 ± 0.70	70.0 ± 3.10	1.00 ± 0.19	0.60 ± 0.11
October, 1983							
Saku, NAGANO	0.638	0.285	2.77	1.6 ± 0.23	5.6 ± 0.79	0.40 ± 0.12	0.10 ± 0.04
November, 1983							
Sakai-gun, FUKUI	0.506	0.207	2.17	5.2 ± 0.33	25.0 ± 1.60	0.10 ± 0.11	0.10 ± 0.05
Gotenba, SHIZUOKA	0.592	0.170	1.99	4.4 ± 0.38	26.0 ± 2.20	1.30 ± 0.19	0.60 ± 0.10
Ibusuki-gun, KAGOSHIMA	0.558	0.169	2.44	2.8 ± 0.29	16.0 ± 1.70	1.10 ± 0.16	0.50 ± 0.07
December, 1983							
Akashi, HYOGO	0.561	0.241	2.23	2.4 ± 0.25	10.0 ± 1.00	0.20 ± 0.09	0.10 ± 0.04
Wakayama, WAKAYAMA	0.396	0.172	1.41	1.8 ± 0.33	11.0 ± 1.90	0.10 ± 0.19	0.10 ± 0.13
(Spinach)							
May, 1983							
Atsumi-gun, AICHI	0.741	0.270	2.38	1.1 ± 0.26	4.2 ± 0.96	0.02 ± 0.11	0.01 ± 0.05
July, 1983							
Oota, SHIMANE	1.950	0.730	8.40	15.0 ± 0.70	21.0 ± 1.00	2.70 ± 0.32	0.30 ± 0.04
September, 1983							
Ishikari, HOKKAIDO	1.460	0.475	6.14	14.0 ± 0.70	30.0 ± 1.50	1.10 ± 0.22	0.20 ± 0.04
October, 1983							
Saku, NAGANO	2.090	1.280	8.29	7.1 ± 0.55	5.6 ± 0.43	2.40 ± 0.29	0.30 ± 0.04
November, 1983							
Gotenba, SHIZUOKA	1.510	1.000	4.62	7.2 ± 0.47	7.2 ± 0.47	10.00 ± 0.50	2.20 ± 0.10

Location	Component			^{90}Sr		^{137}Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
Kurayoshi, TOTTORI	1.950	1.110	7.01	12.0 ± 0.70	10.0 ± 0.60	2.90 ± 0.33	0.40 ± 0.05
Matsuyama, EHIME	1.710	0.644	7.36	7.8 ± 0.51	12.0 ± 0.8	0.20 ± 0.14	0.03 ± 0.02
(Onion)							
July, 1983							
Sennan-gun, OSAKA	0.364	0.117	1.49	1.6 ± 0.27	13.0 ± 2.30	0.20 ± 0.12	0.10 ± 0.08

Figure (4)-1 Sampling Locations of Vegetables

- | | |
|---------------|-----------------|
| 1. Ishikari | 7. Akashi |
| 2. Sakai-gun | 8. Wakayama |
| 3. Saku | 9. Kurayoshi |
| 4. Gotenba | 10. Oota |
| 5. Atsumi-gun | 11. Matsuyama |
| 6. Sennan-gun | 12. Ibusuki-gun |



**(4)-2 Strontium-90 and Cesium-137 in Vegetables (consuming districts)
(from May, 1983 to Nov. 1983)**


— continued from No. 65 of this publication —

Table (4)-2: Strontium-90 and Cesium-137 in Vegetables

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C. U.
(Japanese radish)							
September, 1983							
Sendai, MIYAGI	0.595	0.256	2.53	34.0 ± 0.80	130.0 ± 3.00	1.00 ± 0.16	0.40 ± 0.06
October, 1983							
Akita, AKITA	0.481	0.282	1.82	14.0 ± 0.70	50.0 ± 2.3	0.30 ± 0.16	0.10 ± 0.09
Yamagata, YAMAGATA	0.472	0.317	1.61	2.2 ± 0.35	7.0 ± 1.1	0.40 ± 0.15	0.20 ± 0.10
Kyoto, KYOTO	0.497	0.260	1.81	12.0 ± 0.60	45.0 ± 2.4	2.30 ± 0.26	1.30 ± 0.14
November, 1983							
Shinjuku, TOKYO	0.530	0.332	2.19	6.0 ± 0.32	18.0 ± 1.0	0.60 ± 0.11	0.30 ± 0.05
Niigata, NIIGATA	0.454	0.206	1.82	4.3 ± 0.32	21.0 ± 1.5	0.90 ± 0.13	0.50 ± 0.07
Kanazawa, ISHIKAWA	0.394	0.199	1.63	4.9 ± 0.41	24.0 ± 2.1	1.10 ± 0.19	0.70 ± 0.11
Osaka, OSAKA	0.362	0.178	1.39	2.1 ± 0.29	12.0 ± 1.6	1.50 ± 0.19	1.10 ± 0.14
Okayama, OKAYAMA	0.518	0.285	2.15	19.0 ± 0.60	67.0 ± 2.1	0.80 ± 0.13	0.40 ± 0.06
Saga, SAGA	0.585	0.263	2.57	4.1 ± 0.35	15.0 ± 1.3	0.00 ± 0.08	0.00 ± 0.03
(Spinach)							
May, 1983							
Sendai, MIYAGI	1.660	0.911	6.26	5.0 ± 0.47	5.5 ± 0.51	0.50 ± 0.20	0.10 ± 0.03
June, 1983							
Niigata, NIIGATA	1.520	0.998	5.85	6.5 ± 0.52	6.5 ± 0.52	1.00 ± 0.22	0.20 ± 0.04
October, 1983							
Yamagata, YAMAGATA	1.710	0.740	6.95	2.9 ± 0.35	4.0 ± 0.47	0.70 ± 0.18	0.10 ± 0.03
November, 1983							
Shinjuku, TOKYO	2.250	1.500	9.40	3.4 ± 0.40	2.3 ± 0.26	5.60 ± 0.38	0.60 ± 0.04
Kanazawa, ISHIKAWA	1.360	0.561	5.84	1.2 ± 0.32	2.2 ± 0.56	0.04 ± 0.14	0.01 ± 0.02
Kyoto, KYOTO	1.540	0.394	6.96	1.0 ± 0.30	2.6 ± 0.77	0.40 ± 0.14	0.10 ± 0.02
Okayama, OKAYAMA	1.260	0.594	5.34	1.4 ± 0.33	2.4 ± 0.56	2.20 ± 0.26	0.40 ± 0.05
Matsuyama, EHIME	2.150	0.857	9.29	4.3 ± 0.40	5.0 ± 0.47	0.80 ± 0.18	0.10 ± 0.02
Saga, SAGA	1.93	1.250	7.76	1.2 ± 0.31	0.9 ± 0.25	0.50 ± 0.18	0.10 ± 0.02
(Cabbage)							
October, 1983							
Akita, AKITA	0.577	0.441	2.27	9.5 ± 0.40	22.0 ± 0.90	0.40 ± 0.12	0.20 ± 0.05

Figure (4)-2 Sampling Locations of Vegetables

1. Akita
2. Sendai
3. Yamagata
4. Shinjuku
5. Niigata
6. Kanazawa
7. Kyoto
8. Osaka
9. Okayama
10. Matsuyama
11. Saga



(5) Strontium-90 and Cesium-137 in Tea (Japanese tea)


— continued from No. 63 of this publication —

Table (5): Strontium-90 and Cesium-137 in Tea

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
June, 1983							
Iwata, SHIZUOKA	5.22	3.58	17.3	32 ± 2.7	9.0 ± 0.75	15.0 ± 1.7	0.8 ± 0.10
Tagata-gun, SHIZUOKA	5.10	4.09	18.0	250 ± 6.0	60.0 ± 1.5	88.0 ± 3.2	4.9 ± 0.18
Uji, KYOTO	5.06	3.02	18.4	38 ± 2.8	13.0 ± 0.9	5.0 ± 1.4	0.3 ± 0.07
Yosa-gun, KYOTO	5.94	3.67	21.2	58 ± 3.6	16.0 ± 1.0	30.0 ± 2.3	1.4 ± 0.11
Satsuma-gun, KAGOSHIMA	5.31	2.24	20.7	17 ± 2.4	8.0 ± 1.1	61.0 ± 3.0	2.9 ± 0.14
Kawabe-gun, KAGOSHIMA	5.23	2.91	20.0	59 ± 3.8	20.0 ± 1.3	120.0 ± 4.0	6.2 ± 0.21

Figure (5) Sampling Locations of Tea

1. Iwata
2. Tagata-gun
3. Uji
4. Yosa-gun
5. Satsuma-gun
6. Kawabe-gun



(6) Strontium-90 and Cesium-137 in Sea fish
(from Aug. 1983 to Dec. 1983)

— continued from No. 65 of this publication —

Table (6) Strontium-90 and Cesium-137 in Sea fish

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	(g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
(Hexagrammos otakii)							
August, 1983							
Souma, FUKUSHIMA	1.55	1.83	3.65	0.7 ± 0.28	0.4 ± 0.15	5.3 ± 0.42	1.5 ± 0.12
(Trachurus trachurus)							
July, 1983							
Harimanada, HYOGO	2.48	5.73	3.62	0.6 ± 0.25	0.1 ± 0.04	7.9 ± 0.39	2.2 ± 0.11
September, 1983							
Wakayama, WAKAYAMA	3.42	8.48	3.59	0.9 ± 0.29	0.1 ± 0.03	4.6 ± 0.35	1.3 ± 0.10
November, 1983							
Shizuoka, SHIZUOKA	3.64	9.19	3.25	1.7 ± 0.37	0.2 ± 0.04	5.2 ± 0.38	1.6 ± 0.12
December, 1983							
Odawara, KANAGAWA	3.55	7.16	2.70	0.4 ± 0.26	0.1 ± 0.04	3.7 ± 0.36	1.4 ± 0.13
(Sardinops melanosticta)							
September, 1983							
Yamagata, YAMAGATA	2.73	6.74	2.54	0.0 ± 0.22	0.0 ± 0.03	2.8 ± 0.28	1.1 ± 0.11
December, 1983							
Nagano, NAGANO	2.77	7.31	2.65	0.3 ± 0.32	0.05 ± 0.04	3.7 ± 0.37	1.4 ± 0.14
(Sebastiscus marmoratus)							
October, 1983							
Hamada, SHIMANE	5.34	14.6	3.06	1.4 ± 0.34	0.1 ± 0.02	4.7 ± 0.39	1.5 ± 0.13
(Limanda herzensteini)							
June, 1983							
Sendai, MIYAGI	2.76	7.33	2.30	0.1 ± 0.24	0.01 ± 0.03	2.4 ± 0.28	1.0 ± 0.12
November, 1983							
Mutsu, AOMORI	1.51	1.06	3.85	0.2 ± 0.25	0.20 ± 0.24	4.8 ± 0.40	1.2 ± 0.10
Niigata, NIIGATA	2.88	6.69	3.59	0.3 ± 0.27	0.04 ± 0.04	2.6 ± 0.32	0.7 ± 0.09
Fukui, FUKUI	1.53	2.08	3.26	0.7 ± 0.29	0.30 ± 0.14	5.5 ± 0.40	1.7 ± 0.12

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	(g/kg)	K (g/kg)	pCi/kg	S. U.	pCi/kg	C. U.
(<i>Spratelloides gracilis</i>) December, 1983 Akune, KAGOSHIMA	2.95	6.47	3.71	0.0 ± 0.28	0.0 ± 0.04	7.8 ± 0.46	2.1 ± 0.12
(<i>Oncorhynchus keta</i>) November, 1983 Urakawa, HOKKAIDO	1.47	0.467	4.18	0.1 ± 0.28	0.3 ± 0.60	6.8 ± 0.46	1.6 ± 0.11
(<i>Scomber japonicus</i>) September, 1983 Matsuyama, EHIME	1.29	0.357	3.09	0.3 ± 0.26	0.9 ± 0.74	6.3 ± 0.37	2.0 ± 0.12
November, 1983 Kyoto, KYOTO	1.90	3.08	2.81	0.5 ± 0.27	0.2 ± 0.09	3.9 ± 0.36	1.4 ± 0.13
(<i>Chrysophrys major</i>) July, 1983 Fukuoka, FUKUOKA	1.39	0.413	4.51	0.0 ± 0.13	0.0 ± 0.32	7.4 ± 0.28	1.6 ± 0.06
(<i>Arctoscopus japonicus</i>) December, 1983 Oga, AKITA	2.30	5.01	2.85	0.1 ± 0.26	0.02 ± 0.05	3.8 ± 0.35	1.3 ± 0.12
(<i>Seriola quinqueradiata</i>) October, 1983 Hakui-gun, ISHIKAWA	1.42	1.02	4.03	0.3 ± 0.26	0.3 ± 0.26	9.0 ± 0.46	2.2 ± 0.11
(<i>Mugil cephalus</i>) August, 1983 Ariake, SAGA	1.27	0.389	4.11	0.0 ± 0.23	0.0 ± 0.60	4.4 ± 0.31	1.1 ± 0.08
November, 1983 Oku-gun, OKAYAMA	1.35	0.516	4.11	1.9 ± 0.34	3.7 ± 0.65	5.0 ± 0.35	1.2 ± 0.09
(<i>Decapterus muroadsi</i>) September, 1983 Miyake-Island, TOKYO	1.32	0.915	3.30	0.2 ± 0.24	0.2 ± 0.27	5.6 ± 0.36	1.7 ± 0.11


Sea fish

Japanese name	English name	Scientific name
Ainame	Greenling	<i>Hexagrammos otakii</i>
Aji	Horse mackerel	<i>Trachurus trachurus</i>
Iwashi	Sardine	<i>Sardinops melanosticta</i>
Kasago	Scorpion-fish	<i>Sebastiscus marmoratus</i>
Karei	Flatfish	<i>Limanda herzensteini</i>
Kibinago	Banded blue-sprat	<i>Spratelloides gracilis</i>
Sake	Salmon	<i>Oncorhynchus keta</i>
Saba	Common mackerel	<i>Scomber japonicus</i>
Tai	Sea bream	<i>Chrysophrys major</i>
Hatahata	Japanese sandfish	<i>Arctoscopus japonicus</i>
Hukuragi (Buri)	Yellow-tail	<i>Seriola quinqueradiata</i>
Bora	Gray mullet	<i>Mugil cephalus</i>
Muroaji	Horse-scad mackerel	<i>Decapterus muroadsi</i>

Figure (6) Sampling Locations of Sea fish

1. Urakawa
2. Mutsu
3. Oga
4. Sendai
5. Yamagata
6. Souma
7. Miyake-Island
8. Odawara
9. Niigata
10. Hakui-gun
11. Fukui

12. Shizuoka
13. Nagano
14. Kyoto
15. Harimanada
16. Wakayama
17. Oku-gun
18. Hamada
19. Matsuyama
20. Fukuoka
21. Ariake
22. Akune



(7) Strontium-90 and Cesium-137 in Freshwater fish

— continued from No. 65 of this publication —


Table (7): Strontium-90 and Cesium-137 in Freshwater fish

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
(Carassius auratus)							
July, 1983							
Barato-lake, HOKKAIDO	4.41	13.40	2.88	45.0 ± 1.10	3.4 ± 0.08	3.4 ± 0.31	1.2 ± 0.11
November, 1983							
Toyanogata, NIIGATA	1.25	1.02	3.75	5.5 ± 0.66	5.4 ± 0.64	6.1 ± 0.51	1.6 ± 0.14
December, 1983							
Mikata-lake, FUKUI	0.95	0.503	2.94	3.0 ± 0.63	6.0 ± 1.30	8.5 ± 0.64	2.9 ± 0.22
Uji, KYOTO	5.19	13.30	3.24	49.0 ± 1.30	3.7 ± 0.10	3.8 ± 0.34	1.2 ± 0.10
(Cyprinus carpio)							
September, 1983							
Fukushima, FUKUSHIMA	3.23	8.94	3.07	31.0 ± 1.00	3.5 ± 0.11	3.9 ± 0.37	1.3 ± 0.12
November, 1983							
Hiroshima, HIROSHIMA	1.13	0.706	3.52	3.0 ± 0.48	4.3 ± 0.68	7.7 ± 0.50	2.2 ± 0.14
(Hypomesus transpacificus nipponensis)							
December, 1983							
Suwa-lake, NAGANO	2.12	4.88	2.44	3.8 ± 0.40	0.8 ± 0.08	3.1 ± 0.31	1.3 ± 0.13

Japanese name	English name	Scientific name
Funa	A crucian carp	Carassius auratus
Koi	Carp	Cyprinus carpio
Wakasagi	Pond-smelt	Hypomesus transpacificus nipponensis

Figure (7) Sampling Locations of Freshwater fish

1. Barato-lake
2. Fukushima
3. Toyanogata
4. Mikata-gun
5. Suwa-lake
6. Uji
7. Hiroshima



(8) Strontium-90 and Cesium-137 in Shellfish
(from Jun. 1983 to Nov. 1983)

— continued from No. 65 of this publication —


Table (8): Strontium-90 and Cesium-137 in Shellfish

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
<i>(Venerupis philippinarum)</i>							
June, 1983							
Kitatakagi-gun, NAGASAKI	2.00	2.29	1.83	0.04 ± 0.37	0.02 ± 0.16	1.1 ± 0.27	0.6 ± 0.15
<i>(Turbo cornutus)</i>							
July, 1983							
Sakata, YAMAGATA	2.64	2.33	2.52	0.6 ± 0.28	0.3 ± 0.12	1.5 ± 0.24	0.6 ± 0.09
August, 1983							
Hakui-gun, ISHIKAWA	2.92	2.10	2.26	0.5 ± 0.31	0.2 ± 0.15	1.2 ± 0.27	0.5 ± 0.12
<i>(Pecten yessoensis)</i>							
November, 1983							
Mutsu, AOMORI	2.07	0.564	3.15	0.0 ± 0.31	0.0 ± 0.55	1.6 ± 0.31	0.5 ± 0.10
<i>(Mytilus edulis)</i>							
July, 1983							
Mutsu, AOMORI	2.08	1.10	2.61	0.02 ± 0.28	0.01 ± 0.26	1.0 ± 0.22	0.4 ± 0.09

Japanese name	English name	Scientific name
Asari	Short-necked clam	<i>Venerupis philippinarum</i>
Sazae	Wreath shell	<i>Turbo cornutus</i>
Hotategai	Scallop	<i>Pecten yessoensis</i>
Murasakiigai	Mussel	<i>Mytilus edulis</i>

Figure (8) Sampling Locations of Shellfish

- 1. Mutsu
- 2. Sakata
- 3. Hakui-gun
- 4. Kitatakagi-gun



(9) Strontium-90 and Cesium-137 in Seaweeds
(from May, 1983 to Jun. 1983)

— continued from No. 65 of this publication —


Table (9): Strontium-90 and Cesium-137 in Seaweeds

Location	Component			⁹⁰ Sr		¹³⁷ Cs	
	Ash (%)	Ca (g/kg)	K (g/kg)	pCi/kg	S.U.	pCi/kg	C.U.
(Sargassum fulvellum)							
May, 1983							
Mutsu, AOMORI	2.10	1.13	5.75	2.1 ± 0.28	1.8 ± 0.25	0.6 ± 0.17	0.1 ± 0.03
June, 1983							
Fukaura, AOMORI	2.74	1.83	6.57	3.2 ± 0.50	1.7 ± 0.27	2.0 ± 0.27	0.3 ± 0.04

Japanese name	English name	Scientific name
Hondawara	Gulfweed	Sargassum fulvellum

Figure (9) Sampling Locations of Seaweeds

- 1. Mutsu
- 2. Fukaura



Edited by National Institute of Radiological Sciences, under the supervision of Science and Technology Agency of Japanese Government.