

ISSN 0441-2516

NIRS-RSD-97

# RADIOACTIVITY SURVEY DATA in Japan

Part 2

= Dietary Materials =

NUMBER 97

March 1992

National Institute of Radiological Sciences  
Chiba, Japan

# Radioactivity Survey Data in Japan

Number 97

March 1992 part 2 = Dietary Materials =

---

Contents	Page
Environmental and Dietary Materials (Japan Chemical Analysis Center)	
1. Collection and pretreatment of samples .....	1
2. Preparation of samples for analysis .....	3
3. Separation of Strontium-90 and Cesium-137 .....	3
4. Determination of Stable Strontium, Calcium and Potassium .....	4
5. Counting .....	4
6. Results .....	5
(1) Strontium-90 and Cesium-137 in Total Diet .....	5
(2)-1 Strontium-90 and Cesium-137 in Rice .....	8
(producing districts)	
-2 Strontium-90 and Cesium-137 in Rice .....	9
(consuming districts)	
(3)-1 Strontium-90 and Cesium-137 in Milk .....	10
(producing districts for domestic program)	
-2 Strontium-90 and Cesium-137 in Milk .....	11
(producing districts for WHO program)	
-3 Strontium-90 and Cesium-137 in Milk .....	12
(consuming districts)	
-4 Strontium-90 and Cesium-137 in Milk .....	13
(powderd milk)	
(4)-1 Strontium-90 and Cesium-137 in Vegetables .....	14
(producing districts)	
-2 Strontium-90 and Cesium-137 in Vegetables .....	17
(consuming districts)	
(5) Strontium-90 and Cesium-137 in Tea(Japanese Tea) .....	19
(6) Strontium-90 and Cesium-137 in Sea Fish .....	20
(7) Strontium-90 and Cesium-137 in Freshwater Fish .....	24
(8) Strontium-90 and Cesium-137 in Shellfish .....	25
(9) Strontium-90 and Cesium-137 in Seaweeds .....	27
7. Contents of Figure ( Selected Location ) .....	28

---

## Environmental and Dietary Materials\*

(Japan Chemical Analysis Center)

### 1. Collection and pretreatment of samples

#### (1) Rain and dry fallout

Rain and dry fallout was collected monthly on a sampling tray, approximately 5000 cm<sup>2</sup> in area, which was filled with water to a depth of 1 cm at the beginning of every month.

Strontium and cesium carrier solutions were added after the sample was filtered. The tray was washed with 5 l of distilled water and the washing was combined to the filtrate.

The sample was passed through a cation exchange column (500 ml of Dowex 50W X8, 50~100 mesh, Na form) at a rate flow of 80 ml/min.

#### (2) Airborne dust

Airborne dust was collected by an electrostatic precipitator or a filter air sampler for every three months at a rate of more than 3000 m<sup>3</sup> per month. The sampling was done 1 to 1.5 meters above the ground.

#### (3) Service water and freshwater

Service water, 100 l each, was collected at the intake of the water-treatment plant and at the tap after water was left running for five minutes. Strontium and cesium carriers were added to the filtered water sample. The subsequent process was the same as that described in the section (1). Freshwater was treated in the same way as the service water.

#### (4) Soil

Soil was collected from the location in the spacious and flat area without past surface disturbance caused by dust storms, inflow and outflow due to precipitation, etc.. Any places located under trees in a forest, in a stony area or inside of river banks were avoided. Soil was taken from two layers of different depths, 0-5 cm and 5-20 cm. The soil lumps were crushed by hands and dried in a drying oven regulated at 105°C. The soil was then passed through a 2 mm sieve to remove plant roots and pebbles.

#### (5) Sea water

Sea water was collected at the fixed stations where the effect of terrestrial fresh water from rivers was expected to be negligibly

small. A special consideration was also given to weather conditions. The sampling was carried out when there was no rainfall for the last few days. To prevent contamination, water samples were collected at the bow of a sampling boat just before she stood still by scooping surface water using a polyethylene bucket.

Immediately after the collection, the samples were acidified to a pH lower than 3 by adding concentrated hydrochloric acid in a ratio of 1 ml to 1 l of sea water, and then stored in 20 l polyethylene containers. The sampling equipments as well as containers were thoroughly rinsed with dilute hydrochloric acid and then with distilled water before use. Two hundred milliliters of sea water was also collected at the same stations for the determination of chlorinity.

#### (6) Sea sediments

Sediment was collected in the same area as that for the sea water sample, taking the following criteria into account:

- The depth of water exceeds 1 m at low tide.
- No significant sedimental movement is observed in the vicinity of concern.
- Mud, silt and fine sand are preferable.

A conventional sediment sampling device was used for collecting the top few centimeters of surface sediment. Approximately 4 kg of the sample in wet weight was spread on a stainless steel dish after removal of the pebbles, shells and other foreign materials, and dried in a drying oven regulated at 105°C.

#### (7) Total diet

A full one day ordinary diet including three meals, water, tea and other in-between snacks for five persons was collected as a sample of "total diet".

The sample in a large stainless steel pan was carbonized carefully by direct application of gas flame, and was transferred to a porcelain dish and then ashed at 450°C in an electric muffle furnace.

#### (8) Rice

Polished rice was collected in producing districts at the harvest and in consuming areas when new crops were first put on sale. The sample was carbonized and ashed in a porcelain dish.

---

\* Samples were sent to the Center from 46 contracted prefectures.

(9) Milk

Raw milk was collected in producing districts and commercial milk was purchased in consuming districts. Milk in a stainless steel pan or a porcelain dish was evaporated to dryness followed by carbonization and ashing.

(10) Vegetables

Spinach and Japanese radish were selected as the representatives for leaf vegetables and for non-starch roots, respectively. After removing soil, the edible part of vegetable sample was dried and carbonized in a stainless steel pan or a porcelain dish.

(11) Tea

Five hundred grams of manufactured green tea was collected, carbonized and ashed in a stainless steel pan or a porcelain dish.

(12) Fish, shellfish and seaweeds

a. Sea fish and freshwater fish

Fish was rinsed with water and blotted with a filter paper. Only the edible part was used in case of larger sized fish, and the whole part was used in case of smaller ones. Each sample was weighed and placed in a stainless steel pan or a porcelain dish. After carbonized, the sample was ashed in an electric muffle furnace.

b. Shellfish

Approximately 4 kg of shellfish including the shells was collected or purchased. After removing the shells, it was treated in the same way as that for the sea fish.

c. Seaweeds

Edible seaweeds were collected and rinsed with water to remove sand and other adhering matters on the surface. These were removed of excess water, weighed dried and ashed.

Table 1 shows details of sample collection.

Table 1 Details of sample collection

Sample	Frequency of sampling	Quantity of sample
=Environmental materials=		
(1) Rain and dry fallout		
1. For domestic program	monthly	
2. For WHO program	monthly	
(2) Airborne dust	quarterly	>3000 m <sup>3</sup> /month
(3) Service water and freshwater		
1. Service water (source water)	semiyearly	100 l
2. Service water (tap water)	semiyearly	100 l
3. Freshwater	yearly (fishing season)	100 l
(4) Soil		
1. 0~ 5 cm	yearly	4 kg
2. 5~ 20cm	yearly	4 kg
(5) Sea water	yearly	40 l
(6) Sea sediments	yearly	4 kg
=Dietary materials=		
(7) Total diet	semiyearly	daily amount for 5 persons
(8) Rice		
1. Producing districts	yearly (harvesting season)	5 kg (polished rice)
2. Consuming districts	yearly (harvesting season)	5 kg (polished rice)
(9) Milk		
1. Producing districts for WHO program	quarterly (February, May, August and November)	3 l
2. Producing districts for domestic program	semiyearly (February and August)	3 l

Sample	Frequency of sampling	Quantity of sample
3. Consuming districts	semiyearly (February and August)	3 l
4. Powdered milk	semiyearly (April and October)	2~ 3 kg
(10) Vegetables		
1. Producing districts	yearly (harvesting season)	4 kg
2. Consuming districts	yearly (harvesting season)	4 kg
(11) Tea	yearly (the first harvesting season)	500g (manufactured tea)
(12) Fish, shellfish and seaweeds		
1. Sea fish	yearly (fishing season)	4 kg
2. Freshwater fish	yearly (fishing season)	4 kg
3. Shellfish	yearly (fishing season)	4 kg
4. Seaweeds	yearly (fishing season)	2~ 3 kg

## 2. Preparation of samples for analysis

### (1) Rain, service water and freshwater

Strontium and cesium were eluted with hydrochloric acid from the cation exchange column. The residue of rain sample on the filter paper was ashed in an electric muffle furnace and the ash was dissolved in hydrochloric acid. The insoluble part was filtered and washed. The filtrate and the washings were combined to the previous eluate and used for radiochemical analysis.

### (2) Soil and Sea sediment

Dried soil was crushed to smaller ones than 0.25 mm in size by a crusher. The sieved sample was ashed in an electric muffle furnace regulated at 450°C. The sample was then heated with hydrochloric acid, strontium and cesium carrier solutions and the mixture was heated. The insoluble constituent was filtered off and washed with water.

The dried sample was crushed to smaller ones than 0.25 mm by a crushing machine. The further preparation of the sample was the same as that described in the section 2-(2).

### (3) Rice

The ashed sample was pulverized with a porcelain mortar and passed through a 0.35 mm sieve. The sieved sample to which both strontium and cesium carriers were added, was digested with nitric acid by heating. After the sample was heated again with nitric acid to dryness, strontium and cesium were extracted with hydrochloric acid and water. The insoluble constituent was filtered and washed. The filtrate and washings were combined for subsequent radiochemical analysis.

### (4) Airborne dust, diet, milk, vegetables, fish and shellfish, seaweeds, tea and others

These ashed samples were treated with the same procedure as that described in the section 2-(4).

## 3. Separation of strontium-90 and cesium-137

### (1) Strontium-90

Sample solutions, prepared as in the foregoing sections 2-(1) through 2-(4), were neutralized with sodium hydroxide. After sodium carbonate was added, the precipitate of strontium and calcium carbonates was separated. The supernatant solution was retained for cesium-137 determination. The carbonates were dissolved in hydrochloric acid and strontium and calcium were precipitated as oxalates. The precipitate was dissolved in nitric acid and strontium was separated from calcium by successive fuming nitric acid separation. Iron scavenge was made after addition of ferric iron carrier followed by barium chromate separation after addition of barium carrier to remove radium, its daughters and lead. Strontium was recovered as carbonate, and the precipitate was dried and weighed to determine strontium recovery. The strontium carbonate was dissolved in hydrochloric acid and iron carrier was added. The solution was allowed to stand for two weeks for strontium-90 and yttrium-90 to attain equilibrium. Yttrium-90 was coprecipitated with ferric hydroxide and the precipitate was filtered off, washed and counted.

### (2) Cesium-137

The supernatant separated from the strontium fraction was acidified with hydrochloric

acid. While stirring, cesium was adsorbed on the ammonium molybdophosphate added.

After filtered off and washed with hydrochloric acid the precipitate was dissolved in 2.5 N sodium hydroxide solution. The solution was adjusted to pH 8.2 with hydrochloric acid and allowed to cool. Resultant molybdenum hydroxide which separated out in the solution, was filtered off and washed with water. EDTA was added to the filtrate and washings. Cesium and rubidium were adsorbed on a cation exchange column and cesium was separated from rubidium by eluting with hydrochloric acid.

The eluate was evaporated to dryness and was dissolved. The solution was filtered. Chloroplatinic acid was added to precipitate cesium. The precipitate was filtered onto a tared paper using a demountable filter and washed with water and then ethanol. After drying, the chemical yield of cesium was determined by weighing the precipitate. Cesium-137 radioactivity was measured for this precipitate.

#### 4. Determination of stable strontium, calcium and potassium

A weighed amount of soil or sea sediment was heated in a electric muffle furnace at 450

°C and then treated with hydrochloric acid for extraction. A weighed aliquot of ashed samples of total diet, vegetables, milk, fish, shellfish or seaweeds was digested with hydrofluoric acid and nitric acid.

The extract was made up to an appropriate volume with dilute hydrochloric acid. The sample solution was analyzed for calcium by titration with standard potassium permanganate solution after separating calcium as oxalate. Atomic absorption spectroscopy was applied when appropriate. Stable strontium and potassium were determined by atomic absorption and flame emission spectrometry, respectively.

#### 5. Counting

After the radiochemical separation the mounted precipitates were counted for activity using low background beta counters normally for 60 to 90 min.

Net sample counting rates were corrected for counter efficiency, recovery, self-absorption and decay to obtain the content of strontium-90 and cesium-137 per sample aliquot. From the results, concentrations of these nuclides in the original samples were calculated.

## 6. Results

### (1) Strontium-90 and Cesium-137 in Total Diet (from Nov. 1990 to Mar. 1991)

-continued from No. 95 of this publication-

Table (1): Strontium-90 and Cesium-137 in Total Diet

Location	Ash	Ca	K	<sup>90</sup> Sr		<sup>137</sup> Cs	
	(g/p·d)	(mg/p·d)	(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)
November, 1990							
Ishinomaki, MIYAGI	13.9	540	1620	0.066±0.0092	0.12 ± 0.017	0.083±0.0083	0.051±0.0051
Onagawa-machi, MIYAGI	13.8	578	1610	0.073±0.0098	0.13 ± 0.017	0.075±0.0081	0.047±0.0050
Ichihara, CHIBA	18.1	346	1660	0.066±0.0066	0.19 ± 0.019	0.084±0.0095	0.050±0.0057
Chikura-machi, CHIBA	16.5	512	2230	0.081±0.0067	0.16 ± 0.013	0.090±0.0093	0.040±0.0042
Toyama, TOYAMA	12.2	353	1540	0.034±0.0040	0.097±0.011	0.042±0.0059	0.027±0.0038
Takaoka, TOYAMA	18.4	585	2150	0.084±0.0072	0.14 ± 0.012	0.071±0.0089	0.033±0.0041
Nagano, NAGANO	15.9	716	1830	0.057±0.012	0.079±0.017	0.034±0.0096	0.019±0.0052
Owase, MIE	12.6	412	1720	0.15 ± 0.008	0.36 ± 0.020	0.090±0.0085	0.052±0.0050
Kashihara, NARA	15.1	696	1330	0.046±0.0089	0.066±0.013	0.037±0.0074	0.028±0.0056
Gojou, NARA	11.3	650	1550	0.061±0.0087	0.094±0.013	0.031±0.0067	0.020±0.0043
Tottori, TOTTORI	15.1	511	1910	0.11 ± 0.011	0.21 ± 0.022	0.13 ± 0.010	0.067±0.0053
Fukube-mura, TOTTORI	14.5	601	1980	0.070±0.0058	0.12 ± 0.010	0.051±0.0073	0.026±0.0037
Matsue, SHIMANE	26.2	1090	3720	0.16 ± 0.018	0.14 ± 0.017	0.14 ± 0.015	0.037±0.0040
Kashima-machi, SHIMANE	21.4	1970	2190	0.16 ± 0.015	0.082±0.0077	0.045±0.0091	0.021±0.0041
Okayama, OKAYAMA	16.4	573	2440	0.065±0.013	0.11 ± 0.023	0.11 ± 0.013	0.047±0.0052
Kamisaibara-mura, OKAYAMA							
	13.9	382	1790	0.11 ± 0.008	0.28 ± 0.020	0.34 ± 0.016	0.19 ± 0.0090
Ikata-machi, EHIME	9.80	302	955	0.039±0.0037	0.13 ± 0.012	0.011±0.0045	0.012±0.0047
Fukuoka, FUKUOKA	10.4	266	1020	0.031±0.0057	0.12 ± 0.021	0.027±0.0045	0.026±0.0044
Dazaifu, FUKUOKA	14.2	554	2030	0.098±0.010	0.18 ± 0.018	0.13 ± 0.010	0.062±0.0048
Nagasaki, NAGASAKI	11.6	407	1560	0.049±0.0045	0.12 ± 0.011	0.035±0.0053	0.022±0.0034
Matsuura, NAGASAKI	11.7	498	1470	0.057±0.0047	0.11 ± 0.009	0.043±0.0055	0.029±0.0037
Ookuchi, KAGOSHIMA	13.3	385	1010	0.055±0.0051	0.14 ± 0.013	0.035±0.0069	0.035±0.0068
Sendai, KAGOSHIMA	13.5	450	1260	0.081±0.0062	0.18 ± 0.014	0.063±0.0080	0.050±0.0063
December, 1990							
Sapporo, HOKKAIDO	24.9	725	3180	0.12 ± 0.010	0.17 ± 0.014	0.18 ± 0.015	0.057±0.0046
Iwanai-machi, HOKKAIDO	17.3	410	1770	0.089±0.0075	0.22 ± 0.018	0.081±0.0086	0.046±0.0049
Aomori, AOMORI	15.9	534	2210	0.12 ± 0.009	0.23 ± 0.016	0.080±0.0090	0.036±0.0041
Ajigasawa-machi, AOMORI	15.6	756	2240	0.093±0.0078	0.12 ± 0.010	0.093±0.0093	0.042±0.0042
Morioka, IWATE	13.8	315	1820	0.069±0.010	0.22 ± 0.032	0.055±0.0076	0.030±0.0042
Fukushima, FUKUSHIMA	12.2	434	1570	0.039±0.010	0.089±0.023	0.069±0.010	0.044±0.0065
Mito, IBARAKI	17.7	599	2140	0.097±0.013	0.16 ± 0.022	0.061±0.0095	0.029±0.0045

Location	Ash	Ca	K	<sup>90</sup> Sr		<sup>137</sup> Cs	
	(g/p·d)	(mg/p·d)	(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)
Tokai-mura, IBARAKI	16.1	480	2150	0.092±0.012	0.19 ± 0.025	0.050±0.0082	0.023±0.0038
Shinjuku, TOKYO	13.5	317	1540	0.049±0.053	0.15 ± 0.017	0.033±0.0052	0.021±0.0034
Hachijo-machi, TOKYO	14.7	879	3020	0.065±0.0057	0.074±0.0064	0.077±0.0075	0.025±0.0022
Yokohama, KANAGAWA	12.7	336	1690	0.061±0.0052	0.18 ± 0.015	0.061±0.0071	0.036±0.0042
Hiratsuka, KANAGAWA	14.7	533	2280	0.078±0.0063	0.15 ± 0.012	0.077±0.0084	0.034±0.0037
Utsunomiya, TOCHIGI	12.6	438	1850	0.049±0.0048	0.11 ± 0.011	0.089±0.0080	0.048±0.0043
Kanuma, TOCHIGI	11.8	375	1480	0.063±0.0051	0.17 ± 0.014	0.16 ± 0.010	0.11 ± 0.007
Kashiwazaki, NIIGATA	18.9	581	2580	0.16 ± 0.016	0.28 ± 0.027	0.10 ± 0.011	0.039±0.0043
Nishikawa-machi, NIIGATA	22.6	756	3050	0.11 ± 0.016	0.14 ± 0.021	0.041±0.010	0.014±0.0034
Fukui, FUKUI	15.1	760	2010	0.073±0.0061	0.096±0.0080	0.050±0.0067	0.025±0.0033
Ueda, NAGANO	17.4	606	2420	0.073±0.013	0.12 ± 0.022	0.040±0.0092	0.017±0.0038
Koufu, YAMANASHI	13.3	429	1490	0.043±0.0047	0.099±0.011	0.023±0.0050	0.015±0.0034
Sudama-machi, YAMANASHI	11.9	374	1590	0.040±0.0072	0.11 ± 0.019	0.040±0.0060	0.025±0.0038
Gifu, GIFU	13.4	534	1610	0.055±0.010	0.10 ± 0.019	0.046±0.0079	0.028±0.0049
Tarui-machi, GIFU	9.73	258	1260	0.036±0.0069	0.14 ± 0.027	0.054±0.0059	0.043±0.0047
Tsu, MIE	17.1	345	1760	0.050±0.0061	0.14 ± 0.018	0.10 ± 0.011	0.059±0.0062
Ootsu, SHIGA	15.9	515	1690	0.067±0.012	0.13 ± 0.023	0.065±0.0086	0.038±0.0051
Imazu-machi, SHIGA	12.1	462	1540	0.071±0.0053	0.15 ± 0.012	0.047±0.0066	0.031±0.0043
Kyoto, KYOTO	17.2	836	2530	0.084±0.011	0.10 ± 0.013	0.063±0.0087	0.025±0.0035
Maizuru, KYOTO	14.6	593	1920	0.067±0.0088	0.11 ± 0.015	0.080±0.0084	0.042±0.0044
Osaka, OSAKA	15.4	560	2230	0.069±0.0095	0.12 ± 0.017	0.051±0.0072	0.023±0.0032
Neyagawa, OSAKA	15.2	629	1950	0.051±0.0094	0.082±0.015	0.069±0.0084	0.035±0.0043
Kakogawa, HYOGO	14.5	623	1690	0.066±0.0094	0.11 ± 0.015	0.034±0.0062	0.020±0.0037
Hamasaka-machi, HYOGO	11.3	476	1650	0.062±0.0084	0.13 ± 0.018	0.064±0.0068	0.039±0.0041
Miyoshi, HIROSHIMA	13.6	899	1770	0.081±0.0094	0.090±0.010	0.054±0.0075	0.031±0.0042
Tokushima, TOKUSHIMA	24.2	780	2790	0.074±0.0087	0.096±0.011	0.073±0.010	0.026±0.0036
Takamatsu, KAGAWA	14.9	496	1680	0.066±0.0073	0.13 ± 0.015	0.047±0.0078	0.028±0.0046
Tsuda-machi, KAGAWA	12.7	472	1310	0.041±0.0047	0.087±0.010	0.032±0.0054	0.024±0.0041
Kumamoto, KUMAMOTO	13.8	328	1570	0.052±0.0086	0.16 ± 0.026	0.057±0.0070	0.036±0.0044
Aso-machi, KUMAMOTO	16.9	631	1970	0.11 ± 0.012	0.17 ± 0.020	0.10 ± 0.010	0.052±0.0052
Saiki, OOITA	12.8	609	1810	0.056±0.0086	0.092±0.014	0.069±0.0076	0.038±0.0042
Miyazaki, MIYAZAKI	16.8	495	1810	0.073±0.011	0.15 ± 0.022	0.071±0.0093	0.039±0.0052
Takahara-machi, MIYAZAKI	17.4	608	2830	0.094±0.012	0.15 ± 0.020	0.18 ± 0.013	0.063±0.0047
January, 1991							
Maebashi, GUNMA	16.4	489	2150	0.050±0.011	0.10 ± 0.023	0.043±0.0090	0.020±0.0042
Nakanojo-machi, GUNMA	17.6	584	2270	0.10 ± 0.014	0.18 ± 0.023	0.12 ± 0.012	0.052±0.0052
Hiroshima, HIROSHIMA	13.6	584	1490	0.046±0.0086	0.079±0.015	0.029±0.0064	0.019±0.0043
Yamaguchi, YAMAGUCHI	17.0	646	2240	0.084±0.0071	0.13 ± 0.011	0.17 ± 0.012	0.077±0.0056
Ajisu-machi, YAMAGUCHI	24.3	1180	2710	0.066±0.0080	0.056±0.0068	0.071±0.011	0.026±0.0040



Location	Ash	Ca	K	$^{90}\text{Sr}$		$^{137}\text{Cs}$	
	(g/p·d)	(mg/p·d)	(mg/p·d)	(Bq/p·d)	(Bq/gCa)	(Bq/p·d)	(Bq/gK)
Ooita, OOITA	18.0	670	2280	$0.080 \pm 0.0073$	$0.12 \pm 0.011$	$0.16 \pm 0.013$	$0.072 \pm 0.0056$
Naha, OKINAWA	14.8	687	1820	$0.043 \pm 0.0052$	$0.062 \pm 0.0075$	$0.051 \pm 0.0071$	$0.028 \pm 0.0039$
Ginowan, OKINAWA	16.1	787	2290	$0.076 \pm 0.0064$	$0.097 \pm 0.0082$	$0.11 \pm 0.010$	$0.049 \pm 0.0044$
February, 1991							
Maebashi, GUNMA	16.5	564	2050	$0.086 \pm 0.013$	$0.15 \pm 0.022$	$0.032 \pm 0.0083$	$0.016 \pm 0.0041$
Nakanojo-machi, GUNMA	15.4	605	2310	$0.082 \pm 0.0066$	$0.14 \pm 0.011$	$0.076 \pm 0.0091$	$0.033 \pm 0.0039$
Hamaoka-machi, SHIZUOKA	13.7	636	2040	$0.056 \pm 0.0083$	$0.089 \pm 0.013$	$0.046 \pm 0.0067$	$0.023 \pm 0.0033$
Gifu, GIFU	12.0	484	1770	$0.036 \pm 0.0043$	$0.075 \pm 0.0089$	$0.055 \pm 0.0060$	$0.031 \pm 0.0034$
Tarui-machi, GIFU	9.03	280	1140	$0.044 \pm 0.0066$	$0.16 \pm 0.024$	$0.050 \pm 0.0055$	$0.044 \pm 0.0048$
Wakayama, WAKAYAMA	13.5	585	1420	$0.046 \pm 0.0050$	$0.079 \pm 0.0085$	$0.032 \pm 0.0067$	$0.023 \pm 0.0047$
Koza-machi, WAKAYAMA	12.3	464	1280	$0.068 \pm 0.0053$	$0.15 \pm 0.011$	$0.036 \pm 0.0058$	$0.029 \pm 0.0046$
Okayama, OKAYAMA	18.3	714	2280	$0.082 \pm 0.0087$	$0.12 \pm 0.012$	$0.054 \pm 0.010$	$0.024 \pm 0.0045$
Kamisaibara-mura, OKAYAMA	14.4	486	1690	$0.086 \pm 0.0077$	$0.18 \pm 0.016$	$0.078 \pm 0.0097$	$0.046 \pm 0.0057$
Kamiita-machi, TOKUSHIMA	15.9	373	1940	$0.068 \pm 0.010$	$0.18 \pm 0.028$	$0.075 \pm 0.0083$	$0.039 \pm 0.0043$
March, 1991							
Ookuma-machi, FUKUSHIMA	11.5	253	982	$0.030 \pm 0.0074$	$0.12 \pm 0.029$	$0.083 \pm 0.0079$	$0.085 \pm 0.0080$

(2)-1 Strontium-90 and Cesium-137 in Rice (producing districts)  
(from Aug. 1990 to Dec. 1990)

-continued from No. 95 of this publication-

Table (2)-1: Strontium-90 and Cesium-137 in Rice

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
August, 1990							
Sadowara-machi, MIYAZAKI	0.446	0.027	0.830	0.011 ± 0.0030	0.43 ± 0.11	0.0083 ± 0.0035	0.010 ± 0.0042
September, 1990							
Chiba, CHIBA	0.503	0.047	0.920	0.0085 ± 0.0035	0.18 ± 0.074	0.0084 ± 0.0051	0.0092 ± 0.0055
Toyoshina-machi, NAGANO	0.613	0.054	1.15	0.015 ± 0.0072	0.29 ± 0.14	0.0059 ± 0.0041	0.0051 ± 0.0036
Shiga-machi, SHIGA	0.729	0.037	0.984	0.0039 ± 0.0077	0.11 ± 0.21	0.016 ± 0.0079	0.016 ± 0.0080
Usa, OOLTA	0.721	0.045	0.822	0.0000 ± 0.0080	0.00 ± 0.18	0.0000 ± 0.0037	0.0000 ± 0.0045
October, 1990							
Mito, IBARAKI	0.616	0.035	0.612	0.0093 ± 0.0039	0.27 ± 0.11	0.027 ± 0.0060	0.044 ± 0.0097
Maki-machi, NIIGATA	0.404	0.023	0.671	0.0038 ± 0.0041	0.17 ± 0.18	0.0042 ± 0.0041	0.0062 ± 0.0061
Kosugi-machi, YOYAMA	0.429	0.036	0.652	0.012 ± 0.0029	0.33 ± 0.080	0.022 ± 0.0041	0.033 ± 0.0064
Gifu, GIFU	0.503	0.046	0.850	0.012 ± 0.0057	0.25 ± 0.13	0.0014 ± 0.0029	0.0017 ± 0.0034
Matsusaka, MIE	0.593	0.035	0.973	0.0098 ± 0.0036	0.28 ± 0.10	0.0056 ± 0.0045	0.0058 ± 0.0046
Kashihara, NARA	0.898	0.040	1.11	0.017 ± 0.0052	0.42 ± 0.13	0.014 ± 0.0089	0.012 ± 0.0080
Ishii-machi, TOKUSHIMA	0.657	0.055	0.670	0.025 ± 0.0082	0.46 ± 0.15	0.0033 ± 0.0061	0.0050 ± 0.0091
Tsuda-machi, KAGAWA	0.588	0.039	0.970	0.0064 ± 0.0027	0.16 ± 0.068	0.0050 ± 0.0036	0.0052 ± 0.0037
Koushi-machi, KUMAMOTO	0.466	0.021	0.610	0.0036 ± 0.0020	0.17 ± 0.091	0.0035 ± 0.0028	0.0058 ± 0.0046
November, 1990							
Ishikari-machi, HOKKAIDO	0.587	0.042	0.916	0.017 ± 0.0076	0.42 ± 0.18	0.0003 ± 0.0033	0.0004 ± 0.0036
Takisawa-mura, IWATE	0.625	0.045	0.750	0.0060 ± 0.0030	0.13 ± 0.067	0.40 ± 0.016	0.53 ± 0.021
Fukushima, FUKUSHIMA	0.778	0.034	1.24	0.0000 ± 0.0093	0.00 ± 0.28	0.013 ± 0.0081	0.010 ± 0.0065
Maebashi, GUNMA	0.617	0.031	0.673	0.0077 ± 0.0035	0.25 ± 0.11	0.0000 ± 0.0058	0.0000 ± 0.0087
Utsunomiya, TOCHIGI	0.636	0.033	0.763	0.0097 ± 0.0065	0.30 ± 0.20	0.031 ± 0.0077	0.040 ± 0.010
Kasai, HYOGO	0.465	0.026	0.544	0.0078 ± 0.0027	0.30 ± 0.10	0.0043 ± 0.0045	0.0078 ± 0.0083
Yamaguchi, YAMAGUCHI	0.656	0.033	0.991	0.0085 ± 0.0035	0.26 ± 0.11	0.043 ± 0.0079	0.043 ± 0.0080
December, 1990							
Nagasaka-machi, YAMANASHI	0.567	0.038	0.811	0.0043 ± 0.0060	0.11 ± 0.16	0.015 ± 0.0045	0.018 ± 0.0055
Chikushino, FUKUOKA	0.636	0.049	0.642	0.016 ± 0.0080	0.33 ± 0.16	0.024 ± 0.0051	0.038 ± 0.0079
Kagoshima, KAGOSHIMA	0.591	0.026	0.733	0.012 ± 0.0071	0.47 ± 0.27	0.093 ± 0.0092	0.13 ± 0.013

(2)-2 Strontium-90 and Cesium-137 in Rice(consuming districts)  
(from Sep. 1990 to Jan. 1991)

-continued from No. 95 of this publication-

Table (2)-2: Strontium-90 and Cesium-137 in Rice

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
September, 1990							
Kanazawa, ISHIKAWA	0.641	0.036	0.968	0.013 ± 0.0036	0.35 ± 0.099	0.044 ± 0.0069	0.045 ± 0.0072
Fukui, FUKUI	0.563	0.028	1.06	0.019 ± 0.0068	0.67 ± 0.24	0.0028 ± 0.0053	0.0027 ± 0.0050
October, 1990							
Mito, IBARAKI	0.548	0.035	0.712	0.0021 ± 0.0027	0.062 ± 0.077	0.0042 ± 0.0044	0.0058 ± 0.0061
Shinjuku, TOKYO	0.490	0.039	0.862	0.0040 ± 0.0024	0.10 ± 0.063	0.058 ± 0.0066	0.067 ± 0.0076
Niigata, NIIGATA	0.427	0.024	0.623	0.0051 ± 0.0044	0.21 ± 0.18	0.0037 ± 0.0042	0.0060 ± 0.0068
Hiroshima, HIROSHIMA	0.465	0.038	0.735	0.015 ± 0.0031	0.41 ± 0.083	0.040 ± 0.0051	0.054 ± 0.0069
Matsuyama, EHIME	0.487	0.039	0.760	0.0066 ± 0.0029	0.17 ± 0.075	0.0000 ± 0.0028	0.0000 ± 0.0037
Yonagusuku-mura, OKINAWA	0.547	0.037	0.968	0.0071 ± 0.0029	0.19 ± 0.080	0.016 ± 0.0045	0.016 ± 0.0046
November, 1990							
Sapporo, HOKKAIDO	0.599	0.047	0.952	0.015 ± 0.0068	0.33 ± 0.15	0.018 ± 0.0052	0.019 ± 0.0054
Tajiri-machi, MIYAGI	0.773	0.036	0.943	0.010 ± 0.0037	0.29 ± 0.11	0.017 ± 0.0054	0.018 ± 0.0057
Akita, AKITA	0.656	0.036	0.741	0.024 ± 0.0042	0.67 ± 0.12	0.049 ± 0.0064	0.066 ± 0.0086
Yamagata, YAMAGATA	0.567	0.031	0.669	0.014 ± 0.0032	0.47 ± 0.10	0.097 ± 0.0079	0.15 ± 0.012
Shizuoka, SHIZUOKA	0.424	0.025	0.729	0.010 ± 0.0058	0.40 ± 0.23	0.018 ± 0.0048	0.025 ± 0.0065
Kyoto, KYOTO	0.635	0.036	0.667	0.0062 ± 0.0072	0.17 ± 0.20	0.0071 ± 0.0064	0.011 ± 0.0097
Osaka, OSAKA	0.486	0.037	0.807	0.011 ± 0.0032	0.30 ± 0.087	0.060 ± 0.0066	0.074 ± 0.0082
Kobe, HYOGO	0.524	0.027	0.713	0.0067 ± 0.0028	0.25 ± 0.10	0.012 ± 0.0055	0.017 ± 0.0078
Saga, SAGA	0.510	0.036	0.780	0.0031 ± 0.0027	0.086 ± 0.075	0.0035 ± 0.0034	0.0044 ± 0.0044
December, 1990							
Yokohama, KANAGAWA	0.456	0.026	0.775	0.0048 ± 0.0047	0.19 ± 0.18	0.013 ± 0.0053	0.017 ± 0.0069
Nagoya, AICHI	0.710	0.039	0.684	0.0036 ± 0.0034	0.091 ± 0.087	0.012 ± 0.0074	0.017 ± 0.011
Tottori, TOTTORI	0.534	0.044	0.780	0.012 ± 0.0042	0.27 ± 0.096	0.050 ± 0.0070	0.064 ± 0.0089
Matsue, SHIMANE	0.747	0.041	0.769	0.0020 ± 0.0039	0.047 ± 0.093	0.075 ± 0.010	0.097 ± 0.013
Seto-machi, OKAYAMA	0.567	0.024	1.01	0.0029 ± 0.0027	0.12 ± 0.11	0.0027 ± 0.0055	0.0027 ± 0.0055
Kochi, KOCHI	0.532	0.027	0.819	0.0096 ± 0.0063	0.36 ± 0.23	0.0000 ± 0.0049	0.0000 ± 0.0060
Kasuga, FUKUOKA	0.658	0.045	1.05	0.0005 ± 0.0077	0.01 ± 0.17	0.020 ± 0.0054	0.019 ± 0.0051
January, 1991							
Hirosaki, AOMORI	0.624	0.046	0.936	0.0048 ± 0.0034	0.10 ± 0.073	0.018 ± 0.0065	0.019 ± 0.0070
Shinguu, WAKAYAMA	0.603	0.042	0.941	0.021 ± 0.0076	0.50 ± 0.18	0.017 ± 0.0064	0.018 ± 0.0068
Nagasaki, NAGASAKI	0.408	0.042	0.865	0.013 ± 0.0031	0.31 ± 0.074	0.045 ± 0.0062	0.051 ± 0.0072

(3)-1 Strontium-90 and Cesium-137 in Milk (producing districts for domestic program)  
(from Aug. 1990 to Mar. 1991)

-continued from No. 95 of this publication-

Table (3)-1: Strontium-90 and Cesium-137 in Milk

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	Bq/ℓ	Bq/gCa	Bq/ℓ	Bq/gK
August, 1990							
Aomori, AOMORI	7.27	1.04	1.57	0.13 ± 0.011	0.12 ± 0.010	0.15 ± 0.011	0.098 ± 0.0069
Oshimizu-machi, ISHIKAWA	7.03	1.12	1.55	0.032 ± 0.0075	0.028 ± 0.0067	0.053 ± 0.0073	0.034 ± 0.0047
December, 1990							
Kasamatsu-machi, Gifu	7.05	1.11	1.35	0.055 ± 0.0085	0.049 ± 0.0077	0.038 ± 0.0064	0.028 ± 0.0047
January, 1991							
Fujimi-mura, GUNMA	7.44	1.13	1.66	0.033 ± 0.0078	0.029 ± 0.0069	0.033 ± 0.0065	0.020 ± 0.0039
February, 1991							
Aomori, AOMORI	7.36	1.14	1.50	0.039 ± 0.0074	0.034 ± 0.0065	0.031 ± 0.0061	0.021 ± 0.0041
Takisawa-mura, IWATE	7.40	1.09	1.65	0.025 ± 0.0085	0.022 ± 0.0078	0.10 ± 0.011	0.061 ± 0.0064
Mito, IBARAKI	7.42	1.13	1.45	0.020 ± 0.0081	0.018 ± 0.0072	0.011 ± 0.0048	0.0076 ± 0.0033
Nishinasuno-machi, TOCHIGI	7.09	1.19	1.21	0.048 ± 0.0089	0.040 ± 0.0074	0.050 ± 0.0069	0.042 ± 0.0057
Tonami, TOYAMA	7.53	1.13	1.51	0.033 ± 0.0072	0.029 ± 0.0063	0.037 ± 0.0067	0.024 ± 0.0045
Oshimizu-machi, ISHIKAWA	7.21	1.16	1.55	0.028 ± 0.0073	0.024 ± 0.0063	0.23 ± 0.013	0.15 ± 0.008
Oouchiyama-mura, MIE	7.46	1.13	1.56	0.018 ± 0.0071	0.016 ± 0.0063	0.019 ± 0.0053	0.012 ± 0.0034
Hino-machi, SHIGA	7.47	1.15	1.55	0.018 ± 0.0067	0.016 ± 0.0058	0.0000 ± 0.0041	0.0000 ± 0.0027
Mihara-machi, HYOGO	7.47	1.13	1.67	0.028 ± 0.0075	0.025 ± 0.0067	0.0043 ± 0.0045	0.0026 ± 0.0027
Oouda-machi, NARA	6.40	1.04	1.20	0.027 ± 0.0042	0.026 ± 0.0040	0.0026 ± 0.0030	0.0022 ± 0.0025
Matsuyama, EHIME	7.77	1.19	1.47	0.0060 ± 0.0069	0.0050 ± 0.0058	0.0099 ± 0.0054	0.0067 ± 0.0036
Kamiita-machi, TOKUSHIMA	7.27	1.12	1.35	0.032 ± 0.0084	0.029 ± 0.0075	0.0007 ± 0.0045	0.0005 ± 0.0033
Takasa-machi, KAGAWA	7.71	1.16	1.56	0.025 ± 0.0042	0.021 ± 0.0037	0.0065 ± 0.0078	0.0042 ± 0.0050
Koushi-machi, KUMAMOTO	7.16	1.06	1.49	0.016 ± 0.0070	0.015 ± 0.0066	0.017 ± 0.0049	0.012 ± 0.0033
Kujuu-machi, OOTA	7.60	1.21	1.59	0.035 ± 0.0044	0.029 ± 0.0037	0.083 ± 0.0090	0.052 ± 0.0057
Takahara-machi, MIYAZAKI	7.60	1.11	1.64	0.019 ± 0.0039	0.017 ± 0.0035	0.080 ± 0.0077	0.049 ± 0.0047
March, 1991							
Yachimata-machi, CHIBA	7.55	1.14	1.65	0.039 ± 0.0050	0.034 ± 0.0044	0.069 ± 0.0070	0.042 ± 0.0042
Takane-machi, YAMANASHI	6.98	1.06	1.55	0.035 ± 0.0071	0.0033 ± 0.0067	0.0062 ± 0.0046	0.0040 ± 0.0030

(3)-2 Strontium-90 and Cesium-137 in Milk(producing districts for WHO program)  
(from May. 1990 to Feb. 1991)

-continued from No. 95 of this publication-

Table (3)-2: Strontium-90 and Cesium-137 in Milk

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	Bq/ℓ	Bq/gCa	Bq/ℓ	Bq/gK
May, 1990							
Takamiya-machi, HIROSHIMA	6.82	1.03	1.45	0.012 ± 0.0066	0.011 ± 0.0064	0.0044 ± 0.0045	0.0030 ± 0.0031
Fukuma-machi, FUKUOKA	7.03	0.949	1.69	0.0058 ± 0.0072	0.0061 ± 0.0076	0.048 ± 0.0067	0.028 ± 0.0040
August, 1990							
Hikawa-machi, SHIMANE	7.59	1.10	1.56	0.079 ± 0.0092	0.072 ± 0.0084	0.090 ± 0.0087	0.057 ± 0.0056
November, 1990							
Hokudainoujou, HOKKAIDO	7.43	1.19	1.71	0.056 ± 0.0088	0.047 ± 0.0074	0.28 ± 0.014	0.16 ± 0.008
Hikawa-machi, SHIMANE	7.21	1.05	1.59	0.079 ± 0.0093	0.075 ± 0.0089	0.059 ± 0.0077	0.037 ± 0.0048
Kajiki-machi, KAGOSHIMA	7.46	1.14	1.54	0.022 ± 0.0072	0.019 ± 0.0063	0.041 ± 0.0067	0.026 ± 0.0044
January, 1991							
Nose-machi, OSAKA	7.63	1.13	1.54	0.021 ± 0.0065	0.019 ± 0.0057	0.030 ± 0.0062	0.020 ± 0.0040
Takamiya-machi, HIROSHIMA	6.98	1.03	1.44	0.018 ± 0.0062	0.018 ± 0.0060	0.0074 ± 0.0051	0.0051 ± 0.0035
February, 1991							
Hokudainoujou, HOKKAIDO	7.51	1.18	1.51	0.052 ± 0.0092	0.044 ± 0.0078	0.13 ± 0.010	0.085 ± 0.0068
Hachijo-Island, TOKYO	7.07	1.04	1.41	0.060 ± 0.0091	0.058 ± 0.0088	0.26 ± 0.013	0.19 ± 0.010
Nishikawa-machi, NIIGATA	7.81	1.08	1.63	0.044 ± 0.0095	0.041 ± 0.0088	0.39 ± 0.018	0.24 ± 0.011
Katsuyama, FUKUI	7.50	1.24	1.55	0.034 ± 0.0085	0.028 ± 0.0069	0.027 ± 0.0067	0.018 ± 0.0043
Hikawa-machi, SHIMANE	6.83	0.884	1.63	0.029 ± 0.0072	0.033 ± 0.0081	0.038 ± 0.0064	0.023 ± 0.0040
Kochi, KOCHI	7.61	1.16	1.44	0.058 ± 0.0091	0.050 ± 0.0078	0.020 ± 0.0053	0.014 ± 0.0037
Yasu-machi, FUKUOKA	7.54	1.14	1.55	0.029 ± 0.0041	0.025 ± 0.0036	0.017 ± 0.0069	0.011 ± 0.0044
Kajiki-machi, KAGOSHIMA	7.63	1.18	1.60	0.025 ± 0.0069	0.021 ± 0.0059	0.037 ± 0.0071	0.023 ± 0.0045

(3)-3 Strontium-90 and Cesium-137 in Milk(consuming districts)  
(from Aug. 1990 to Mar. 1991)

-continued from No. 95 of this publication-

Table (3)-3: Strontium-90 and Cesium-137 in Milk

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(g/ℓ)	Ca(g/ℓ)	K(g/ℓ)	Bq/ℓ	Bq/gCa	Bq/ℓ	Bq/gK
August, 1990							
Matsue, SHIMANE	7.14	1.03	1.47	0.025 ± 0.0067	0.024 ± 0.0064	0.049 ± 0.0068	0.033 ± 0.0046
December, 1990							
Yonagusuku-mura, OKINAWA	7.02	1.06	1.51	0.020 ± 0.0069	0.019 ± 0.0065	0.011 ± 0.0048	0.0075 ± 0.0032
January, 1991							
Osaka, OSAKA	7.28	1.10	1.52	0.032 ± 0.0067	0.029 ± 0.0061	0.013 ± 0.0050	0.0084 ± 0.0033
Shinguu, WAKAYAMA	6.22	0.975	1.29	0.016 ± 0.0059	0.017 ± 0.0061	0.017 ± 0.0047	0.014 ± 0.0036
Hiroshima, HIROSHIMA	6.93	1.02	1.45	0.030 ± 0.0071	0.029 ± 0.0069	0.011 ± 0.0048	0.0079 ± 0.0033
February, 1991							
Sapporo, HOKKAIDO	6.91	1.06	1.43	0.046 ± 0.0089	0.043 ± 0.0084	0.090 ± 0.0082	0.063 ± 0.0058
Yamagata, YAMAGATA	7.27	1.10	1.41	0.040 ± 0.0084	0.036 ± 0.0077	0.023 ± 0.0055	0.016 ± 0.0039
Shinjuku, TOKYO	7.25	1.06	1.55	0.025 ± 0.0078	0.023 ± 0.0073	0.028 ± 0.0057	0.018 ± 0.0037
Niigata, NIIGATA	8.08	1.17	1.60	0.045 ± 0.0096	0.039 ± 0.0082	0.13 ± 0.011	0.079 ± 0.0069
Nagano, NAGANO	6.87	1.06	1.42	0.044 ± 0.0089	0.041 ± 0.0084	0.018 ± 0.0064	0.012 ± 0.0045
Shizuoka, SHIZUOKA	7.11	1.08	1.54	0.040 ± 0.0075	0.037 ± 0.0070	0.040 ± 0.0064	0.026 ± 0.0042
Nagoya, AICHI	7.40	1.08	1.53	0.031 ± 0.0095	0.029 ± 0.0088	0.013 ± 0.0058	0.0088 ± 0.0038
Yonago, TOTTORI	7.34	1.11	1.55	0.024 ± 0.0041	0.022 ± 0.0037	0.023 ± 0.0050	0.015 ± 0.0032
Matsue, SHIMANE	7.23	1.08	1.49	0.038 ± 0.0079	0.035 ± 0.0073	0.016 ± 0.0056	0.011 ± 0.0037
Okayama, OKAYAMA	7.08	1.07	1.52	0.030 ± 0.0039	0.028 ± 0.0036	0.0072 ± 0.0059	0.0047 ± 0.0039
Yamaguchi, YAMAGUCHI	6.88	1.04	1.39	0.034 ± 0.0040	0.033 ± 0.0038	0.0099 ± 0.0061	0.0071 ± 0.0044
Matsuyama, EHIME	7.03	1.06	1.50	0.021 ± 0.0072	0.020 ± 0.0068	0.042 ± 0.0066	0.028 ± 0.0044
Kochi, KOCHI	7.30	1.08	1.39	0.024 ± 0.0081	0.023 ± 0.0075	0.0018 ± 0.0044	0.0013 ± 0.0032
Chikushino, FUKUOKA	7.30	1.11	1.51	0.035 ± 0.0044	0.032 ± 0.0040	0.030 ± 0.0070	0.020 ± 0.0047
Nagasaki, NAGASAKI	5.34	0.801	1.16	0.015 ± 0.0034	0.019 ± 0.0043	0.011 ± 0.0037	0.0092 ± 0.0031
Kagoshima, KAGOSHIMA	7.45	1.13	1.55	0.026 ± 0.0068	0.023 ± 0.0060	0.034 ± 0.0066	0.022 ± 0.0042
March, 1991							
Fukushima, FUKUSHIMA	7.44	1.12	1.58	0.058 ± 0.0089	0.052 ± 0.0080	0.023 ± 0.0058	0.015 ± 0.0037
Yokohama, KANAGAWA	7.23	1.07	1.51	0.0071 ± 0.0067	0.0066 ± 0.0062	0.023 ± 0.0065	0.015 ± 0.0043
Fukui, FUKUI	7.11	1.07	1.53	0.027 ± 0.0079	0.025 ± 0.0073	0.021 ± 0.0062	0.014 ± 0.0040

(3)-4 Strontium-90 and Cesium-137 in Milk(powderd milk)

-continued from No. 95 of this publication-

Table (3)-4: Strontium-90 and Cesium-137 in Milk

Market Milk	Component			<sup>90</sup> Sr		<sup>137</sup> Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet		Bq/gK	
December, 1990									
Sample A	8.07	12.5	17.5	0.54 ± 0.032	0.044 ± 0.0026	0.67	± 0.032	0.038	± 0.0018
Sample B	2.52	3.38	5.92	0.088 ± 0.0088	0.026 ± 0.0026	0.42	± 0.016	0.071	± 0.0027
Sample C	2.46	3.86	5.29	0.093 ± 0.0095	0.024 ± 0.0025	0.56	± 0.018	0.11	± 0.003
Sample D	7.92	12.2	17.3	0.66 ± 0.036	0.054 ± 0.0030	2.3	± 0.06	0.13	± 0.003
Sample E	2.58	3.84	5.34	0.088 ± 0.0092	0.023 ± 0.0024	0.37	± 0.015	0.068	± 0.0028
Sample F	2.49	3.69	5.28	0.068 ± 0.0085	0.019 ± 0.0023	0.37	± 0.015	0.071	± 0.0028

\*Skim milk

(4)-1 Strontium-90 and Cesium-137 in Vegetables (producing districts)  
(from May. 1990 to Mar. 1991)

-continued from No. 95 of this publication-

Table (4)-1: Strontium-90 and Cesium-137 in Vegetables

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
(Japanese radish)							
May, 1990							
Tahara-machi, AICHI	0.617	0.177	2.54	0.043 ± 0.0069	0.24 ± 0.039	0.0079 ± 0.0067	0.0031 ± 0.0026
Kasai, HYOGO	0.831	0.146	3.35	0.056 ± 0.0076	0.39 ± 0.052	0.0000 ± 0.0075	0.0000 ± 0.0022
July, 1990							
Oota, SHIMANE	0.642	0.204	2.63	0.47 ± 0.019	2.3 ± 0.09	0.12 ± 0.009	0.046 ± 0.0035
August, 1990							
Ishikari-machi, HOKKAIDO	0.585	0.221	2.52	0.27 ± 0.009	1.2 ± 0.04	0.025 ± 0.0049	0.0099 ± 0.0019
October, 1990							
Tamayama-mura, IWATE	0.522	0.291	2.07	0.051 ± 0.0047	0.18 ± 0.016	0.0043 ± 0.0039	0.0021 ± 0.0019
Takamatsu, KAGAWA	0.550	0.241	2.25	0.012 ± 0.0056	0.051 ± 0.023	0.0041 ± 0.0034	0.0018 ± 0.0015
November, 1990							
Sannohe-machi, AOMORI	0.514	0.264	2.15	0.15 ± 0.008	0.58 ± 0.030	0.014 ± 0.0049	0.0065 ± 0.0023
Fukushima, FUKUSHIMA	0.654	0.284	2.55	0.057 ± 0.010	0.20 ± 0.036	0.0012 ± 0.0046	0.0005 ± 0.0018
Chiba, CHIBA	0.500	0.292	1.95	0.14 ± 0.009	0.48 ± 0.033	0.0039 ± 0.0052	0.0020 ± 0.0027
Utsunomiya, TOCHIGI	0.674	0.293	2.84	0.31 ± 0.011	1.1 ± 0.04	0.0024 ± 0.0041	0.0009 ± 0.0014
Kosugi-machi, TOYAMA	0.412	0.251	1.52	0.54 ± 0.016	2.2 ± 0.06	0.012 ± 0.0048	0.0078 ± 0.0032
Kanazu-machi, FUKUI	0.480	0.160	2.12	0.084 ± 0.013	0.53 ± 0.080	0.0000 ± 0.0056	0.0000 ± 0.0026
Saku, NAGANO	0.403	0.221	1.47	0.014 ± 0.0035	0.063 ± 0.016	0.0000 ± 0.0037	0.0000 ± 0.0025
Gotenba, SHIZUOKA	0.561	0.252	2.36	0.17 ± 0.010	0.69 ± 0.040	0.074 ± 0.0082	0.031 ± 0.0035
Gifu, Gifu	0.440	0.206	1.58	0.050 ± 0.0099	0.24 ± 0.048	0.0027 ± 0.0046	0.0017 ± 0.0029
Adogawa-machi, SHIGA	0.395	0.128	1.52	0.20 ± 0.016	1.6 ± 0.12	0.0050 ± 0.0052	0.0033 ± 0.0034
Shime-machi, FUKUOKA	0.535	0.215	2.61	0.027 ± 0.0063	0.13 ± 0.029	0.0041 ± 0.0036	0.0016 ± 0.0014
Koushi-machi, KUMAMOTO	0.569	0.390	2.04	0.11 ± 0.010	0.28 ± 0.026	0.0000 ± 0.0036	0.0000 ± 0.0017
December, 1990							
Maebashi, GUNMA	0.540	0.309	2.11	0.041 ± 0.0077	0.13 ± 0.025	0.0033 ± 0.0034	0.0016 ± 0.0016
Takane-machi, YAMANASHI	0.401	0.248	1.46	0.16 ± 0.013	0.66 ± 0.054	0.0000 ± 0.0045	0.0000 ± 0.0031
Meiwa-machi, MIE	0.552	0.141	2.51	0.031 ± 0.068	0.22 ± 0.048	0.015 ± 0.0045	0.0058 ± 0.0018
Kashihara, NARA	0.522	0.166	2.21	0.093 ± 0.0078	0.56 ± 0.047	0.0048 ± 0.0044	0.0022 ± 0.0020
Kokufu-machi, TOTTORI	0.475	0.208	2.01	0.13 ± 0.010	0.61 ± 0.046	0.0000 ± 0.0030	0.0000 ± 0.0015
Hiroshima, HIROSHIMA	0.515	0.188	2.00	0.017 ± 0.0095	0.089 ± 0.051	0.017 ± 0.0059	0.0087 ± 0.0030



Location	Component			<sup>90</sup> Sr				<sup>137</sup> Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet		Bq/gCa		Bq/kgwet		Bq/gK	
Yuya-machi, YAMAGUCHI	0.486	0.197	1.65	0.16	± 0.010	0.79	± 0.052	0.010	± 0.0035	0.0060	± 0.0021
Kubokawa-machi, KOCHI	0.670	0.169	2.90	0.13	± 0.009	0.76	± 0.055	0.021	± 0.0057	0.0074	± 0.0020
Ishii-machi, TOKUSHIMA	0.594	0.238	2.56	0.046	± 0.0087	0.19	± 0.037	0.0003	± 0.0035	0.0001	± 0.0014
Usa, OOLTA	0.662	0.179	2.59	0.082	± 0.011	0.46	± 0.063	0.0049	± 0.0045	0.0019	± 0.0017
Takanabe-machi, MIYAZAKI	0.605	0.261	2.58	0.20	± 0.008	0.77	± 0.032	0.0062	± 0.0066	0.024	± 0.0025
Kaimon-machi, KAGOSHIMA	0.557	0.221	2.28	0.13	± 0.007	0.58	± 0.033	0.023	± 0.0050	0.010	± 0.0022
January, 1991 Shinguu, WAKAYAMA	0.521	0.134	2.10	0.039	± 0.0067	0.29	± 0.050	0.0000	± 0.0029	0.0000	± 0.0014
(Onion)											
July, 1990 Kumatori-machi, OSAKA	0.328	0.129	1.26	0.061	± 0.0068	0.47	± 0.053	0.021	± 0.0061	0.017	± 0.0048
(Cabbage)											
July, 1990 Oota, SHIMANE	1.19	1.67	3.19	1.5	± 0.03	0.91	± 0.018	0.23	± 0.012	0.073	± 0.0036
November, 1990 Sannohe-machi, AOMORI	0.532	0.401	2.01	0.20	± 0.008	0.50	± 0.021	0.10	± 0.008	0.051	± 0.0040
January, 1991 Kumatori-machi, OSAKA	0.621	0.371	1.87	0.023	± 0.0069	0.061	± 0.019	0.0017	± 0.0039	0.0009	± 0.0021
(Spinach)											
May, 1990 Tahara-machi, AICHI	1.21	0.420	4.95	0.033	± 0.0061	0.078	± 0.014	0.017	± 0.0077	0.0034	± 0.0016
August, 1990 Ishikari-machi, HOKKAIDO	1.32	0.400	5.40	0.10	± 0.009	0.25	± 0.024	0.015	± 0.0090	0.0027	± 0.0017
October, 1990 Toyama, TOYAMA	1.67	0.970	6.20	0.53	± 0.019	0.54	± 0.020	0.027	± 0.0087	0.0044	± 0.0014
Takamatsu, KAGAWA	1.69	0.717	6.92	0.054	± 0.0078	0.075	± 0.011	0.0084	± 0.0064	0.0012	± 0.00092
November, 1990 Fukushima, FUKUSHIMA	1.82	1.45	5.93	0.40	± 0.022	0.27	± 0.015	0.0095	± 0.0066	0.0016	± 0.0011
Chiba, CHIBA	1.70	0.568	7.23	0.22	± 0.010	0.38	± 0.018	0.021	± 0.0059	0.0029	± 0.00082
Fukui, FUKUI	1.94	0.488	8.74	0.049	± 0.011	0.10	± 0.023	0.0052	± 0.0058	0.00060	± 0.00067
Saku, NAGANO	2.23	0.729	9.50	0.058	± 0.012	0.079	± 0.016	0.13	± 0.012	0.013	± 0.0012
Gotenba, SHIZUOKA	1.40	0.502	5.71	0.067	± 0.0072	0.13	± 0.014	0.19	± 0.012	0.033	± 0.0022
Gifu, GUFU	1.54	0.597	5.86	0.11	± 0.013	0.18	± 0.021	0.0025	± 0.0042	0.00043	± 0.00071

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
Kusu-machi, MIE	1.79	0.945	7.08	0.044 ± 0.0098	0.046 ± 0.010	0.011 ± 0.0059	0.0016 ± 0.00084
Kasai, HYOGO	1.39	0.336	6.08	0.080 ± 0.0080	0.24 ± 0.024	0.0000 ± 0.0044	0.0000 ± 0.00072
Kurayoshi, TOTTORI	1.70	0.681	6.02	0.099 ± 0.014	0.15 ± 0.020	0.15 ± 0.013	0.025 ± 0.0022
Matsuyama, EHIME	1.66	0.844	6.02	0.052 ± 0.0063	0.062 ± 0.0074	0.015 ± 0.0060	0.0024 ± 0.0010
Shime-machi, FUKUOKA	1.82	1.12	7.06	0.081 ± 0.0074	0.072 ± 0.0066	0.0076 ± 0.0052	0.0011 ± 0.00074
Koushi-machi, KUMAMOTO	1.75	0.561	7.12	0.045 ± 0.013	0.081 ± 0.024	0.028 ± 0.0077	0.0039 ± 0.0011
December, 1990							
Maebashi, GUNMA	1.64	1.49	5.51	0.081 ± 0.0066	0.054 ± 0.0044	0.084 ± 0.0092	0.015 ± 0.0017
Takane-machi, YAMANASHI	1.80	0.661	7.63	0.050 ± 0.0089	0.076 ± 0.013	0.0089 ± 0.0065	0.0012 ± 0.00086
Ritsuto-machi, SHIGA	1.49	0.678	6.06	0.072 ± 0.012	0.11 ± 0.017	0.0047 ± 0.0062	0.0008 ± 0.0010
Kashihara, NARA	1.14	0.479	4.55	0.15 ± 0.015	0.31 ± 0.031	0.011 ± 0.0060	0.0024 ± 0.0013
Hiroshima, HIROSHIMA	1.31	0.694	5.03	0.041 ± 0.0096	0.060 ± 0.014	0.014 ± 0.0061	0.0027 ± 0.0012
Yuya-machi, YAMAGUCHI	1.53	0.473	6.33	0.23 ± 0.015	0.49 ± 0.032	0.027 ± 0.0063	0.0043 ± 0.0010
Kubokawa-machi, KOCHI	1.55	1.54	4.87	0.12 ± 0.012	0.081 ± 0.0081	0.22 ± 0.014	0.045 ± 0.0028
Usa, OOTA	2.14	0.507	9.12	0.11 ± 0.013	0.21 ± 0.026	0.0097 ± 0.0059	0.0011 ± 0.00064
Takanabe-machi, MIYAZAKI	1.55	1.07	5.10	0.30 ± 0.013	0.28 ± 0.012	0.34 ± 0.016	0.067 ± 0.0031
February, 1991							
Kaimon-machi, KAGOSHIMA	1.62	0.962	4.96	0.29 ± 0.013	0.30 ± 0.014	0.54 ± 0.022	0.11 ± 0.004
March, 1991							
Ishii-machi, TOKUSHIMA	1.42	0.575	5.34	0.042 ± 0.0058	0.073 ± 0.010	0.0000 ± 0.0061	0.0000 ± 0.0011
(Chinese cabbage)							
October, 1990							
Tamayama-mura, IWATE	0.542	0.437	2.12	0.17 ± 0.016	0.40 ± 0.036	0.031 ± 0.0062	0.015 ± 0.0029
November, 1990							
Utsunomiya, TOCHIGI	0.683	0.717	2.19	0.087 ± 0.0080	0.12 ± 0.011	0.011 ± 0.0057	0.0051 ± 0.0026
January, 1991							
Shinguu, WAKAYAMA	0.569	0.299	2.09	0.14 ± 0.011	0.46 ± 0.037	0.027 ± 0.0051	0.013 ± 0.0024

(4)-2 Strontium-90 and Cesium-137 in Vegetables (consuming districts)  
(from May, 1990 to Mar. 1991)

-continued from No. 95 of this publication-

Table (4)-2: Strontium-90 and Cesium-137 in Vegetables

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
(Japanese radish)							
September, 1990 Sendai, MIYAGI	0.546	0.211	2.48	0.048 ± 0.0074	0.23 ± 0.035	0.0051 ± 0.0048	0.0021 ± 0.0019
October, 1990 Yamagata, YAMAGATA	0.434	0.353	1.60	0.033 ± 0.0098	0.095 ± 0.028	0.0040 ± 0.0044	0.0025 ± 0.0027
Kyoto, KYOTO	0.518	0.153	2.13	0.098 ± 0.012	0.64 ± 0.080	0.052 ± 0.0079	0.025 ± 0.0037
November, 1990 Akita, AKITA	0.519	0.287	2.02	0.34 ± 0.022	1.2 ± 0.08	0.061 ± 0.0080	0.030 ± 0.0040
Shinjuku, TOKYO	0.612	0.345	2.48	0.12 ± 0.007	0.35 ± 0.020	0.012 ± 0.0043	0.0049 ± 0.0017
Niigata, NIIGATA	0.413	0.203	1.49	0.0047 ± 0.0091	0.023 ± 0.045	0.0055 ± 0.0043	0.0037 ± 0.0029
Kanazawa, ISHIKAWA	0.542	0.257	2.22	0.096 ± 0.0096	0.37 ± 0.037	0.0000 ± 0.0034	0.0000 ± 0.0015
Osaka, OSAKA	0.472	0.150	2.07	0.026 ± 0.0065	0.17 ± 0.043	0.048 ± 0.0059	0.023 ± 0.0029
Okayama, OKAYAMA	0.428	0.204	1.51	0.30 ± 0.019	1.5 ± 0.10	0.011 ± 0.0051	0.0074 ± 0.0034
Saga, SAGA	0.544	0.183	2.45	0.094 ± 0.0088	0.51 ± 0.048	0.0000 ± 0.0040	0.0000 ± 0.0017
Yonagusuku-mura, OKINAWA	0.718	0.284	2.63	0.070 ± 0.010	0.25 ± 0.035	0.0000 ± 0.0027	0.0000 ± 0.0010
January, 1991 Nagasaki, NAGASAKI	0.509	0.182	2.28	0.091 ± 0.0097	0.50 ± 0.053	0.016 ± 0.0041	0.0070 ± 0.0018
March, 1991 Yokohama, KANAGAWA	0.468	0.250	1.65	0.051 ± 0.0069	0.20 ± 0.028	0.0018 ± 0.0023	0.0011 ± 0.0014
(Cabbage)							
November, 1990 Akita, AKITA	0.513	0.563	1.72	0.20 ± 0.017	0.36 ± 0.030	0.064 ± 0.0079	0.037 ± 0.0046
(Spinach)							
May, 1990 Sendai, MIYAGI	1.55	0.491	6.04	0.060 ± 0.0063	0.12 ± 0.013	0.013 ± 0.0056	0.0021 ± 0.00092
June, 1990 Niigata, NIIGATA	1.79	1.53	7.29	0.067 ± 0.0068	0.044 ± 0.0044	0.025 ± 0.0065	0.0034 ± 0.00089
October, 1990 Yamagata, YAMAGATA	1.48	0.437	6.25	0.087 ± 0.0089	0.20 ± 0.020	0.0079 ± 0.0065	0.0013 ± 0.0010

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
November, 1990							
Shinjuku, TOKYO	1.61	0.456	6.95	0.19 ± 0.019	0.41 ± 0.041	0.0091 ± 0.0065	0.0013 ± 0.00093
Kanazawa, ISHIKAWA	1.47	0.520	6.03	0.052 ± 0.011	0.10 ± 0.021	0.028 ± 0.0076	0.0047 ± 0.0013
Kyoto, KYOTO	1.91	1.08	5.39	0.13 ± 0.015	0.12 ± 0.013	0.018 ± 0.0073	0.0032 ± 0.0014
Osaka, OSAKA	1.80	0.597	7.82	0.13 ± 0.015	0.22 ± 0.025	0.022 ± 0.0068	0.0029 ± 0.00086
Okayama, OKAYAMA	1.62	0.502	6.44	0.050 ± 0.011	0.10 ± 0.023	0.0000 ± 0.0036	0.00000 ± 0.00055
Matsuyama, EHIME	1.71	0.630	6.66	0.076 ± 0.0069	0.12 ± 0.011	0.014 ± 0.0062	0.0022 ± 0.00093
Saga, SAGA	1.70	1.04	7.03	0.064 ± 0.0067	0.061 ± 0.0064	0.011 ± 0.0049	0.0016 ± 0.00070
December, 1990							
Yonagusuku-mura, OKINAWA	1.72	0.612	7.14	0.029 ± 0.0046	0.047 ± 0.0076	0.013 ± 0.0057	0.0018 ± 0.00080
January, 1991							
Nagasaki, NAGASAKI	1.42	0.568	5.55	0.062 ± 0.012	0.11 ± 0.022	0.0093 ± 0.0068	0.0017 ± 0.0012
March, 1991							
Yokohama, KANAGAWA	1.54	0.870	4.89	0.17 ± 0.010	0.19 ± 0.011	0.0097 ± 0.0060	0.0020 ± 0.0012

(5) Strontium-90 and Cesium-137 in Tea(Japanese Tea)  
(from May. 1990 to Jun. 1990)

-continued from No. 95 of this publication-

Table (5): Strontium-90 and Cesium-137 in Tea(Japanese Tea)

Location	Component			<sup>90</sup> Sr				<sup>137</sup> Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kg		Bq/gCa		Bq/kg		Bq/gK	
May, 1990											
Shirakawa-machi, GIFU	4.64	1.88	16.6	0.55	± 0.074	0.29	± 0.040	0.44	± 0.052	0.027	± 0.0032
Mihune-machi, KUMAMOTO	4.70	2.99	16.6	0.35	± 0.065	0.12	± 0.022	0.32	± 0.049	0.019	± 0.0029
Ue-mura, KUMAMOTO	5.33	2.62	19.1	1.1	± 0.10	0.44	± 0.039	0.67	± 0.068	0.035	± 0.0036
June, 1990											
Nara, NARA	5.17	2.63	16.7	0.47	± 0.074	0.18	± 0.028	0.23	± 0.044	0.014	± 0.0027
Nara, NARA	5.39	2.52	18.1	0.67	± 0.091	0.27	± 0.036	0.80	± 0.069	0.044	± 0.0038
Miyanojo-machi, KAGOSHIMA	5.68	2.90	20.0	1.0	± 0.10	0.35	± 0.034	1.2	± 0.09	0.058	± 0.0043
Chiran-machi, KAGOSHIMA	4.84	2.31	17.7	0.28	± 0.067	0.12	± 0.029	1.8	± 0.10	0.10	± 0.006

(6) Strontium-90 and Cesium-137 in Sea Fish  
(from Apr. 1990 to Feb. 1991)

-continued from No. 95 of this publication-

Table (6): Strontium-90 and Cesium-137 in Sea Fish

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK	
(Hexagrammos otakii) August, 1990 Souma, FUKUSHIMA	1.66	1.99	3.79	0.0080±0.0032	0.0040±0.0016	0.16 ±0.011	0.042 ±0.0030	
(Trachurus japonicus) November, 1990 Shizuoka, SHIZUOKA	3.51	7.53	3.22	0.018 ±0.0039	0.0023±0.00051	0.21 ±0.015	0.065±0.0047	
December, 1990 Odawara, KANAGAWA	1.55	0.292	4.80	0.0000±0.0092	0.000 ±0.032	0.25 ±0.016	0.052 ±0.0033	
January, 1991 Shinguu, WAKAYAMA	2.62	5.35	2.38	0.0000±0.0080	0.0000±0.0015	0.12 ±0.012	0.051 ±0.0049	
(Branchiostegus sp.) November, 1990 Nagasaki, NAGASAKI	1.01	0.411	2.93	0.007 ±0.013	0.017 ±0.032	0.15 ±0.015	0.052 ±0.0050	
(Sardinops melanosticta) September, 1990 Yamagata, YAMAGATA	3.00	6.42	2.40	0.0097±0.0088	0.0015±0.0014	0.081 ±0.011	0.034 ±0.0045	
January, 1991 Nagano, NAGANO	3.07	7.46	2.73	0.0006±0.0096	0.0001±0.0013	0.095 ±0.011	0.035 ±0.0040	
(Sebastiscus marmoratus) April, 1990 Hamada, SHIMANE	5.61	14.4	2.01	0.029 ±0.0080	0.0020±0.00056	0.14 ±0.012	0.068 ±0.0062	
(Katsuwonus pelamis) June, 1990 Tosa, KOCHI	1.15	0.119	3.48	0.0048±0.0086	0.040 ±0.073	0.43 ±0.018	0.12 ±0.005	
(Limanda herzensteini) June, 1990 Sendai, NIYAGI	3.28	6.73	3.40	0.0075±0.0038	0.0011±0.00056	0.085 ±0.012	0.025 ±0.0036	
November, 1990 Mutsu, AOMORI	1.47	0.982	3.84	0.0000±0.0036	0.0000±0.0037	0.17 ±0.014	0.044 ±0.0036	

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs		
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK	
Iwamuro-mura, NIIGATA	1.47	0.687	3.90	0.0076±0.0063	0.011 ±0.0092	0.13 ±0.019	0.034 ±0.0049	
Mikuni-machi, FUKUI	1.52	2.00	3.09	0.011 ±0.0054	0.0054±0.0027	0.11 ±0.017	0.037 ±0.0054	
Aji-machi, KAGAWA	1.46	0.245	4.37	0.0031±0.0036	0.013 ±0.015	0.15 ±0.013	0.035 ±0.0030	
February, 1991								
Ootake, HIROSHIMA	2.88	6.19	3.76	0.014 ±0.0090	0.0023±0.0015	0.13 ±0.012	0.035 ±0.0032	
(Sillago sihama)								
June, 1990								
Minamichita-machi, AICHI	3.75	9.07	2.67	0.014 ±0.0097	0.0015±0.0011	0.099 ±0.012	0.037 ±0.0045	
(Spratelloides gracilis)								
December, 1990								
Akune, KAGOSHIMA	3.02	5.90	3.13	0.0077±0.0039	0.0013±0.00066	0.17 ±0.014	0.055 ±0.0043	
(Oncorhynchus keta)								
September, 1990								
Urakawa-machi, HOKKAIDO	1.32	0.573	3.71	0.0026±0.0034	0.0046±0.0059	0.31 ±0.016	0.084 ±0.0044	
(Scomber japonicus)								
August, 1990								
Matsuyama, EHIME	1.30	0.352	3.99	0.0075±0.0054	0.021 ±0.015	0.21 ±0.017	0.053 ±0.0043	
November, 1990								
Kyoto, KYOTO	1.12	0.191	3.16	0.0027±0.0046	0.014 ±0.024	0.23 ±0.018	0.073 ±0.0057	
Osaka, OSAKA	1.31	0.189	4.10	0.0041±0.0039	0.022 ±0.021	0.20 ±0.015	0.050 ±0.0037	
December, 1990								
Sakaiminato, TOTTORI	1.34	0.492	3.58	0.010 ±0.0033	0.021 ±0.0068	0.14 ±0.011	0.040 ±0.0031	
(Chrysophrys major)								
July, 1990								
Oga, AKITA	5.66	12.7	3.25	0.066 ±0.0064	0.0051±0.00050	0.21 ±0.017	0.065 ±0.0051	
Fukuoka, FUKUOKA	1.46	0.420	4.50	0.0000±0.0086	0.000 ±0.020	0.26 ±0.016	0.057 ±0.0035	
(Caesio chrysozonus cuvier)								
November, 1990								
Yonagusuku-mura, OKINAWA	3.79	9.24	4.16	0.013 ±0.0039	0.0014±0.00042	0.24 ±0.016	0.059 ±0.0037	
(Seriola quinqueradiata)								
October, 1990								
Togi-machi, ISHIKAWA	1.40	0.953	3.73	0.0062±0.0051	0.0065±0.0054	0.24 ±0.019	0.065 ±0.0052	
(Mugil cephalus)								
August, 1990								
Morodomi-machi, SAGA	1.17	0.399	3.64	0.0047±0.0030	0.012 ±0.0075	0.12 ±0.011	0.034 ±0.0030	

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
November, 1990 Ushimado-machi, OKAYAMA	1.31	0.489	3.89	0.0038±0.0089	0.008 ±0.018	0.15 ±0.012	0.038 ±0.0032
(Decapterus muroadsi)							
October, 1990 Miyake-Island, TOKYO	1.36	1.27	3.12	0.0091±0.0042	0.0071±0.0033	0.16 ±0.014	0.050 ±0.0046
(Sebastes inermis)							
January, 1991 Yamaguchi, YAMAGUCHI	4.95	12.8	3.40	0.028 ±0.0053	0.0022±0.00041	0.17 ±0.014	0.050 ±0.0041
(Ammodytes personatus Girard)							
April, 1990 Akashi, HYOGO	2.48	3.89	3.82	0.0015±0.0083	0.0004±0.0021	0.067 ±0.011	0.018 ±0.0028



# Sea Fish

Japanese name	English name	Scientific name
Ainame	Greenling	Hexagrammos otakii
Aji	Horse mackerel	Trachurus japonicus
Amadai	Tilefish	Branchiostegus sp.
Iwashi	Sardine	Sardinops melanosticta
Kasago	Scorpion-fish	Sebastiscus marmoratus
Katsuo	Bonito	Katsuwonus pelamis
Karei	Flatfish	Limanda herzensteini
Kisu	Sillago	Sillago sihama
Kibinago	Banded blue-sprat	Spratelloides gracilis
Sake	Salmon	Oncorhynchus keta
Saba	Common mackerel	Scomber japonicus
Tai	Sea bream	Chrysophrys major
Takasago	Black-tipped fusilier	Caesio chrysozonus cuvier
Fukuragi	Yellow-tail	Seriola quinqueradiata
Bora	Gray mullet	Mugil cephalus
Muroaji	Horse-scad mackerel	Decapterus muroadsi
Webaru	Black Rockfish	Sebastes inermis
Ikanago	Sando lance	Ammodytes personatus Girard

(7) Strontium-90 and Cesium-137 in Freshwater Fish  
(from Jul. 1990 to Dec. 1990)

-continued from No. 95 of this publication-

Table (7): Strontium-90 and Cesium-137 in Freshwater Fish

Location	Component			<sup>90</sup> Sr				<sup>137</sup> Cs			
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet		Bq/gCa		Bq/kgwet		Bq/gK	
(Cyprinus carpio)											
August, 1990											
Akita, AKITA	3.37	8.41	2.56	2.3	± 0.04	0.28	± 0.004	0.59	± 0.026	0.23	± 0.010
September, 1990											
Fukushima, FUKUSHIMA	4.07	11.9	2.71	0.96	± 0.025	0.081	± 0.0021	0.18	± 0.015	0.065	± 0.0056
November, 1990											
Shobara, HIRISHIMA	1.04	0.511	2.74	0.093	± 0.010	0.18	± 0.020	0.15	± 0.016	0.053	± 0.0057
(Carassius auratus)											
July, 1990											
Barato-lake, HOKKAIDO	4.28	9.93	2.79	0.0084±0.0049		0.00085±0.00049		0.088 ± 0.010		0.031 ± 0.0037	
November, 1990											
Toyanogata, NIIGATA	1.15	0.459	3.19	0.068	± 0.0083	0.15	± 0.018	0.16	± 0.015	0.050	± 0.0047
December, 1990											
Mikata-lake, FUKUI	3.07	6.57	2.64	0.81	± 0.020	0.12	± 0.003	0.27	± 0.018	0.10	± 0.007
Uji, KYOTO	4.69	10.6	2.64	1.1	± 0.02	0.11	± 0.002	0.052	± 0.0093	0.020	± 0.0035
(Hypomesus transpacificus nipponensis)											
December, 1990											
Suwa-lake, NAGANO	2.27	7.75	2.24	0.089	± 0.0083	0.011	± 0.0011	0.15	± 0.014	0.069	± 0.0062

Freshwater Fish

Japanese name	English name	Scientific name
Koi	Carp	Cyprinus carpio
Funa	A crucian carp	Carassius auratus
Wakasagi	Pond-smelt	Hypomesus transpacificus nipponensis

(8) Strontium-90 and Cesium-137 in Shellfish  
(from May. 1990 to Feb. 1991)

-continued from No. 95 of this publication-

Table (8): Strontium-90 and Cesium-137 in Shellfish

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
(Venerupis philipinarum)							
May, 1990							
Takaki-machi, NAGASAKI	2.35	1.57	2.76	0.000 ± 0.022	0.000 ± 0.014	0.037 ± 0.020	0.013 ± 0.0072
(Saxidomus purpuratus)							
June, 1990							
Minamichita-machi, AICHI	1.50	0.328	2.52	0.000 ± 0.017	0.000 ± 0.052	0.022 ± 0.016	0.0087 ± 0.0063
(Ostrea gigas)							
February, 1991							
Hatsukaichi, HIROSHIMA	1.64	0.635	2.43	0.019 ± 0.014	0.031 ± 0.022	0.055 ± 0.012	0.023 ± 0.0048
(Turbo cornutus)							
May, 1990							
Ryotsu, NIIGATA	2.04	0.561	3.20	0.025 ± 0.020	0.045 ± 0.036	0.047 ± 0.018	0.015 ± 0.0055
June, 1990							
Sakata, YAMAGATA	2.69	2.09	2.46	0.0000 ± 0.0080	0.0000 ± 0.0038	0.049 ± 0.0096	0.020 ± 0.0039
September, 1990							
Togi-machi, ISHIKAWA	2.83	1.53	1.67	0.0041 ± 0.0057	0.0027 ± 0.0037	0.053 ± 0.011	0.032 ± 0.0067
(Pecten Yessoensis)							
November, 1990							
Mutsu, AOMORI	1.32	0.255	2.21	0.0033 ± 0.0036	0.013 ± 0.014	0.053 ± 0.0085	0.024 ± 0.0038
February, 1991							
Yamada-machi, IWATE	2.19	0.304	3.12	0.000 ± 0.010	0.000 ± 0.034	0.045 ± 0.0085	0.015 ± 0.0027
(Mytilus edulis)							
June, 1990							
Mutsu, AOMORI	2.23	0.437	2.26	0.0000 ± 0.0045	0.000 ± 0.010	0.035 ± 0.0094	0.016 ± 0.0041

## Shellfish

Japanese name	English name	Scientific name
Asari	Short-necked clam	Venerupis phillipinarum
Ohasari		Saxidomus purpuratus
Kaki	Oyster	Ostrea gigas
Sazae	Wreath shell	Turbo cornutus
Hotategai	Scallop	Pecten Yessoensis
Murasakiigai	Mussel	Mytilus edulis

(9) Strontium-90 and Cesium-137 in Seaweeds  
(from Apr. 1990 to Feb. 1991)

-continued from No. 95 of this publication-

Table (9): Strontium-90 and Cesium-137 in Seaweeds

Location	Component			<sup>90</sup> Sr		<sup>137</sup> Cs	
	Ash(%)	Ca(g/kg)	K(g/kg)	Bq/kgwet	Bq/gCa	Bq/kgwet	Bq/gK
(Undaria pinnatifida)							
April, 1990							
Togi-machi, ISHIKAWA	3.68	0.779	6.34	0.0000 ± 0.0070	0.0000 ± 0.0090	0.035 ± 0.0088	0.0056 ± 0.0014
May, 1990							
Fukaura-machi, AOMORI	1.90	0.927	4.04	0.025 ± 0.0050	0.027 ± 0.0053	0.021 ± 0.0063	0.0051 ± 0.0016
Mutsu, AOMORI	1.11	0.683	1.95	0.018 ± 0.0038	0.027 ± 0.0056	0.021 ± 0.0053	0.011 ± 0.0027
Ryotsu, NIIGATA	3.88	0.997	6.29	0.030 ± 0.045	0.030 ± 0.0045	0.036 ± 0.0066	0.0057 ± 0.0011
June, 1990							
Sakata, YAMAGATA	2.99	1.77	3.45	0.045 ± 0.011	0.025 ± 0.0060	0.044 ± 0.010	0.013 ± 0.0029
February, 1991							
Minamichita-machi, AICHI	2.54	0.853	6.22	0.031 ± 0.0045	0.037 ± 0.0053	0.027 ± 0.0066	0.0044 ± 0.0011
Hiroshima, HIROSHIMA	1.46	0.749	3.23	0.025 ± 0.0089	0.034 ± 0.012	0.020 ± 0.0069	0.0063 ± 0.0021
Shimabara, NAGASAKI	2.65	0.761	10.4	0.040 ± 0.011	0.052 ± 0.014	0.030 ± 0.0067	0.0029 ± 0.00064

Seaweeds

Japanese name	English name	Scientific name
Wakame	Wakame seaweed	Undaria pinnatifida

## \* \* \* Total Diet \* \* \*




Fig. 1

\* \* \* Milk (producing districts for domestic program) \* \* \*




Fig.2-1

\* \* \* Milk (producing districts for WHO program) \* \* \*




Fig. 2-2



\* \* \* Milk (consuming districts) \* \* \*




Fig.2-3

\* \* \*    Vegetables (producing districts)    \* \* \*  
[ Spinach ]




Fig. 3-1

\* \* \* Vegetables (consuming districts) \* \* \*

[ Spinach ]




Fig.3-2

\* \* \* Sea Fish \* \* \*

[ *Scomber japonicus* ]




Fig. 4

\*\*\* Shellfish \*\*\*

( Turbo cornutus )





Fig.5

\* \* \* Seaweeds \* \* \*

[ *Undaria pinnatifida* ]

< Strontium-90 >



< Cesium-137 >





Fig.6

# \*\* Sampling Locations in Japan \*\*

- |                 |                |
|-----------------|----------------|
| 1 : Sapporo     | 23 : Tsu       |
| 2 : Aomori      | 24 : Kyoto     |
| 3 : Morioka     | 25 : Osaka     |
| 4 : Akita       | 26 : Tottori   |
| 5 : Sendai      | 27 : Kobe      |
| 6 : Yamagata    | 28 : Wakayama  |
| 7 : Fukushima   | 29 : Okayama   |
| 8 : Niigata     | 30 : Matsue    |
| 9 : Mito        | 31 : Takamatsu |
| 10 : Utsunomiya | 32 : Hiroshima |
| 11 : Chiba      | 33 : Kochi     |
| 12 : Urawa      | 34 : Matsuyama |
| 13 : Shinjuku   | 35 : Yamaguchi |
| 14 : Nagano     | 36 : Oita      |
| 15 : Yokohama   | 37 : Fukuoka   |
| 16 : Koufu      | 38 : Saga      |
| 17 : Toyama     | 39 : Miyazaki  |
| 18 : Kanazawa   | 40 : Nagasaki  |
| 19 : Shizuoka   | 41 : Kagoshima |
| 20 : Fukui      | 42 : Naha      |
| 21 : Nagoya     |                |
| 22 : Otsu       |                |



Notice for Changing of the Address and the Telephone Number

National Institute of Radiological Sciences, Japan, has changed the addresss and the telephone number as follows, from April 1992.

Address(from April 1,1992) : Anagawa 4-9-1, Inage-ku, 263 Japan

Telephone Number(from April 29,1992): 81-43-251-2111

Fax Number(from April 29,1992) : 81-43-256-9616